

**National Center for Health Statistics** 

Guide to Completing the Facility
Worksheets for the Certificate of Live
Birth and Report of Fetal Death

(2003 revision)

Updated May 2016



CDC

CONTROL OF THE PREVIOUS C

# Training for completing medical and health information for the birth certificate and report of fetal death is available online!

To access

# "Applying Best Practices for Reporting Medical and Health Information on Birth Certificates"

go to:

http://www.cdc.gov/nchs/training/BirthCertificateElearning.

## **Table of Contents**

| Instructions  | Pregnancy resulted from infertility treatment   | 18 |
|---|---|----|
| How to Use This Guide                                   | Fertility-enhancing drugs, artificial insemination, or intrauterine                         |    |
| Mother  | insemination  |    |
|   | Assisted reproductive technology  |    |
| Facility Information                                    | Mother had a previous cesarean delivery   | 19 |
| Facility name   | Infections present and/or treated during this pregnancy                                     | 20 |
| Facility ID   | Gonorrhea   |    |
| City, town, or location of birth                        | Syphilis  |    |
| County of birth   | Chlamydia   |    |
| Place where birth occurred (Birthplace)                 | Hepatitis B   |    |
| There where of the occurred (Bittiplace)                | Hepatitis C   |    |
| Prenatal Care and Pregnancy History                     | Obstetric procedures  |    |
| Date of first prenatal care visit                       | External cephalic version   | 22 |
| Total number of prenatal care visits for this pregnancy | Labor and Delivery  |    |
| Date last normal menses began                           | Date of birth   | 23 |
| Number of previous live births now living               | Time of birth   | 23 |
| Number of previous live births now dead                 | Certifier's name and title.   | 23 |
| Date of last live birth                                 | Date certified  | 23 |
| Number of other pregnancy outcomes                      | Principal source of payment   | 24 |
| Date of last other pregnancy outcome                    | Infant's medical record number  | 24 |
| Risk factors in this pregnancy                          | Was the mother transferred to this facility for maternal medical or                         |    |
| Diabetes  | fetal indications for delivery?   | 25 |
| Prepregnancy  | Attendant's name, title, and ID   | 26 |
| Gestational   | Mother's weight at delivery   | 26 |
| Hypertension  | Characteristics of labor and delivery   | 27 |
| Prepregnancy  | Induction of labor  |    |
| Gestational   | Augmentation of labor   |    |
| Eclampsia16Previous preterm births17                    | Steroids (glucocorticoids) for fetal lung maturation received by the mother before delivery |    |

#### **Table of Contents—Continued**

| Labor and Delivery—Continued  |    | Cyanotic congenital heart disease                       | 42 |
|---|----|---|----|
| Antibiotics received by the mother during delivery                  | 20 | Congenital diaphragmatic hernia                         |    |
| Clinical chorioamnionitis diagnosed during labor or maternal        | 2) | Omphalocele   |    |
| temperature $\geq 38^{\circ}\text{C} (100.4^{\circ}\text{F})$       | 29 | Gastroschisis   |    |
| Epidural or spinal anesthesia during labor                          |    | Limb reduction defect.                                  |    |
| Method of delivery  |    | Cleft lip with or without cleft palate                  |    |
| Fetal presentation at birth   |    | Cleft palate alone                                      |    |
| Final route and method of delivery                                  |    | Down syndrome   |    |
| If cesarean, was a trial of labor attempted?                        |    | Suspected chromosomal disorder                          |    |
| Maternal morbidity  | 32 | Hypospadias   |    |
| Maternal transfusion  |    | Was the infant transferred within 24 hours of delivery? |    |
| Third- or fourth-degree perineal laceration                         |    | Is the infant living at the time of the report?         |    |
| Ruptured uterus   |    | Is the infant being breastfed at discharge?             |    |
| Unplanned hysterectomy  | 33 | Method of disposition                                   |    |
| Admission to an intensive care unit                                 | 33 | iviethod of disposition                                 | 4/ |
| Newborn Information   |    | Index of Items  | 48 |
| Birthweight or weight of fetus                                      | 34 |   |    |
| Obstetric estimate of gestation at delivery                         | 34 |   |    |
| Sex of child  |    |   |    |
| Apgar score   | 37 |   |    |
| Plurality.  | 37 |   |    |
| If not a single birth, order born in the delivery                   | 37 |   |    |
| If not a single birth, number of infants in the delivery born alive | 37 |   |    |
| Abnormal conditions of the newborn                                  | 38 |   |    |
| Assisted ventilation required immediately following delivery        | 38 |   |    |
| Assisted ventilation required for more than six hours               | 39 |   |    |
| NICU admission.   | 40 |   |    |
| Newborn given surfactant replacement therapy                        | 40 |   |    |
| Antibiotics received by the newborn for suspected neonatal sepsis   | 40 |   |    |
| Seizure or serious neurologic dysfunction                           | 41 |   |    |

Anencephaly. .

## **How to Use This Guide**

This guide was developed to assist in completing the facility worksheets for the revised Certificate of Live Birth and Report of Fetal Death (birth certificate [BC], facility worksheet for the report of fetal death [FDFWS], report of fetal death [FDR]).

| Definitions   | Instructions   | Sources   | Keywords and abbreviations  |
|---|--|---|---|
| Defines the items in the order they appear on the facility worksheet. | Provides specific instructions for completing each item. | Identifies the sources in the medical records where information for each item can be found. The specific records available will differ somewhat from facility to facility. The source listed first is considered the best or preferred source. Please use this source whenever possible. All subsequent sources are listed in order of preference. The precise location within the records where an item can be found is further identified by under and or.  Example:  To determine whether gestational diabetes is recorded as a "Risk factor in this pregnancy" (item #14) in the records:  The first or best source is the prenatal care record.  Within the prenatal care record, information on diabetes may be found under:  • Medical history  • Previous obstetric (OB) history  • Problem list or initial risk assessment  • Historical risk summary  • Complications of previous pregnancies  • Factors this pregnancy | <ul> <li>▶ Identifies alternative, usually synonymous terms and common abbreviations and acronyms for items. The keywords and abbreviations given in this guide are not intended as inclusive. Facilities and practitioners will likely add to the lists.</li> <li>Example:         <ul> <li>Keywords and abbreviations for prepregnancy diabetes are:</li> <li>DM—Diabetes mellitus</li> <li>Type 1 diabetes</li> <li>IDDM—Insulin dependent diabetes mellitus</li> <li>Type 2 diabetes</li> <li>Noninsulin dependent diabetes mellitus</li> <li>Class B DM</li> <li>Class C DM</li> <li>Class C DM</li> <li>Class T DM</li> <li>Class T DM</li> <li>Class H DM</li> </ul> </li> <li>▶ Medications commonly used for items.</li> <li>Example: "Clomid" for "Assisted reproduction treatment."</li> </ul> |

## **How to Use This Guide—Continued**

This guide was developed to assist in completing the facility worksheets for the revised Certificate of Live Birth and Report of Fetal Death. (birth certificate [BC], facility worksheet for the report of fetal death [FDFWS], report of fetal death [FDR])

| Definitions         | Instructions  | Sources                                     | Keywords and abbreviations  |
|---------------------|---|---|---|
|                     |   |   | ► Look for is used to indicate terms that may be associated with, but are not synonymous with, an item. Terms listed under look for may indicate that an item should be reported for the pregnancy, but additional information will be needed before it can be determined whether the item should be reported.  Example: "Trial of labor" for "cesarean delivery" |
| MISSING INFORMATION | Where information for an item cannot be worksheet). | located, please check "unknown" or write "t | unknown" (if using the paper copy of the  |

# Guide to Completing the Facility Worksheet for the Certificate of Live Birth and Report of Fetal Death

| Definitions   | Instructions   | Sources   | Keywords and abbreviations |
|---|--|-----------|----------------------------|
|   |  |           |                            |
| Mother  |  |           |                            |
| The woman who gave birth to, or delivered the infant.   | All birth certificate information reported for the mother should be for the woman who delivered the infant.  In cases of surrogacy or gestational carrier, the information reported should be for the surrogate or the gestational carrier, that is, the woman who delivered the infant.   |           |                            |
|   | FACILITY IN  | FORMATION |                            |
| 1. Facility name (BC #5, FDFWS #1, F                    | DR #8)   |           |                            |
| The name of the facility where the delivery took place. | Enter the name of the facility where the birth occurred.  If this birth did not occur in a hospital or freestanding birthing center, enter the street and number of the place where the birth occurred.  If this birth occurred en route to a hospital or freestanding birthing center, that is, in a moving conveyance, enter the city, town, village, or location where the child was first removed from the conveyance.  If the birth occurred in international waters or air space, enter "boat" or "plane." |           |                            |

| Definitions  | Instructions  | Sources | Keywords and abbreviations |
|--|---|---------|----------------------------|
| 2. Facility ID (BC #17, FDFWS #2, FD   | R #9)   |         |                            |
| National Provider Identifier.  | Enter the facility's National Provider Identifier (NPI) number.  If no NPI, enter the state hospital code.  |         | NPI                        |
| 3. City, town, or location of birth (BC  | #6, FDFWS #3, FDR #5)   |         |                            |
| The name of the city, town, township, village, or other location where the birth occurred. | Enter the name of the city, town, township, village, or other location where the birth occurred.  If the birth occurred in international waters or air space, enter the location where the infant was first removed from the boat or plane. |         |                            |
| 4. County of birth (BC #7, FDFWS #4,   | FDR #6)   |         |                            |
| The name of the county where the birth occurred.   | Enter the name of the county where the birth occurred.  If the birth occurred in international waters or air space, enter the name of the county where the infant was removed from the boat or plane.                                       |         |                            |

| Definitions   | Instructions   | Sources  | Keywords and abbreviations       |
|---|--|--|----------------------------------|
| 5. Place where birth occurred (Birthpl  | ace) (BC #26, FDFWS #5, FDR #7)  |  |                                  |
| The type of place where the birth occurred.  Hospital  Freestanding birthing center No direct physical connection with an operative delivery center.  Home birth The birth occurred at a private residence.  Clinic/doctor's office Other | Check the box that best describes the type of place where the birth occurred.  If home birth is checked, check whether the home birth was planned. If unknown whether a planned home birth, write "unknown."  Specify taxi, train, plane, etc. | 1st Admission history and physical (H&P) under—General Admission under—  • Admitted from home, doctor's office, other or—  • Problem list or findings  2nd Delivery record under—  • Delivery information  • Labor and delivery summary (L&D)  • Maternal obstetric (OB) or labor summary under—delivery  3rd Basic admission data  4th Progress notes or Note | FBC–Freestanding birthing center |

| ge 10   |   | The state of the s | in the Certificate of Live Birth and Report of Fetar |  |
|---|---|--|--|--|
| Definitions   | Instructions  | Sources  | Keywords and abbreviations                           |  |
| PRENATAL CARE AND PREGNANCY HISTORY  The prenatal care record is the preferred source for items 6 through 16. If the prenatal care record is not in the mother's file, please contact the prenatal care provider and obtain a copy of the record.   |   |  |  |  |
| 6. Date of first prenatal care visit (BC  | #29a, FDFWS #6a, FDR #23a)  |  |  |  |
| The date a physician or other health care professional first examined or counseled the pregnant woman for the pregnancy.  | Enter the month, day, and year of the first prenatal care visit.  If date information is incomplete, enter all parts of the date that are known. Report "unknown" for any parts of the date that are missing.  If mother's earliest prenatal care records are not available (i.e., the date of the first prenatal care visit is unavailable), report "unknown."  If "no prenatal care," check the box and enter "0" for item "total number of prenatal care visits."  | 1st Prenatal care record <i>under</i> —  • Intake information  • Initial physical examination  • Prenatal visit flow sheet  • Current pregnancy  2nd Initial physical examination  | PNC–Prenatal care                                    |  |
| 7. Total number of prenatal care visits   | for this pregnancy (BC #30)   |  |  |  |
| The total number of visits recorded in the record.  A prenatal visit is one in which the physician or other health care professional examines or counsels the pregnant woman for her pregnancy.  Do not include visits for laboratory and other testing in which a physician or health care professional did not examine or counsel the pregnant woman. | Access the most recent prenatal records available. If up-to-date records are not available, contact the prenatal care provider for the most current information.  Count the prenatal visits recorded in the record. Exclude visits for laboratory and other tests or classes in which the mother was not seen by a physician or other health care professional for pregnancy-related care. If it is not clear whether the mother was seen by a physician or other health care professional, include the visit(s) in the total number. | 1st Prenatal care record under— Prenatal visit flow sheet (count visits)   | PNC-Prenatal care                                    |  |
| continued on next page  | continued on next page  |  |  |  |

| Definitions  | Instructions   | Sources  | Keywords and abbreviations |  |  |
|--|--|--|----------------------------|--|--|
| 7. Total number of prenatal care visits  | 7. Total number of prenatal care visits for this pregnancy (BC #30)—Continued  |  |                            |  |  |
| Do not include classes, such as childbirth classes, where the physician or health care professional did not provide individual care to the pregnant woman. | Count only visits recorded in the most current record available. Do not estimate additional prenatal visits when the prenatal record is not up to date.  Enter the total number of prenatal visits.  If no visits are recorded, enter "0." If the number of prenatal visits is equal to 0, the "no prenatal care" box should also be checked for item "Date of first prenatal care visit." | See above  | See above                  |  |  |
| 8. Date last normal menses began (BC   | #30, FDFWS #8, FDR #32)  |  |                            |  |  |
| The date the mother's last normal menstrual period began.  This item is used to compute the gestational age of the infant.                                 | Enter <u>all</u> known parts of the date the mother's last normal menstrual period began. Report "unknown" for any parts of the date that are missing. <u>Do not estimate the date.</u>  | <ul> <li>1st Prenatal care record <i>under</i>—</li> <li>Menstrual history</li> <li>Labor and delivery nursing admission triage form</li> <li>2nd Admission history and physical (H&amp;P) <i>under</i>—Medical history</li> </ul> | LMP–Last menstrual period  |  |  |

| Definitions   | Instructions   | Sources  | Keywords and abbreviations   |  |  |
|---|--|--|--|--|--|
| 9. Number of previous live births now                     | 9. Number of previous live births now living (BC #35a, FDFWS #9, FDR #29a)   |  |  |  |  |
| The total number of previous liveborn infants now living. | Do not include this infant.  Include all previous live-born infants who are still living.  For multiple deliveries:  Include all live-born infants before this infant in the pregnancy.  If the first born, do not include this infant.  If the second born, include the first born, etc.  If no previous live-born infants, check "none."  See "Attachment to the Facility Worksheet for the Live Birth Certificate for Multiple Births." | <ul> <li>1st Prenatal care record under—</li> <li>Intake information</li> <li>Gravida section—L (living)—last number in series</li> <li>Para section—L—last number in series</li> <li>Pregnancy history information</li> <li>Previous OB history</li> <li>Past pregnancy history</li> <li>2nd Labor and delivery nursing admission triage form under—Patient data</li> <li>3rd Admission history and physical (H&amp;P)</li> </ul> | L-Now living  Look for:  G-Gravida-Total number of pregnancies  P-Para-Previous live births and fetal deaths > 28 weeks of gestation  T-Term-Delivered at 37 to 40 weeks gestation |  |  |

| Definitions   | Instructions   | Sources   | Keywords and abbreviations |  |  |
|---|--|---|----------------------------|--|--|
| 10. Number of previous live births now                  | 10. Number of previous live births now dead (BC #35b, FDFWS #10, FDR #29b)   |   |                            |  |  |
| The total number of previous liveborn infants now dead. | Do not include this infant.  Include all previous live-born infants who are no longer living.  For multiple deliveries:  Include all live-born infants before this infant in the pregnancy who are now dead.  If the first born, do not include this infant.  If the second born, include the first born, etc.  If no previous live-born infants now dead, check "none."  See "Attachment to the Facility Worksheet for the Live Birth Certificate for Multiple Births." | 1st Prenatal care record <i>under</i> —  • Pregnancy history information—comments, complications  • Previous OB history—comments, complications  • Past pregnancy history—comments, complications  2nd Admission history and physical (H&P) | See above<br>Expired       |  |  |
| 11. Date of last live birth (BC #35c, FD                | FWS #11, FDR #29c)   |   |                            |  |  |
| The date of birth of the last live-born infant.         | If applicable, enter the month and year.  Include live-born infants now living and now dead.  If date information is incomplete, enter all parts of the date that are known. Report "unknown" for any parts of the date that are missing. Do not estimate or guess a date.   | 1st Prenatal care record <i>under</i> —  • Pregnancy history information—date  • Previous OB history—date  • Past pregnancy history—date  2nd Admission history and physical (H&P)  | DOB–Date of birth          |  |  |

| Definitions  | Instructions   | Sources  | Keywords and abbreviations   |  |
|--|--|--|--|--|
| 12. Number of other pregnancy outcomes (BC #36a)   |  |  |  |  |
| Total number of other pregnancy outcomes that did not result in a live birth.  Includes pregnancy losses of any gestational age.  Examples: spontaneous or induced losses or ectopic pregnancy | Include all previous pregnancy losses that did not result in a live birth.  If no previous pregnancy losses, check "none."  For multiple deliveries:  Include any losses regardless of gestational age that occurred before the delivery of this infant. This could include losses occurring in this pregnancy or in a previous pregnancy. | 1st Prenatal care record under—  • Gravida section—"A" (abortion or miscarriage)  • PARA section—"A"  • Pregnancy history information—comments, complications  • Previous OB history—comments, complications  • Past pregnancy history—comments, complications  2nd Labor and delivery nursing admission triage form  3rd Admission history and physical (H&P) | Miscarriages Fetal demise AB-Abortion induced SAB-Spontaneous abortion TAB-Therapeutic abortion Abortion spontaneous Septic abortion Ectopic pregnancy Tubal pregnancy FDIU-Fetal death in utero IUFD-Intrauterine fetal death |  |
| 13. Date of last other pregnancy outcom  | me (BC #36b)   |  |  |  |
| The date the last pregnancy that did not result in a live birth ended.  Includes pregnancy losses at any gestational age.  Examples: spontaneous or induced losses or ectopic pregnancy        | If applicable, enter the month and year.  If date information is incomplete, enter all parts of the date that are known. Report "unknown" for any parts of the date that are missing. Do not estimate or guess a date.   | 1st Prenatal care record <i>under</i> —  • Pregnancy history information  • Previous OB history  • Past pregnancy history  2nd Admission history and physical (H&P)  |  |  |
| 14. Risk factors in this pregnancy (BC   | #41, FDFWS #14, FDR #36)   |  |  |  |
| Risk factors of the mother during this pregnancy.  | Check all boxes that apply. The mother may have more than one risk factor.  If the mother has none of the risk factors, check "none of the above."   | See below  | See below  |  |

| Definitions                                       | Instructions   | Sources  | Keywords and abbreviations   |  |  |  |
|---|--|--|--|--|--|--|
| 14. Risk factors in this pregnancy (BC            | 14. Risk factors in this pregnancy (BC #41, FDFWS #14, FDR #36)—Continued                      |  |  |  |  |  |
| Diabetes Glucose intolerance requiring treatment. | If diabetes is present, check either prepregnancy or gestational diabetes.  Do not check both. | <ul> <li>1st Prenatal care record <i>under</i>—</li> <li>Medical history</li> <li>Previous OB history <i>under</i>—summary of previous pregnancies</li> </ul>  |  |  |  |  |
| Prepregnancy Diagnosis before this pregnancy.     |  | <ul> <li>Problem list <i>or</i>—initial risk assessment</li> <li>Historical risk summary</li> <li>Complications of previous pregnancies</li> <li>Factors this pregnancy</li> <li>2nd Labor and delivery nursing admission triage form <i>under</i>—</li> <li>Medical complications</li> <li>Comments</li> <li>3rd Admission history and physical (H&amp;P) <i>under</i>—</li> <li>Current pregnancy history</li> </ul> | Prepregnancy: DM-Diabetes mellitus Type 1 diabetes IDDM-Insulin dependent diabetes mellitus Type 2 diabetes Non-insulin dependent diabetes mellitus Class B DM Class C DM Class C DM Class F DM Class R DM Class R DM Class H DM |  |  |  |
| Gestational Diagnosis during this pregnancy.      |  | <ul> <li>Medical history</li> <li>Previous OB history underpregnancy related</li> <li>Problem list or findings</li> <li>4th Delivery record under-</li> <li>Maternal OB or labor summary</li> <li>Labor and delivery admission history</li> <li>Labor summary record</li> </ul>  | Gestational: GDM-Gestational diabetes mellitus IDGDM-Insulin dependent gestational diabetes mellitus Class A1 or A2 diabetes mellitus  |  |  |  |

| Definitions  | Instructions   | Sources      | Keywords and abbreviations   |  |  |
|--|--|--------------|--|--|--|
| 14. Risk factors in this pregnancy (BC   | 14. Risk factors in this pregnancy (BC #41, FDFWS #14, FDR #36)—Continued                              |              |  |  |  |
| Hypertension Elevation of blood pressure above normal for age, sex, and physiological condition.   | If hypertension is present, check either prepregnancy or gestational hypertension. Do not check both.  | See Diabetes |  |  |  |
| Prepregnancy (chronic) Diagnosis prior to the onset of this pregnancy—does not include gestational (pregnancy-induced hypertension [PIH]). |  |              | Prepregnancy: CHT–Chronic hypertension Benign essential hypertension Essential hypertension Preexisting hypertension |  |  |
| Gestational Diagnosis in this pregnancy (Pregnancy-induced hypertension or preeclampsia).  |  |              | Gestational: PIH–Pregnancy-induced hypertension Preeclampsia Eclampsia Transient hypertension HELLP Syndrome         |  |  |
| Eclampsia Hypertension with proteinuria with generalized seizures or coma. May include pathologic edema.                                   | If eclampsia is present, one type of hypertension (either gestational or prepregnancy) may be checked. | See Diabetes | See Hypertension   |  |  |

| Definitions   | Instructions                       | Sources  | Keywords and abbreviations    |
|---|------------------------------------|--|-------------------------------|
| 14. Risk factors in this pregnancy (BC  | #41, FDFWS #14, FDR #36)—Continued |  |                               |
| Previous preterm births History of pregnancy(ies) terminating in a live birth of less than 37 completed weeks of gestation. |                                    | <ul> <li>1st Prenatal care record under—</li> <li>• Medical history</li> <li>• Previous OB history under—summary of previous pregnancies</li> <li>• Problem list or—initial risk assessment</li> <li>• Historical risk summary</li> <li>• Complications of previous pregnancies</li> <li>2nd Labor and delivery nursing admission triage form under—</li> <li>• Medical complications</li> <li>• Comments</li> <li>3rd Admission history and physical (H&amp;P) under—</li> <li>• Medical history</li> <li>• Previous OB history under—pregnancy related</li> <li>• Problem list/findings</li> </ul> | PTL-Preterm labor P-Premature |

| Definitions   | Instructions  | Sources   | Keywords and abbreviations  |  |  |  |
|---|---|---|---|--|--|--|
| 14. Risk factors in this pregnancy (BC  | 14. Risk factors in this pregnancy (BC #41, FDFWS #14, FDR #36)—Continued   |   |   |  |  |  |
| Pregnancy resulted from infertility treatment Any assisted reproductive treatment used to initiate the pregnancy.  Includes: - Drugs (e.g., Clomid or Pergonal) - Artificial insemination - Technical procedures (e.g., in vitro fertilization) | Check if any infertility treatment was used to initiate the pregnancy. This information is included on both the mother's and the facility worksheets. Check "yes" if treatment is reported on either worksheet. | <ul> <li>1st Prenatal care record under—</li> <li>Medical history</li> <li>Current pregnancy history</li> <li>Problem list or—initial risk assessment</li> <li>Medications this pregnancy</li> <li>2nd Labor and delivery nursing admission triage form under—</li> <li>Comments</li> <li>Medications</li> <li>3rd Admission history and physical (H&amp;P) under—</li> <li>Current pregnancy history</li> <li>Problem list/findings</li> </ul> | See below   |  |  |  |
| Fertility-enhancing drugs, artificial insemination, or intrauterine insemination  Any fertility-enhancing drugs (e.g., Clomid or Pergonal), artificial insemination, or intrauterine insemination used to initiate the pregnancy.               | Check if specific therapy (drugs or insemination) was used. This information is included on both the mother's and the facility worksheets. Check "yes" if treatment is reported on either worksheet.            | See Pregnancy resulted from infertility treatment   | Fertility-enhancing drugs, or artificial or intrauterine insemination:  Medications Clomid, Serophene Pergonal Metrodin Profasi Progesterol Crinone (progesterone gel) Follistim FSH–Follicle stimulating hormone Gonadotropins Hcg–Human chorionic gonadotropin IUI– Intrauterine insemination |  |  |  |

| Definitions   | Instructions  | Sources  | Keywords and abbreviations   |  |  |  |
|---|---|--|--|--|--|--|
| 14. Risk factors in this pregnancy (BC  | 14. Risk factors in this pregnancy (BC #41, FDFWS #14, FDR #36)—Continued   |  |  |  |  |  |
| Assisted reproductive technology (e.g., in-vitro fertilization [IVF] gamete intrafallopian transfer [GIFT])  Any assisted reproductive technology ([ART]/technical procedures [e.g., IVF, GIFT, or ZIFT]) used to initiate the pregnancy. | Check if assisted reproductive therapy was used. This information is included on both the mother's and the facility worksheets. Check "yes" if treatment is reported on either worksheet. | See Pregnancy resulted from infertility treatment  | Assisted reproductive technology: ART Artificial insemination AIH-Artificial insemination by husband AID/DI-Artificial insemination by donor In vitro fertilization IVF-ET-In vitro fertilization embryo transfer GIFT-Gamete intrafallopian transfer ZIFT-Zygote intrafallopian transfer Ovum donation Donor embryo Embryo adoption |  |  |  |
| Mother had a previous cesarean delivery Previous delivery by extracting the fetus, placenta, and membranes through an incision in the mother's abdominal and uterine walls.  If yes, how many?  | If the mother has had a previous cesarean delivery, indicate the number of previous cesarean deliveries she has had.  | 1st Prenatal care record <i>under</i> —  • Past pregnancy history  • Past OB history  • Problem list <i>or</i> —initial risk assessment  2nd Labor and delivery nursing admission triage form <i>under</i> —Comments  3rd Admission history and physical (H&P) <i>under</i> —  • Past OB history  • Past pregnancy history <i>under</i> —problem list/findings | C/S-Cesarean section Repeat C/S VBAC-Vaginal delivery after cesarean LSTCS (or LTCS) low segment transverse cesarean section Classical cesarean section Low vertical C/S Low transverse C/S Look for: TOL-Trial of labor   |  |  |  |

| Definitions   | Instructions   | Sources   | Keywords and abbreviations   |  |
|---|--|---|--|--|
| 15. Infections present and/or treated during this pregnancy (BC #42)  |  |   |  |  |
| Infections present at the time of the pregnancy diagnosis or a confirmed diagnosis during the pregnancy with or without documentation of treatment.  Documentation of treatment during this pregnancy is adequate if a definitive diagnosis is not present in the available record. | Check all boxes that apply. The mother may have more than one infection.  If the mother has none of the infections, check "none of the above." | See below   | "+" indicates that the test for the infection was positive and the woman has the infection.  "-" indicates that the test was negative, and the woman does not have the infection.  Look for: treatment or Rx for specific infection. |  |
| Gonorrhea A positive test or culture for Neisseria gonorrhoeae.   |  | 1st Prenatal record under—  • Infection history  • Sexually transmitted diseases  • Problem list  • Complications this pregnancy  • Factors this pregnancy  • Medical history  2nd Labor and delivery nursing admission triage form under— Comments  3rd Admission history and physical (H&P) under—  • Current pregnancy history  • Medical history  • Medical history  • Problem list/findings  4th Delivery record under—  • Maternal OB/labor summary  • Labor and delivery admission | GC Gonorrheal Gonococcal Treatment or Rx for Gonorrhea NAAT–Nucleic amplification tests  |  |

| Definitions   | Instructions                           | Sources       | Keywords and abbreviations  |
|---|--|---------------|---|
| 15. Infections present and/or treated de  | uring this pregnancy (BC #42)—Continue | d             |   |
| Syphilis (also called lues) A positive test for Treponema pallidum.                     |  | See Gonorrhea | TP-PA-T. pallidum particle agglutination STS-Serologic test for syphilis RPR-Rapid plasma regain VDRL-Venereal disease research laboratories FTA-AS-Fluorescent antibody test Lues Treatment or Rx for syphilis or lues |
| Chlamydia A positive test for Chlamydia trachomatis.                                    |  | See Gonorrhea | Treatment or Rx for chlamydia   |
| Hepatitis B (HBV, serum hepatitis) A positive test for the hepatitis B virus.           |  | See Gonorrhea | Hep B HBV Treatment or Rx for hepatitis B   |
| Hepatitis C (non A or non B hepatitis [HCV]) A positive test for the hepatitis C virus. |  | See Gonorrhea | Hep C HCV Treatment or Rx for hepatitis C   |

| Definitions  | Instructions  | Sources   | Keywords and abbreviations  |
|--|---|---|---|
| 16. Obstetric procedures (BC #43)  |   |   |   |
| Medical treatment or invasive or manipulative procedure performed during this pregnancy to treat the pregnancy or to manage labor or delivery.   | If the mother has had none of the procedures, check "none of the above."  | See below   | See below   |
| External cephalic version  Attempted conversion of a fetus from a nonvertex to a vertex presentation by external manipulation.  Successful Fetus was converted to a vertex presentation.  Failed Fetus was not converted to a vertex presentation. | If checked, also indicate whether the procedure was a success or a failure.  If more than one attempt, report results of most recent attempt. | 1st Prenatal care record under—  • Problem list  • Historical risk summary  • Complications this pregnancy  • Factors this pregnancy  2nd Labor and delivery nursing admission triage form under—  • Complications  • Comments  3rd Admission history and physical (H&P) under—  • Current pregnancy history  • Medical history  • Problem list/findings  4th Delivery record under—  • Maternal OB/labor summary  • Labor and delivery admission history  • Labor summary record | Successful version: Breech version External version  Failed version: Unsuccessful external version Attempted version Failed version Look for: malpresentation |

| Definitions  | Instructions   | Sources  | Keywords and abbreviations |  |  |
|--|--|--|----------------------------|--|--|
|  | LABOR AND DELIVERY   |  |                            |  |  |
| 17. Date of birth (BC #4, FDFWS #16, I   | FDR #4)  |  |                            |  |  |
| The infant's date of birth.  | Enter the month, day, and four-digit year of birth.  If the date of birth of the infant is unknown because the infant is a foundling, enter the date the infant was found. | 1st Labor and delivery <i>under</i> — Delivery record  2nd Newborn admission H&P | DOB–Date of birth          |  |  |
| 18. Time of birth (BC #2, FDFWS #17,   | FDR #2)  |  |                            |  |  |
| The infant's time of birth.  | Enter the time the infant was born based on a 24-hour clock (military time). If time of birth is unknown (foundling), enter "unknown."                                     | 1st Labor and delivery <i>under</i> — Delivery record  2nd Newborn admission H&P |                            |  |  |
| 19. Certifier's name and title (BC #11)  |  |  |                            |  |  |
| The individual who certified to the fact that the birth occurred:  M.D. (doctor of medicine)  D.O. (doctor of osteopathy)  Hospital administrator or designee  CNM/CM (certified nurse midwife or certified midwife)  Other midwife (midwife other than CNM/CM)  Other (specify) | Enter the name and title of the individual who certified to the fact that the birth occurred.  The individual may be, but need not be, the same as the attendant at birth. |  |                            |  |  |
| 20. Date certified (BC #12)  | 20. Date certified (BC #12)  |  |                            |  |  |
| The date the birth was certified.  | Enter the date the birth was certified.  |  |                            |  |  |

| Definitions   | Instructions   | Sources   | Keywords and abbreviations |
|---|--|---|----------------------------|
| 21. Principal source of payment (BC #3  | 8)   |   |                            |
| The primary source of payment for the delivery at the time of delivery:  Private insurance (Blue Cross/Blue Shield, Aetna, etc.)  Medicaid (or a comparable state program)  Self-pay (no third party identified)  Other (Indian Health Service; CHAMPUS or TRICARE; other government [federal, state, or local]; or charity)  The principal source of payment is important public health information and is needed to monitor access to care during delivery. | Check the box that best describes the primary source of payment for this delivery. If more than one source of payment for the delivery is recorded, choose the source that appears to pay for most of the delivery.  Check the source of payment for the delivery, not the payer for the newborn care or prenatal care, if different.  If "other" is checked, specify the payer.  If the principal source of payment is not known, enter "unknown" in the space.  If unsure what source of payment a given insurance falls under, check with the billing office. | 1st Hospital face sheet 2nd Admitting office face sheet   |                            |
| 22. Infant's medical record number (Bo  | C #48)   |   |                            |
| The medical record number assigned to the newborn.  | Enter the medical record number.   | 1st Infant's medical record addressograph plate  2nd Admitting office face sheet under-History number |                            |

| Definitions   | Instructions  | Sources  | Keywords and abbreviations |
|---|---|--|----------------------------|
| 23. Was the mother transferred to this  | facility for maternal medical or fetal indic  | cations for delivery? (BC #28)   |                            |
| Transfers include hospital to hospital, birth facility to hospital, etc. Does not include home to hospital. | If the mother was transferred from another <u>facility</u> to this facility for medical reasons related to the pregnancy, check "yes."  If yes, enter the name of the facility the mother was transferred from. If the name of the facility is not known, enter "unknown."  Check "no" if the mother was transferred from home. | 1st Labor and delivery nursing admission triage form under—  • Reason for admission  • Comments  2nd Admission history and physical (H&P)  3rd Labor & delivery – Delivery record  • Maternal OB/labor summary  • Labor and delivery admission history  • Labor summary record |                            |

| Definitions  | Instructions  | Sources   | Keywords and abbreviations |  |  |  |
|--|---|---|----------------------------|--|--|--|
| 24. Attendant's name, title, and ID (BC  | 24. Attendant's name, title, and ID (BC #27, FDFWS #21, FDR #14)  |   |                            |  |  |  |
| The name, title, and National Provider Identifier (NPI) number of the person responsible for delivering the child.  M.D. (doctor of medicine)  D.O. (doctor of osteopathy)  CNM/CM (certified nurse midwife or certified midwife)  Other midwife (midwife other than CNM/CM)  Other (specify)  The attendant at birth is the individual physically present at the delivery who is responsible for the delivery. For example, if an intern or nurse midwife delivers an infant under the supervision of an obstetrician who is present in the delivery room, the obstetrician should be reported as the attendant. If the obstetrician is not physically present, the intern or nurse midwife must be reported as the | Enter the name, title, and NPI number of the person responsible for delivering the child.  Check one box to specify the attendant's title. If "other" is checked, enter the specific title of the attendant. Examples include nurse, father, police officer, and EMS technician.  This item should be completed by the facility. If the birth did not occur in a facility, the attendant or certifier should complete it. | 1st Delivery record under— Signature of delivery attendant (medical)  |                            |  |  |  |
| attendant.   |   |   |                            |  |  |  |
| 25. Mother's weight at delivery (BC #33  | 3)  |   |                            |  |  |  |
| The mother's weight at the time of delivery.   | Enter the mother's weight at the time of delivery. Use pounds in whole numbers only. For example, enter 140½ pounds as 140 pounds.  If the mother's delivery weight is unknown, enter "unknown."  | 1st Labor and delivery nursing admission triage form <i>under</i> —Physical assessment—Weight  2nd Admission history and physical (H&P) <i>under</i> —Physical examination—Weight | Wgt–Weight                 |  |  |  |

| Definitions  | Instructions   | Sources   | Keywords and abbreviations   |
|--|--|---|--|
| 26. Characteristics of labor and deliver   | ry (BC #45)  |   |  |
| Information about the course of labor and delivery.  | Check all characteristics that apply.  If none of the characteristics of labor and delivery apply, check "none of the above."  | See below   | See below  |
| Induction of labor Initiation of uterine contractions by medical or surgical means for the purpose of delivery before the spontaneous onset of labor (i.e., before labor has begun).  Examples of methods include, but are not limited to: artificial rupture of membranes, balloons, oxytocin, prostaglandin, laminaria, or other cervical ripening agents. | Check this item if medication was given or procedures to induce labor were performed before labor began. If it is not clear whether medication or procedures were performed before or after labor had begun, review records to determine when labor began and when medications were given or procedures performed. If this information is unclear or unavailable, check with the birth attendant.  Induction of labor should be checked even if the attempt to initiate labor is not successful or the induction follows a spontaneous rupture of the membrane without contractions.  NOTE: Does not include augmentation of labor, which applies only after labor or contractions have begun. | 1st Delivery record under— Maternal OB/labor summary  • Labor and delivery admission history  • Labor summary record  2nd Physician progress note  3rd Labor and delivery nursing admission triage form | IOL-Induction of labor Pit Ind-Pitocin induction ROM/NIL-Amniotomy induction or induction for rupture of membranes, not in labor AROM-Artificial rupture of membranes done before labor Balloons Oxytocin Prostaglandin Laminaria Cervidil |

| Definitions   | Instructions   | Sources   | Keywords and abbreviations  |  |  |
|---|--|---|---|--|--|
| 26. Characteristics of labor and deliver  | 26. Characteristics of labor and delivery (BC #45)—Continued   |   |   |  |  |
| Augmentation of labor Stimulation of uterine contractions by drug or manipulative technique with the intent to reduce the time of delivery (i.e., after labor has begun).   | Check this item if medication was given or procedures to augment labor were performed after labor began. If it is not clear whether medication or procedures were performed before or after labor had begun, review records to determine when labor began and when medications were given or procedures performed. If this information is unclear or unavailable, check with the birth attendant.  NOTE: Do not include if induction of labor was performed. | Same as 1st and 2nd sources for Induction of labor  | Pit stim-Pitocin stimulation Pit aug-Pit augmentation AROM-Artificial rupture of membranes done during labor Cervidil     |  |  |
| Steroids (glucocorticoids) for fetal lung maturation received by the mother before delivery  Steroids received by the mother prior to delivery to accelerate fetal lung maturation. Typically administered in anticipation of preterm (less than 37 completed weeks of gestation) delivery.  Steroids include: betamethasone, dexamethasone, or hydrocortisone specifically given to accelerate fetal lung maturation.  Does not include steroid medication given to the mother for anti-inflammatory treatment before or after delivery. | Three conditions must be met for this item. Check this item when 1) steroid medication was given to the mother 2) prior to delivery 3) for fetal lung maturation.  Steroids may be administered to the mother prior to admittance to the hospital for delivery. Review the mother's prenatal care and other hospital records for mention of steroid administration for this purpose.   | <ul> <li>1st Delivery record under—</li> <li>Maternal OB/labor summary—comments</li> <li>Labor summary record—comments</li> <li>2nd Maternal medication record</li> <li>3rd Newborn admission H&amp;P</li> <li>4th Maternal physician order sheet</li> <li>5th Prenatal care records</li> </ul> | Medications (before delivery): Betamethasone Betamethasone phosphate Beta-PO4 Betamethasone acetate Beta-Ac Dexamethasone |  |  |

| Definitions  | Instructions   | Sources   | Keywords and abbreviations  |  |  |
|--|--|---|---|--|--|
| 26. Characteristics of labor and deliver   | 26. Characteristics of labor and delivery (BC #45)—Continued   |   |   |  |  |
| Antibiotics received by the mother during delivery Includes antibacterial medications given systemically (intravenous or intramuscular) to the mother in the interval between the onset of labor and the actual delivery. Includes: Ampicillin Penicillin Clindamycin Erythromycin Gentamicin Cefotaxime Ceftriaxone   | Mother should have undergone labor, regardless of method of delivery.  Check the timing of the administration of the antibacterial medications. Check this item only if medications were received systemically by the mother during labor. If information on onset of labor cannot be determined from the records, check with the birth attendant. | See steroids (glucocorticoids)  | Medications (during delivery): Ampicillin Penicillin Clindamycin Erythromycin Gentamicin Cefotaxime Ceftriaxone Vancomycin Look for: SBE (sub-acute bacterial endocarditis) prophylaxis GBS positive or GBS + (Group B streptococcus) Maternal fever Mother febrile |  |  |
| Clinical chorioamnionitis diagnosed during labor or maternal temperature ≥ 38°C (100.4°F) Clinical diagnosis of chorioamnionitis during labor made by the delivery attendant. Usually includes more than one of the following: fever, uterine tenderness or irritability, leukocytosis, fetal tachycardia, maternal tachycardia, or malodorous vaginal discharge.  Any recorded maternal temperature at or above 38°C (100.4°F). | Check that recorded maternal temperature is at or above 38°C (100.4°F).  | <ul> <li>1st Delivery record under—</li> <li>Maternal OB/labor summary—comments/complications</li> <li>Labor summary record—comments/complications</li> <li>2nd Newborn admission H&amp;P</li> <li>3rd Physician progress note</li> <li>4th Maternal vital signs record under—Temperature recordings</li> </ul> | Chorioamnionitis Chorio Temp > 38°C or 100.4°F  Look for: Maternal fever Mother febrile   |  |  |

| Definitions  | Instructions   | Sources  | Keywords and abbreviations  |
|--|--|--|---|
| 26. Characteristics of labor and deliver   | y (BC #45)—Continued   |  |   |
| Epidural or spinal anesthesia during labor  Administration to the mother of a regional anesthetic to control the pain of labor.  Delivery of the agent into a limited space with the distribution of the analgesic effect limited to the lower body. | Mother should have undergone labor, regardless of method of delivery.      | <ul> <li>1st Delivery record <i>under</i>—</li> <li>Maternal OB labor summary <i>under</i>—analgesia or anesthesia</li> <li>Labor summary record <i>under</i>—analgesia or anesthesia</li> </ul> | Epidural analgesia<br>Epid. given<br>Spinal given   |
| 27. Method of delivery (BC #46, FDFW   | <br>/S #23, FDR #38)   |  |   |
| The physical process by which the complete delivery of the fetus was affected.   | Complete sections C and D.   | See below  | See below   |
| C. Fetal presentation at birth  Cephalic—presenting part of the fetus listed as vertex, occiput anterior (OA), or occiput posterior (OP).  Breech—presenting part of the fetus listed as breech, complete breech, frank breech, or footling breech.  | Check one of the three boxes.  Check only the final presentation at birth. | 1st Delivery record <i>under</i> —Fetal birth presentation   | Cephalic: Vertex-OA, OP, LOA, ROA, LOP, ROP, LOT, ROT Face-LMA, LMT, LMP, RMA, RMP, RMT Brow Sinciput Mentum-chin  Breech: (Buttocks, sacrum) Frank breech-LSA, LST, LSP, RSP, RST Single footling breech |
| continued on next page   |  |  | Double footling breech Complete breech  continued on next page  |

| Definitions  | Instructions               | Sources  | Keywords and abbreviations  |
|--|----------------------------|--|---|
| 27. Method of delivery (BC #46, FDFV   | VS #23, FDR #38)—Continued |  |   |
| C. Fetal presentation at birth—Continued  Other—any other presentation not listed above.   | See above                  | See above  | Other: Shoulder Transverse lie Funis Compound   |
| D. Final route and method of delivery  Vaginal/spontaneous  Delivery of the entire fetus through the vagina by the natural force of labor with or without manual assistance from the delivery attendant.  Vaginal/forceps  Delivery of the fetal head through the vagina by the application of obstetrical forceps to the fetal head.  Vaginal/vacuum  Delivery of the fetal head through the vagina by the application of a vacuum cup or ventouse to the fetal head. | Check one of the boxes.    | 1st Delivery record <i>under</i> –Method of delivery  2nd Newborn admission H&P  3rd Recovery room record <i>under</i> – Maternal data–Delivered | Vaginal/spontaneous: VAG Del-Vaginal delivery SVD-Spontaneous vaginal delivery  Vaginal/forceps: LFD-Low forceps delivery  Vaginal/vacuum: Vac Ext vacuum |
| Cesarean  Extraction of the fetus, placenta, and membranes through an incision in the maternal abdominal and uterine walls.  |                            |  | Cesarean: C/S—Cesarean section LSTCS—Low segment transverse Look for: TOL—Trial of labor  |

| Definitions   | Instructions   | Sources  | Keywords and abbreviations   |
|---|--|--|--|
| 27. Method of delivery (BC #46, FDFV  | VS #23, FDR #38)—Continued   |  |  |
| If cesarean, was a trial of labor attempted?  Labor was allowed, augmented, or induced with plans for a vaginal delivery.   | Check "yes" or "no."   |  | TOL–Trial of labor   |
| 28. Maternal morbidity (BC #47, FDF   | WS #24, FDR #39)   |  |  |
| Serious complications experienced by the mother associated with labor and delivery.  Maternal transfusion Includes infusion of whole blood or packed red blood cells associated with labor and delivery.  | Check all boxes that apply.  If the mother has none of the complications, check "none of the above." | 1st Delivery record <i>under</i> —  • Labor summary  • Delivery summary  2nd Physician delivery notes or Operative notes  3rd Intake & output form | Transfused Blood transfusion Look for: PRBC-Packed red blood cells Whole blood |
| Third- or fourth-degree perineal laceration  3° laceration extends through the perineal skin, vaginal mucosa, perineal body, and partially or completely through the anal sphincter.  4° laceration is all of the above with extension through the rectal mucosa. |  | 1st Delivery record <i>under</i> —  • Episiotomy section  • Lacerations section  2nd Recovery room record <i>under</i> — Maternal data—Delivered   | 4th degree lac.  4° LAC degree  3rd degree lac.  3° LAC degree  3a  3b  3c     |

| Definitions   | Instructions               | Sources  | Keywords and abbreviations  |
|---|----------------------------|--|---|
| 28. Maternal morbidity (BC #47, FDF)  | WS #24, FDR #39)—Continued |  |   |
| Ruptured uterus  Tearing of the uterine wall. Uterine rupture is a full-thickness disruption of the uterine wall that also involves the overlying visceral peritoneum (uterine serosa).  Does not include uterine dehiscence, in which the fetus, placenta, and umbilical cord remain contained with the uterine cavity. Does not include a silent or incomplete rupture or an asymptomatic separation. |                            | 1st Delivery record <i>under</i> —  • Delivery summary note—  • Comments or Complications  2nd Operative note  3rd Physician progress note |   |
| Unplanned hysterectomy Surgical removal of the uterus that was not planned before the admission. Includes an anticipated, but not definitively planned, hysterectomy.   |                            | See Ruptured uterus  | Hysterectomy  Look for: laparotomy  |
| Admission to an intensive care unit<br>Any admission, planned or<br>unplanned, of the mother to a facility<br>or unit designated as providing<br>intensive care.  |                            | 1st Physician progress note 2nd Transfer note  | ICU-Intensive care unit MICU-Medical intensive care unit SICU-Surgical intensive care unit L&D ECU-Labor and delivery emergency care unit |

| Guide to Completing the Facility Worksheet for the Certaneate of Live Birth and Report of Feder 1   |  |   |  |  |
|---|--|---|--|--|
| Definitions   | Instructions   | Sources   | Keywords and abbreviations   |  |
| NEWBORN INFORMATION   |  |   |  |  |
| 29. Birthweight or weight of fetus (BC a  | #49, FDFWS #25, FDR #18c)  |   | 1  |  |
| The weight of the infant at birth.  | Enter the weight (in grams) of the infant at birth.  Do not convert pounds (lbs.) and ounces (oz.) to grams.  If the weight in grams is unavailable, enter the birth weight in lbs. and oz.  | 1st Delivery record <i>under</i> — Infant data        | BW-Birthweight Gms-Grams kg-Kilograms Lbs-Pounds oz-Ounces                                   |  |
| 30. Obstetric estimate of gestation at d  | elivery (BC #50, FDFWS #26, FDR #18d)  |   |  |  |
| The best obstetric estimate of the infant's gestational age (OE) in completed weeks is based on the clinician's final estimate of gestation.  For the clinician:  The best estimated delivery date (EDD) is determined by the date of the last menstrual period (LMP) if confirmed by early ultrasound. If ultrasound is not performed or is unknown, the best EDD is determined by the LMP. If the LMP is unknown or inconsistent with the early ultrasound, the best EDD is determined by early ultrasound. For deliveries resulting from the use of assisted reproductive technology, the best EDD is based on the known date of fertilization.*  Accurate pregnancy dating is important to improve outcomes and is a research and public health imperative. The best estimated due date should be documented clearly in the medical records.* | Enter the final best obstetric estimate of the infant's gestational age in completed weeks.  The OE should be based on the clinician's estimate of gestational age at delivery. Look for the most recent gestational age estimate in the records (this can often be found in the labor and delivery records under "gestational age").  If the most recent gestational age is dated on or after the infant's date of delivery, enter this estimate.  If the most recent gestational age is dated before the date of delivery, add the number of days between the most recent gestational age and the date of delivery to the gestational age estimate. For example, if the most recent gestational age in the records is 32 weeks, 5 days and is dated 3/24, and the date of delivery is 3/31, add 7 days to the gestational age for the final total of 33 weeks, 5 days. | 1st OB admission H&P under- • Weeks • Gestational age | Gestation weeks (wks) weeks gestational age GA-Gestational age EGA-Estimated gestational age |  |

| Definitions Inst   | ructions Sources   | Keywords and abbreviations |  |  |  |
|--|--|----------------------------|--|--|--|
| 30. Obstetric estimate of gestation at delivery (BC #50, FDFWS #26, FDR #18d)—Continued  |  |                            |  |  |  |
| Adopted from: ACOG reVITALize Obstetric Data Definitions" vailable from: https://www.acog. rg/-/media/Departments/Patient- afety-and-Quality-Improvement/ Ol4reVITALizeObstetricDataDe nitionsV10.pdf) and "Method for stimating Due Date" (available from: ttp://www.acog.org/Resources-And- ublications/Committee-Opinions/ ommittee-on-Obstetric-Practice/ lethod-for-Estimating-Due-Date).  If both the EDD known, enter "ur  If the date of de the EDD, then ur calculation:  1) Find the mos | ge for the OE is leted weeks. If the action of a week (e.g., and always round down ole week (e.g., and as a stetric estimate of the found, search for all estimated due date late the OE are not alknown."  Stivery occurs before the following and the date of delivery the date of delivery the date of delivery to follow the date of delivery the date of delivery to follow the date of delivery to follow the | See above                  |  |  |  |

| Definitions                                 | Instructions  | Sources  | Keywords and abbreviations  |  |  |
|---|---|--|---|--|--|
| 30. Obstetric estimate of gestation at d    | 30. Obstetric estimate of gestation at delivery (BC #50, FDFWS #26, FDR #18d)—Continued   |  |   |  |  |
| See above                                   | If the date of delivery occurs after the EDD, then use the following calculation:   | See above                                      | See above   |  |  |
|   | Find the most recent best EDD (this may only be found in the prenatal care records) and the date of delivery.                               |  |   |  |  |
|   | 2) Calculate the difference between the date of delivery and EDD in days (e.g., date of delivery of 1/5/2015 – EDD of 12/28/2014 = 8 days). |  |   |  |  |
|   | 3) Add the difference between the date of delivery and the EDD to 280 days (e.g., 280 + 8 = 288 days).                                      |  |   |  |  |
|   | 4) Divide the total number of days from step 3 by 7 days (e.g., 288 days ÷ 7 days = 41.1 weeks) to determine the OE in weeks.               |  |   |  |  |
|   | 5) Enter the OE in completed weeks (e.g., 41 weeks).  |  |   |  |  |
| 31. Sex of child (BC #3, FDFWS #27, FDR #3) |   |  |   |  |  |
| The sex of the infant.                      | Enter whether the infant is male, female, or if the sex of the infant is ambiguous, enter "unknown."  | 1st Delivery record <i>under</i> — Infant data | M-Male F-Female A-Ambiguous or Not yet determined (same as unknown) U-Unknown |  |  |

| Definitions   | Instructions  | Sources  | Keywords and abbreviations   |
|---|---|--|--|
| 32. Apgar score (BC #51)  |   |  |  |
| A systematic measure for evaluating the physical condition of the infant at specific intervals following birth.   | Enter the infant's Apgar score at 5 minutes.  If the score at 5 minutes is less than 6, enter the infant's Apgar score at 10 minutes.   | 1st Delivery record <i>under</i> —Infant data                  |  |
| 33. Plurality (BC #52, FDFWS #28, FD  | PR #33)   |  |  |
| The number of fetuses delivered live or dead at any time in the pregnancy regardless of gestational age, or if the fetuses were delivered at different dates in the pregnancy.  "Reabsorbed" fetuses (those that are not delivered: expulsed or extracted from the mother) should not be counted. | Enter the number of fetuses delivered in this pregnancy.  If two or more live births in this delivery, see "Attachment to the Facility Worksheet for the Live Birth Certificate for Multiple Births." | 1st Delivery record  2nd Admission history and physical (H&P)  | Single Twin, triplet, quadruplet, etc. Multiple (a, b, c) or (1, 2, 3)   |
| 34. If not a single birth, order born in t  | the delivery (BC #53, FDFWS #29, FDR #3   | 34)  |  |
| The order born in the delivery, live born or fetal death (1st, 2nd, 3rd, 4th, 5th, 6th, 7th, etc.).   | If this is a single birth, leave this item blank.  Include all live births and fetal deaths from this pregnancy.  | 1st Delivery record <i>under</i> —Birth order  2nd Infant data | Baby A, B, or Baby 1, 2, etc. Twin A, B, or Twin 1, 2 Triplet A, B, C, or Triplet 1, 2, 3, etc. Look for: Birth order or Set order |
| 35. If not a single birth, number of infa   | ents in the delivery born alive (FDFWS #30  | 0)   |  |
| The number of infants in this delivery born alive at any point in the pregnancy.  | If this is a single birth, leave this item blank.  If this is not a single birth, specify the number of infants in this delivery born alive at any point in the pregnancy. Include this birth.        | 1st Delivery record 2nd Admission history and physical (H&P)   | Look for:<br>Condition   |

| Definitions  | Instructions   | Sources  | Keywords and abbreviations   |
|--|--|--|--|
| 36. Abnormal conditions of the newbor  | n (BC #54)   |  |  |
| Disorders or significant morbidity experienced by the newborn.   | Check all boxes that apply.  If none of the conditions apply, check "none of the above." | See below  | See below  |
| Assisted ventilation required immediately following delivery Infant given manual breaths for any duration with bag and mask or bag and endotracheal tube within the first several minutes from birth.  Excludes free-flow (blow-by) oxygen only, laryngoscopy for aspiration of meconium, nasal cannula, and bulb suction. |  | 1st Labor delivery summary under— Infant Data or Breathing | Bag and mask ventilation Intubation Intubation and PPV–Positive pressure ventilation PPV bag/mask or ET–Positive pressure ventilation via bag, mask, or endotracheal intubation IPPV bag–Intermittent positive pressure ventilation via bag IPPV ET–Intermittent positive pressure ventilation via endotracheal intubation O2 via ET–Oxygen via endotracheal intubation Oxygen |

| Definitions   | Instructions   | Sources                                 | Keywords and abbreviations  |
|---|--|---|---|
| 36. Abnormal conditions of the newbor   | n (BC #54)—Continued                                       |   |   |
| Assisted ventilation required for more than six hours  Infant given mechanical ventilation (breathing assistance) by any method for more than six hours.  Includes conventional, high frequency, or continuous positive pressure (CPAP).  Excludes free-flow oxygen only, laryngoscopy for aspiration of meconium, and nasal cannula. | Count the number of hours of mechanical ventilation given. | 1st Newborn respiratory care flow sheet | If in use for more than 6 hours:  CPAP-Continuous positive airway pressure  IPPV-Intermittent positive pressure ventilation  HFV-High frequency ventilation  IMV-Intermittent mandatory volume ventilation  HFOV-High frequency oscillatory ventilation  IPPV-Intermittent positive pressure ventilation  PIP-Peak inspiratory pressure  PEEP-Positive end expiratory pressure  CMV-Continuous mandatory ventilation  HFPPV-High frequency positive pressure ventilation  HFPPV-High frequency flow interruption ventilation  HFJV-High frequency jet ventilation  Inhaled nitric oxide |

| Definitions   | Instructions   | Sources  | Keywords and abbreviations  |
|---|--|--|---|
| 36. Abnormal conditions of the newbor   | rn (BC #54)—Continued  |  |   |
| NICU admission  Admission into a facility or unit staffed and equipped to provide continuous mechanical ventilatory support for a newborn.  | Include NICU admission at any time during the infant's hospital stay following delivery.  Do not include units that do not provide continuous mechanical ventilation. Do not include well-baby nurseries or special care nurseries (i.e., Level II nursery).  Do not include if the newborn was taken to the NICU for observation but is not admitted to the NICU. | 1st Labor and delivery summary record <i>under</i> —Disposition <i>under</i> —  • Intensive care nursery (ICN)  • Special care nursery (SCN) | ICN-Intensive care nursery SCN-Special care nursery NICU-Neonatal intensive care unit PICU-Pediatric intensive care unit Level II nursery   |
| Newborn given surfactant replacement therapy Endotracheal instillation of a surface-active suspension for treating surfactant deficiency due to preterm birth or pulmonary injury resulting in respiratory distress.  Includes both artificial and extracted natural surfactant.                  | Check both 1st and 2nd sources before completion.  | 1st Labor and delivery summary under—Neonatal medication  2nd Newborn medication administration record                                       | If given to newborn after birth:  Medications (given to newborn): Surfactant Survanta Exosurf Curosurf Infasurf   |
| Antibiotics received by the newborn for suspected neonatal sepsis Any antibacterial drug (penicillin, ampicillin, gentamicin, cefotaxime, etc.) given systemically (intravenous or intramuscular). Does not include antibiotics given to infants who are not suspected of having neonatal sepsis. |  | 1st Newborn medication administration record   | Medications (given to newborn for sepsis): Nafcillin, Chloramphenicol Penicillin, Penicillin G Ampicillin, Gentamicin, Kanamycin, Cefotaxime, Cefoxitin, Vancomycin, Acyclovir, Amikacin, Ceftazidime, Ceftriaxone, Cefazolin |

| Definitions   | Instructions                | Sources  | Keywords and abbreviations  |
|---|-----------------------------|--|---|
| 36. Abnormal conditions of the newbor   | n (BC #54)—Continued        |  |   |
| Seizure or serious neurologic dysfunction Seizure is any involuntary repetitive, convulsive movement or behavior. Serious neurologic dysfunction is severe alteration of alertness. Excludes: - Lethargy or hypotonia in the absence of other neurologic findings - Symptoms associated with CNS congenital anomalies |                             | 1st Newborn H&P  2nd Physician progress notes <i>under</i> — Neuro examination             | Seizures Tonic/Clonic/Clonus Twitching Eye rolling Rhythmic jerking Hypotonia Obtundation Stupor Coma HIE-Hypoxic-ischemic encephalopathy |
| 37. Congenital anomalies of the newbor  | en (BC #55)                 |  |   |
| Malformations of the newborn diagnosed prenatally or after delivery.  | Check all boxes that apply. |  |   |
| Anencephaly Partial or complete absence of the brain and skull.  Also called anencephalus, acrania, or absent brain.  Also includes infants with craniorachischisis (anencephaly with a contiguous spine defect).   |                             | 1st Labor and delivery summary record <i>under</i> –Infant data  2nd Newborn admission H&P | Anencephalus Acrania Absent brain Craniorachischisis Exencephaly Hydraencephaly   |

| Definitions  | Instructions | Sources                             | Keywords and abbreviations  |
|--|--------------|-------------------------------------|---|
| 37. Congenital anomalies of the newborn (BC #55)—Continued   |              |                                     |   |
| Meningomyelocele/Spina bifida Spina bifida is herniation of the meninges or spinal cord tissue through a bony defect of spine closure. |              | See Anencephaly                     | Meningocele<br>Rachischisis   |
| Meningomyelocele is herniation of meninges and spinal cord tissue.   |              |                                     |   |
| Meningocele (herniation of meninges without spinal cord tissue) should also be included in this category.                              |              |                                     |   |
| Both open and closed (covered with skin) lesions should be included.   |              |                                     |   |
| <u>Do not include</u> spina bifida occulta (a midline bony spinal defect without protrusion of the spinal cord or meninges).           |              |                                     |   |
| Cyanotic congenital heart disease Congenital heart defects that cause cyanosis.  |              | 1st Physician progress notes under— | TGA-Transposition of the great arteries TOF-Tetralogy of Fallot Pulmonary or pulmonic valvular atresia Tricuspid atresia Truncus arteriosus TAPVR-Total/partial anomalous pulmonary venous return with or without obstruction COA-Coarctation of the aorta HLHS-Hypoplastic left heart syndrome IAA-Interrupted aortic arch |

| Definitions   | Instructions         | Sources  | Keywords and abbreviations |
|---|----------------------|--|----------------------------|
| 37. Congenital anomalies of the newbor  | n (BC #55)—Continued |  |                            |
| Congenital diaphragmatic hernia  Defect in the formation of the diaphragm allowing herniation of abdominal organs into the thoracic cavity.   |                      | 1st Infant H&P  2nd Labor and delivery summary record <i>under</i> —Infant data  |                            |
| Omphalocele A defect in the anterior abdominal wall in which the umbilical ring is widened, allowing herniation of abdominal organs into the umbilical cord.  The herniating organs are covered by a nearly transparent membranous sac (different from gastroschisis [see below]), although this sac may rupture. Also called exomphalos.  Do not include umbilical hernia (completely covered by skin) in this category. |                      | 1st Labor and delivery summary record <i>under</i> —Infant data  2nd Admission history and physical (H&P) <i>under</i> —G.I. | Exomphalos                 |
| Gastroschisis  An abnormality of the anterior abdominal wall, lateral to the umbilicus, resulting in herniation of the abdominal contents directly into the amniotic cavity. Differentiated from omphalocele by the location of the defect and the absence of a protective membrane.  |                      | See Omphalocele  |                            |

| Definitions   | Instructions   | Sources  | Keywords and abbreviations   |
|---|--|--|--|
| 37. Congenital anomalies of the newborn (BC #55)—Continued  |  |  |  |
| Limb reduction defect excluding congenital amputation and dwarfing syndromes.  Complete or partial absence of a portion of an extremity, secondary to failure to develop. |  | 1st Labor and delivery summary record <i>under</i> —Infant data  2nd Newborn H&P | Look for: Amniotic bands ABS-Amniotic band syndrome  |
| Cleft lip with or without cleft palate Incomplete closure of the lip. May be unilateral, bilateral, or median.  |  | See Limb reduction defect  | Cleft lip (unilateral, bilateral, or median)   |
| Cleft palate alone Incomplete fusion of the palatal shelves. May be limited to the soft palate or may extend into the hard palate.  |  | See Limb reduction defect  |  |
| Cleft palate in the presence of cleft lip should be included in the category above.   |  |  |  |
| Down syndrome Trisomy 21–A chromosomal abnormality caused by the presence of all or part of a third copy of chromosome 21.  Karyotype confirmed Karyotype pending         | Check if a diagnosis of Down syndrome, Trisomy 21 is confirmed or pending. | 1st Infant progress notes 2nd Genetic consult                                    | Trisomy 21 Positive (confirmed) Possible Down (pending) Rule out (R/O) Down (pending) Trisomy 21 mosaicism |

| Definitions  | Instructions  | Sources   | Keywords and abbreviations  |
|--|---|---|---|
| 37. Congenital anomalies of the newbor   | rn (BC #55)—Continued   |   |   |
| Suspected chromosomal disorder Includes any constellation of congenital malformations resulting from or compatible with known syndromes caused by detectable defects in chromosome structure.  Karyotype confirmed Karyotype pending | Check if a diagnosis of a suspected chromosomal disorder is confirmed or pending. (May include Trisomy 21.) | See Down syndrome   | Trisomy and then a number such as: 13–Patau's syndrome 17 or 18–Edward syndrome Positive (confirmed) Possible trisomy(pending) Rule out (R/O) (pending) |
| Hypospadias Incomplete closure of the male urethra resulting in the urethral meatus opening on the ventral surface of the penis. Includes:   |   | 1st Labor and delivery summary under—Infant data  2nd Newborn H&P under— Genitourinary (GU) |   |
| <ul> <li>First degree (on the glans ventral to the tip)</li> <li>Second degree (in the coronal sulcus)</li> <li>Third degree (on the penile shaft)</li> </ul>  |   |   |   |

| Definitions   | Instructions   | Sources                                     | Keywords and abbreviations |
|---|--|---|----------------------------|
| 38. Was the infant transferred within 2   | 4 hours of delivery? (BC #56)  |   |                            |
| Transfer status of the infant from this facility to another within 24 hours after delivery. | Check "yes" if the infant was transferred from this facility to another within 24 hours of delivery.  Enter the name of the facility to which the infant was transferred.  If the name of the facility is not known, enter "unknown."  If the infant was transferred more than once, enter the name of the first facility to which the infant was transferred. | 1st Infant progress notes 2nd Transfer form | Look for: Disposition      |
| 39. Is infant living at time of the report  | ? (BC #57)   |   |                            |
| Information on the infant's survival.   | Check "yes" if the infant is living.  Check "yes" if the infant has already been discharged to home care.  Check "no" if it is known that the infant has died. If the infant has died, make sure that a death certificate is filed.  If the infant was transferred and the status is known, indicate the known status.   | 1st Infant progress notes                   |                            |

| Definitions   | Instructions  | Sources   | Keywords and abbreviations  |
|---|---|---|---|
| 40. Is the infant being breastfed at disc   | harge? (BC #58)   |   |   |
| Information on whether the infant was receiving breastmilk or colostrum during the period between birth and discharge from the hospital.  Breastfeeding refers to the establishment of breastmilk through the action of breastfeeding or pumping (expressing).  Include any attempt to establish breastmilk production during the period between birth and discharge from the hospital.  Include if the infant received formula in addition to being breastfed.  Does not include the intent to breastfeed. | Check "yes" if the infant was breastfed at any time before being discharged from the hospital.  Check "no" if the infant was not breastfed before being discharged from the hospital. | 1st Labor and delivery summary record <i>under</i> —Infant data  2nd Maternal progress note  3rd Newborn flow record <i>under</i> —Feeding  4th Lactation consult | Pumping Lactation consultation LATCH score (Latch on, Audible swallow, Type of nipple, Comfort, and Help—used to measure position and attachment of the baby on the breast) Breast pump Breast pump protocol Breast milk MM-Mother's milk FBM-Fresh breast milk Attempt to breastfeed |
| 41. Method of disposition* (FDFWS #3  | 2, FDR #13)   |   | I   |
| Burial Cremation Hospital disposition Donation Removal from state Other (specify) * Applicable to fetal deaths only.  | Check only one method.  | 1st Labor and delivery summary record <i>under</i> —Infant data  2nd Nursing note  3rd Attending death note  4th Social work note                                 |   |

The use of trade names is for identification only and does not imply endorsement by the Centers for Disease Control and Prevention,

U.S. Department of Health and Human Services.

## **Index of Items**

| $\mathbf{A}$   | County of birth  |
|--|--|
| Abnormal conditions of the newborn   | Cyanotic congenital heart disease (Congenital anomalies of the newborn) 42 |
| Admission to an intensive care unit (Maternal morbidity)   | D  |
| Anencephaly (Congenital anomalies of the newborn)  |  |
| Antibiotics received by the mother during delivery   | Date certified   |
| (Characteristics of labor and delivery)  |  |
| Antibiotics received by the newborn for suspected neonatal sepsis (Abnormal conditions of the newborn) | Date of birth  |
| Assisted reproductive technology (Risk factors in this pregnancy) 19                                   | Date of last live birth  |
| Assisted ventilation required for more than six hours  | Date of last other pregnancy outcome                                       |
| (Abnormal conditions of the newborn)   | Diabetes (Risk factors in this pregnancy)                                  |
| Assisted ventilation required immediately following delivery   | Prepregnancy (Risk factors in this pregnancy)                              |
| (Abnormal conditions of the newborn)   | Gestational (Risk factors in this pregnancy)                               |
| Attendant's name, title, and ID  | Down syndrome (Congenital anomalies of the newborn)                        |
| Augmentation of labor (Characteristics of labor and delivery)  | _  |
| Apgar score  | E  |
| В  | Eclampsia (Risk factors in this pregnancy)                                 |
|  | Epidural or spinal anesthesia during labor (Characteristics of             |
| Birthweight or weight of fetus   | labor and delivery)  |
| C  | External cephalic version (Obstetric procedures)                           |
| Certifier's name and title   | $\mathbf{F}$   |
| Characteristics of labor and delivery  | Facility ID  |
| Chlamydia (Infections present and/or treated during this pregnancy) 21                                 | Facility name  |
| City, town, or location of birth   | Fertility-enhancing drugs, artificial insemination, or                     |
| Cleft lip with or without cleft palate (Congenital anomalies of the newborn) 44                        | intrauterine insemination (Risk factors in this pregnancy)                 |
| Cleft palate alone (Congenital anomalies of the newborn)   | Fetal presentation at birth (Method of delivery)                           |
| Clinical chorioamnionitis diagnosed during labor or maternal   | Final route and method of delivery (Method of delivery)                    |
| temperature ≥ 38°C (100.4°F) (Characteristics of labor and delivery) 29                                |  |
| Congenital anomalies of the newborn  | G  |
| Congenital diaphragmatic hernia (Congenital anomalies of the newborn) 43                               | Gastroschisis (Congenital anomalies of the newborn)                        |

| Gonorrhea (Infections present or treated during this pregnancy) 20          | N   |
|---|---|
| Н   | Newborn given surfactant replacement therapy (Abnormal conditions of the newborn) |
| Hepatitis B (Infections present and/or treated during this pregnancy) 21    | NICU admission (Abnormal conditions of the newborn)                               |
| Hepatitis C (Infections present and/or treated during this pregnancy) 21    | Number of other pregnancy outcomes  |
| Hypertension (Risk factors in this pregnancy)                               | Number of previous live births now dead   |
| Prepregnancy (Risk factors in this pregnancy)                               | Number of previous live births now living   |
| Gestational (Risk factors in this pregnancy)                                |   |
| Hypospadias (Congenital anomalies of the newborn)                           | 0   |
| I   | Obstetric estimate of gestation at delivery                                       |
| If cesarean, was a trial of labor attempted? (Method of delivery)           | Obstetric procedures  |
| If not a single birth, number of infants in the delivery born alive         | Omphalocele (Congenital anomalies of the newborn)                                 |
| If not a single birth, order born in the delivery                           | P   |
| Induction of labor (Characteristics of labor and delivery)                  |   |
| Infant's medical record number  | Place where birth occurred (Birthplace)   |
| Infections present and/or treated during this pregnancy                     | Plurality   |
| Is the infant being breastfed at discharge?                                 | Pregnancy resulted from infertility treatment (Risk factors in this pregnancy)    |
| Is the infant living at the time of the report?                             | Previous preterm births (Risk factors in this pregnancy)                          |
|   | Principal source of payment   |
| L   |   |
| Limb reduction defect (Congenital anomalies of the newborn)                 | R   |
| NA.   | Risk factors in this pregnancy  |
| M   | Ruptured uterus (Maternal morbidity)  |
| Maternal morbidity  | $\mathbf{S}$  |
| Maternal transfusion (Maternal morbidity)                                   |   |
| Meningomyelocele/Spina bifida (Congenital anomalies of the newborn) 42      | Seizure or serious neurologic dysfunction (Abnormal conditions of the newborn)    |
| Method of delivery  | Sex of child  |
| Method of disposition   | Steroids (glucocorticoids) for fetal lung maturation received by                  |
| Mother  | the mother before delivery (Characteristics of labor and delivery)                |
| Mother had a previous cesarean delivery (Risk factors in this pregnancy) 19 | Suspected chromosomal disorder (Congenital anomalies of the newborn) 4:           |
| Mother's weight at delivery   | Syphilis (Infections present and/or treated during this pregnancy)                |

## Page 50

Guide to Completing the Facility Worksheet for the Certificate of Live Birth and Report of Fetal Death

| I |   |  |
|---|---|--|
|   | Third- or fourth-degree perineal laceration (Maternal morbidity)                                    |  |
|   | Time of birth   |  |
|   | Total number of prenatal care visits for this pregnancy   |  |
| U |   |  |
|   | Unplanned hysterectomy (Maternal morbidity)   |  |
| V | V   |  |
|   | Was the infant transferred within 24 hours of delivery?   |  |
|   | Was the mother transferred to this facility for maternal medical or fetal indications for delivery? |  |