

# MetReader: Documentation and User's Guide

Hans F. Schwaiger

June 26, 2018

## 1 Introduction

MetReader is a library written in fortran 90 that provides an interface to numerical weather prediction (NWP) data, or other forms of meteorological data, such as radiosonde or other 1-D data.

This library was originally written as a component of the USGS volcanic ash transport and dispersion model, Ash3d. However, since it is useful for programs other than Ash3d, this interface to NWP files is provided as a separate repository that can either be compiled as a library or simply compiled directly with other source code.

NWP data are generally made available by agencies (NCEP, NOAA, NASA, etc.) in a variety of formats (NetCDF, GRIB1, GRIB2, ASCII); each product having its own data structure, naming convention, units, etc. This library isolates the calling program from the peculiarities of interfacing with a particular NWP product. Data can be returned to the calling program on the native grid of the NWP product, or on any grid needed by the calling program. Projection and interpolation of NWP data to the required grid, along with any rotation of velocity vectors to grid-relative, is calculated internally by MetReader.

## 2 Installation

This library requires two additional libraries made available on GitHub and USGS GitLab:

- HoursSince  
This is a library that takes a date and time, and calculates the number of hours since January 1, 1900 (or any arbitrary base year).  
(<https://github.com/hschwaiger-usgs/HoursSince>)
- projection  
This is a library that calculates commonly-used projected coordinates.  
(<https://github.com/hschwaiger-usgs/projection>)

These libraries are installed by default into `/opt/USGS`. If they are installed elsewhere, then you will need to edit the makefile to point to your installation. Additionally, the default makefile will build MetReader with both NetCDF and GRIB2 enabled. If either of these libraries are unavailable on your system, you can deactivate those options by setting the

corresponding flags to F in the makefile. On a RHEL/CentOS system, you can install the data format libraries by typing

```
sudo yum install netcdf netcdf-devel grib_api grib_api-devel
```

or for Ubuntu

```
sudo apt-get install libnetcdf* libgrib-api*
```

To compile MetReader as a library, simple type:

```
make all
```

This will build the requested components of the library. If GRIB2 is enabled, this will also build the GRIB2 indexer, `gen_GRIB2_index`. This is a tool that generates an index file of the GRIB records which speeds access time to individual records substantially.

To install the library, module files and tools, edit the `INSTALLDIR` variable of the makefile (the default is `/opt/USGS`) and type:

```
make install
```

This will also install scripts that can be used to download the 0.5 degree GFS forecast files and the NCEP 2.5-degree Reanalysis files.

You will need to have write permission in `/${INSTALLDIR}` or install as root. Assuming the libraries `projection` and `hourssince` are installed in the default location, installation will result in the following directory structure:

```
/opt/USGS
|-- bin
|   |-- autorun_scripts
|   |   |-- autorun_gfs0.5deg.sh
|   |   |-- autorun_NCEP_50YearReanalysis.sh
|   |   |-- convert_gfs0.5deg.sh
|   |   |-- get_gfs0.5deg.sh
|   |   |-- get_NCEP_50YearReanalysis.sh
|   |   |-- grib2nc.sh
|   |-- gen_GRIB2_index
|   |-- HoursSince1900
|   |-- ncMet_check
|   |-- ncMetSonde
|   |-- ncMetTraj_B
|   |-- ncMetTraj_F
|   |-- yyyymmddhh_since_1900
|-- include
|   |-- metreader.mod
|   |-- projection.mod
'-- lib
    |-- libhourssince.a
    |-- libMetReader.a
    |-- libprojection.a
```

## 3 Usage

MetReader can read a variety of input data formats (1-D, 3-D, ASCII, NetCDF, GRIB2) and provide access to the meteorological data on either the native grid of the NWP data, or interpolated onto a finer regular grid. Before the meteorological data can be accessible, several preliminary steps must first be completed.

### 3.1 Preliminary meta-data

The calling program needs to prepare several parameters that define the type of meteorological data to be read.

- `iw` :: windfile class (the distinction between 3 and 4 will be obsolete soon)
  - 1 : 1-D wind sounding
  - 2 : 3-D grid is read from a ASCII file
  - 3 : single, multistep 3-D datafile
  - 4 : multiple 3-D datafiles
  - 5 : special case for NCEP 50-year reanalysis and NOAA reanalysis
- `iwf` :: windfile format number (linked to specific product in Table 8.2 of Appendix 8.2)
- `igrid` :: NCEP Grid ID
- `idf` :: Data file format code (ASCII, NetCDF, GRIB, etc)
  - 1 : ASCII
  - 2 : NetCDF
  - 3 : GRIB Edition 2
  - 4 : GRIB Edition 1 (Not yet implemented; used by NARR and NOAA-CIRES)
- `iwfiles` :: number of windfiles to be read
- `SimStartHour` :: The starting time needed given as hours since 1900-01-01 00:00Z  
This can alternatively be given as hours since a different base year as described below.
- `SimDuration` :: duration needed in hours

Additionally, there are several other parameters that could be set, if needed.

- `MR_BaseYear` :: The default value for this is 1900, but must be reassigned by the calling program if the NWP files to be read contain data prior to 1900.
- `MR_useLeap` :: The default is `.true.` but can be set to `.false.` if the NWP files use a calendar without leap years (e.g. some paleoclimate CAM files)
- `MR_iHeightHandler` :: This code defines how MetReader behaves when values at altitudes above those available in the NWP files are requested. Default value is 2.

- 1 : Stop the program
  - 2 : Return wind values equal to the value at the highest available node; temperature values remain constant from 11-20 km, then increase by 2° C/km.
- `MR_iwf_template` :: This is the file name of the template for the custom NetCDF files described in section 5. This is only required to be set if `iwf=0`. Currently, the custom windfile specification is only implemented for `idf=2` (NetCDF).
  - `call MR_Reset_Memory` :: If MetReader had been initialized and used for one set of NWP files, but later during the execution of the calling program, a subsequent set of NWP files is needed, this subroutine can be called to free all allocated memory.

**Step 1:** The space needed for storing the NWP file meta-data must be allocated.

```
call MR_Allocate_FullMetFileList(iw,iwf,igrid,idf,iwfiles)
```

This allocates the variable `MR_windfiles(1:MR_iwindfiles)`.

**Step 2:** The calling program must now fill the list of file names `MR_windfiles()` with each name not exceeding 130 characters in length. For the special cases for the NCEP/NCAR Reanalysis 1 or the NOAA-CIRES 20th Century Reanalysis (`iwf=25` or `27`), MetReader expects a specific structure to the directory holding the NWP files. For these reanalysis product, the list of files is only one string (`iwfiles=1`) with the ‘file’ given as the root directory to the NWP files. For example, for the NCEP/NCAR Reanalysis 1:

```
iwfiles = 1
MR_windfiles(1) = '/data/WindFiles/NCEP'
```

where the files are stored as follows:

```
/data/WindFiles/NCEP
|-- 2016
|   |-- air.2016.nc
|   |-- hgt.2016.nc
|   |-- omega.2016.nc
|   |-- shum.2016.nc
|   |-- uwnd.2016.nc
|   '-- vwnd.2016.nc
|-- 2017
|   |-- air.2017.nc
|   |-- hgt.2017.nc
|   |-- omega.2017.nc
|   |-- shum.2017.nc
|   |-- uwnd.2017.nc
|   '-- vwnd.2017.nc
```

Similarly, for the NOAA-CIRES 20th Century Reanalysis product:

```
iwfiles = 1
MR_windfiles(1) = '/data/WindFiles/NOAA'
```

where the files are stored as follows:

```
/data/WindFiles/NOAA
|-- 2008
|   |-- GRIB
|   |   |-- pgrbanl_mean_2008_HGT_pres.grib
|   |   |-- pgrbanl_mean_2008_RH_pres.grib
|   |   |-- pgrbanl_mean_2008_TMP_pres.grib
|   |   |-- pgrbanl_mean_2008_UGRD_pres.grib
|   |   |-- pgrbanl_mean_2008_VGRD_pres.grib
|   |   |-- pgrbanl_mean_2008_VVEL_pres.grib
|   |-- pgrbanl_mean_2008_HGT_pres.nc
|   |-- pgrbanl_mean_2008_RH_pres.nc
|   |-- pgrbanl_mean_2008_TMP_pres.nc
|   |-- pgrbanl_mean_2008_UGRD_pres.nc
|   |-- pgrbanl_mean_2008_VGRD_pres.nc
|   |-- pgrbanl_mean_2008_VVEL_pres.nc
|-- 2009
|   |-- ...
```

Note that the GRIB files are in version 1 format and cannot yet be directly read by MetReader. They must first be converted to NetCDF and placed as described above.

**Step 3:** Once the names of the NWP files are specified, the files must be queried to determine the scope of the available data.

```
call MR_Read_Met_DimVars([iyear])
```

The `iyear` argument is optional, but is expected for `iwf=25` or `27` since MetReader needs to know whether to allocate space for 365 or 366 days. Once the subroutine is called, each of the NWP files is checked for existence, then the following public variables are set:

- All the projection parameters for the NWP grid.
- `Met_dim_names()` :: The names of the dimensions in the file in the order of: time, pressure1, y, x, pressure2 (for Vz), pressure3 (for RH)
- `Met_var_names()` :: The names of the variables in the file according to the Table 8.1 of Appendix 8.1:
- `Met_var_conversion_factor()` :: For each of the variables, if the units is not as listed above, then the corresponding factor is set that converts to the expected units. For example, if precipitation rate is given in mm/hr, then the conversion factor to the expected units (kg/m<sup>2</sup>s) is 1/3600 or  $2.778 \times 10^{-4}$ .

- `Met_var_IsAvailable()` :: set to `.true.` if the NWP file provides this variable.
- `nx_fullmet,ny_fullmet,np_fullmet` :: lengths of grid for x (lon), y (lat) and p
- `x_fullmet_sp,y_fullmet_sp,p_fullmet_sp` :: values of the grid in km (degrees) and hPa
- `IsLatLon_MetGrid` :: `.true.` if the grid is specified in longitude and latitude
- `IsGlobal_MetGrid` :: `.true.` if the grid is periodic in longitude. (Note: mapping across the poles is not yet implemented).
- `IsRegular_MetGrid` :: `.true.` if dx, dy (dlon, dlat) is constant

**Step 4:** Next, the calling program needs to define the type of grid onto which MetReader will interpolate values. Namely, whether or not the computational grid is projectected, and if so, which projection and associated values. This grid used by the calling program can be independent of the grid used by the NWP files. The parameters in the list for the subroutine below are the values needed by the `libprojection.a` library. Below is an example call from `Ash3d`.

```
call MR_Set_CompProjection(IsLatLon,A3d_iprojflag,A3d_lam0,      &
                          A3d_phi0,A3d_phi1,A3d_phi2,        &
                          A3d_k0_scale,A3d_radius_earth)
```

If `IsLatLon=.true.` then all other projection parameters are ignored. Similarly, some of the parameters are required and others not, depending on the projection used. Currently, `libprojection.a` has implemented Polar stereographic (`iprojflag=1`), Albers Equal Area (`iprojflag=2`), UTM (`iprojflag=3`), Lambert conformal conic (`iprojflag=4`), and Mercator (`iprojflag=5`).

**Step 5:** The final preparatory step is to evaluate the windfiles provided with the spatial and temporal requirements of the calling program.

```
call MR_Initialize_Met_Grids(nxmax,nymax,nzmax,              &
                             x(1:nxmax),y(1:nymax),z(1:nzmax), &
                             IsPeriodic)
```

`nxmax,nymax,nzmax` are the sizes of the computational grid. `x,y,z` are the single-precision grid values in km (or degrees). `IsPeriodic` indicates whether or not the computational grid is periodic in x. This subroutine also determines the size of the sub-grid of the full NWP data that is needed by the calling program.

```
call MR_Set_Met_Times(SimStartHour, SimDuration)
```

Here `SimStartHour` and `SimDuration` give the initial time (in hours since 1900 (or other base year)) and the length of time needed in hours. This subroutine opens all the files listed in `MR_windfiles()` and verifies that the files provided cover the requested time span. Then a

list is prepared of all the time steps and files needed to fully cover the requested time with the step index (`istep=1`) corresponding to the step at or just prior to `SimStartHour`. The hours between time steps is saved in `MR_MetStep_Interval()` and does not need to be constant.

### 3.2 Reading meteorological data

When handling data from the NWP files, there are three grids that MetReader uses:

- `metP` :: the native grid of the NWP file clipped to span the computational needs (possible re-ordered)
- `metH` :: the horizontal grid of the NWP file, but with the vertical coordinate mapped from the pressure grid of the NWP file to the altitude grid of the computational grid.
- `compH` :: the full x,y,z of the computational grid.

Once the computational grid is specified and the sub-grid of the NWP is determined in step 5 above, the following public work spaces are allocated.

```
MR_dum2d_met_int(nx_submet,ny_submet)      :: integer
MR_dum2d_met(nx_submet,ny_submet)         :: float
MR_dum3d_metP(nx_submet,ny_submet,np_fullmet) :: float
MR_dum3d2_metP(nx_submet,ny_submet,np_fullmet) :: float
MR_dum3d_meth(nx_submet,ny_submet,nz_comp)  :: float
MR_dum2d_comp_int(nx_comp,ny_comp)         :: integer
MR_dum2d_comp(nx_comp,ny_comp)            :: float
MR_dum3d_compH(nx_comp,ny_comp,nz_comp)    :: float
```

`nx_comp`, `ny_comp`, `np_comp` are the number of nodes in the computational grid and `nx_submet`, `ny_submet`, `np_submet` are the number of nodes of the NWP grid that is needed to span the required computational grid.

To read data from the meteorological files, the fundamental subroutine is

```
call MR_Read_3d_MetP_Variable(ivar,istep)
```

where `ivar` is the code for the variable of interests (see Table 8.1) and `istep` is the index of the sequence of time steps set up based on `SimStartHour` and `SimDuration` in Step 5 above. This subroutine provides a common interface regardless of the data format (ASCII, NetCDF, GRIB2, etc.), data structure (latitude might be top-to-bottom or bottom-to-top, pressure might be surface-to-top or top-down, grid might be staggered, longitude might be  $-180 \rightarrow 180$  or  $0 \rightarrow 360$ ). In all cases for 3-D variables, what is returned is the public work array `MR_dum3d_metP`. This array is consistently ordered such that `x` and `y` (or `lon` and `lat`) increase with index (North to South NWP grids are flipped) and with `p` ordered from the surface to top (pressure is high to low).

If 3-D data is needed on the `metH` grid,

```
call MR_Read_3d_Meth_Variable(ivar,istep)
```

can be used which first populates `MR_dum3d_metP`, then interpolates onto `MR_dum3d_metH`.

For simply reading meteorological data onto the computational grid, use

```
call MR_Read_3d_Met_Variable_to_CompGrid(ivar,istep,[IsNext])
```

This subroutine reads the variable `ivar` from the NWP file into `MR_dum3d_metP`, then interpolates these values onto `MR_dum3d_metH`, finally interpolating onto `MR_dum3d_compH`. The optional argument `IsNext` is used when velocities are used at different points in the program and need to be saved. `IsNext` indicates that the velocity values `MR_v[x,y]_metP_next` should be copied to `MR_v[x,y]_metP_last`.

To interpolate 3-D data onto the altitude levels of the computational grid, first the geopotential height data must be read. This could be achieved by the subroutine `MR_Read_3d_MetP_Variable` described above, but as a simulation advances, we typically need to have the geopotential height data for the previous and the next time steps so that values can be interpolated between steps. The subroutine

```
call MR_Read_HGT_arrays(istep,[reset_first_time])
```

is a specialized reader for `ivar=1`. The first time this is called, both `istep` and `istep+1` are called, populating the two public variables `MR_geoH_metP_last` and `MR_geoH_metP_next`. Subsequent calls copy `MR_geoH_metP_next` to `MR_geoH_metP_last` and reads the next values from `istep+1`. If the optional argument `reset_first_time` is given, both `istep` and `istep+1` are directly read.

Similar read subroutines are available for 2-D variables.

```
call MR_Read_2d_Met_Variable(ivar,istep)
```

Given a 2-D variable identified with `ivar`, this populates the public work array `MR_dum2d_met`, where again, `1:nx_subset,1:ny_subset` corresponds to the subset of nodes of the NWP file that is needed to span the computational grid.

```
call MR_Read_2d_Met_Variable_to_CompGrid(ivar,istep)
```

can be called similar to `MR_Read_2d_Met_Variable` which directly reads the variable needed and returns the values interpolated onto the public work space `MR_dum2d_comp`.

### 3.2.1 Regridding meteorological data

Internal to `MetReader` are a variety of regridding subroutines to interpolate from the `metP` grid onto `metH` or `compH`. In some circumstances, data is initially needed just on the native NWP grid, but then derived values might be required on the full computational grid. For example, if air viscosity is needed for a particle fall model, temperature and pressure values on the NWP grid can be read. Then viscosities can be calculated for each of the corresponding nodes of the NWP grid. To interpolate these values from the `metP` grid to the `compH` grid, the subroutine

```
call MR_Regrid_MetP_to_CompGrid(istep)
```



can be used. This takes the variable `MR_dum3d_metP` and interpolates it onto `MR_dum3d_compH`. Similarly,

```
call MR_Regrid_MetP_to_MetH(istep)
```

interpolates `MR_dum3d_metP` to `MR_dum3d_metH` and

```
call MR_Regrid_Met2d_to_Comp2d
```

interpolates `MR_dum2d_met` to `MR_dum2d_comp`.

### 3.2.2 Wind velocity vectors

In some cases, the wind velocity vectors need to be mapped from the coordinate system of the NWP files to the computational grid. Velocity components on the computational grid are always provided as grid-relative. For some NWP files, such as the North American Regional Reanalysis (NARR), the NWP grid is projected, yet the wind components are provided as Earth-relative. In other cases, if the NWP data are provided on a projected grid, but the computational grid is either lon/lat or on a different projection, the grid-relative, projected velocity components must be converted to that needed for the computational grid.

If the wind data are provided as grid-relative on a projected grid and something else is needed, the subroutine

```
call MR_Rotate_UV_GR2ER_Met(MR_iMetStep_Now)
```

will read both U and V components and decompose the vector into earth-relative Easterly and Northerly wind components with values at the `metP` grid nodes. These components are put in the variables `MR_u_ER_metP` and `MR_v_ER_metP`. Once we have Earth-relative velocity components (either from direct read of lon/lat data, from the subroutines `MR_Rotate_UV_GR2ER_Met` or from direct read of NARR data), if the computational grid is not Earth-relative, then these Earth-relative components can be subsequently decomposed into grid-relative components on the computational grid with

```
call MR_Rotate_UV_ER2GR_Comp(MR_iMetStep_Now)
```

This subroutine returns the U and V components on the `compH` grid through the two dummy work space variables `MR_dum3d_compH` and `MR_dum3d_compH_2`

### 3.3 Additional Functions

**Horizontal derivatives** MetReader has subroutines for calculating horizontal derivatives of variables on the `metP` grid. This is used for calculating velocity gradients for diffusivity calculations, but could be applied to any variable. The two subroutines are `MR_De1MetP_Dx` and `MR_De1MetP_Dy`. Both read the values from `MR_dum3d_metP` and return derivative values on `MR_dum3d2_metP`.

**U.S. Standard Atmosphere** MetReader also has several functions for calculating values from the U.S. Standard Atmosphere.

- `MR_Temp_US_StdAtm(zin)` :: Returns a temperature in K given a height in km.
- `MR_Z_US_StdAtm(pin)` :: Returns a height in km given a pressure in hPa.
- `MR_Pres_US_StdAtm(zin)` :: Returns a pressure in hPa given a height in km.

## 4 Supported types of meteorological data

MetReader can read meteorological data in a variety of different formats.

### 4.1 1-D ASCII data

The simplest format for meteorological data is just a 1-D wind profile in ASCII format. To read this type, set `iw = 1`. The particular format of the 1-D data can be specified by the `iwf`, where 1 indicates a user-specified column of values and 2 indicates that the files are from the global radiosonde data available from <https://ruc.noaa.gov/raobs/> or from <http://weather.uwyo.edu>. In both these cases, there is no NCEP grid ID associated with the sonde points so `MR_iGridCode` is interpreted as the number of profile points. Each point at each time must have its own file containing the profile data. `MR_idataFormat` is not needed (ASCII is assumed from `iw = 1`). Profiles at multiple times can be included by repeating the sequence of profile locations at the later times. All points must have profiles for all times and files for subsequent time steps must be given in the same order. Also, all profile files must be in the same format (projection, columns of data). For example:

```
iw      = 1
iwf     = 2
igrd    = 3
MR_iwindfiles = 6      # This must be an even multiple of MR_iGrid
PADQ_2018060100.dat  # 00Z radio sonde at Kodiak, AK
PAKN_2018060100.dat  # 00Z radio sonde at King Salmon, AK
PANC_2018060100.dat  # 00Z radio sonde at Anchorage, AK
PADQ_2018060112.dat  # 12Z radio sonde at Kodiak, AK
PAKN_2018060112.dat  # 12Z radio sonde at King Salmon, AK
PANC_2018060112.dat  # 12Z radio sonde at Anchorage,
```

#### 4.1.1 User-specified profiles

User-specified profiles describe minimally, the horizontal velocity as a function of height at a point (or points) in the domain. Beyond this constraint, the format is fairly general. The format is as follows.

```
L01 string header line
L02 time(hr) nlev [ncol] [ivar(ncol)]
L03 x/lon y/lat [LLflag] [projection parameters]
L04 data
```

```

:
Ln EOF

```

Line 1 is a header line that is ignored

Line 2 contains at least the time of the profiles and the number of lines of data to read. Optionally, the number of columns of data can be specified followed by a sequence of integers specifying the variable ID's (see Table 8.1) for the corresponding columns. If neither of these optional arguments `ncol` `ivar(ncol)` are given, then `MetReader` tests for the number of columns present. If there are 3, then the data are read in as altitude (in m), wind speed (in m/s), and wind direction (in degrees E of N, with the convention of where the wind is coming from). If 5 columns are present, the fourth and fifth columns correspond to pressure (in hPa) and temperature (in °C), respectively. If the optional parameters on line 2 are given with the number of columns of data and corresponding variables, then the variables can be read in any order, but must include some measure of height and some measure of the wind vector. Height can include either altitude (in m), or pressure (in hPa), or both. If only one of these two is given, the other is filled according to the U.S. Standard Atmosphere. The measure of the wind vector can be given either in U and V, or speed and direction. If temperature is not provided, it is calculated from the pressure or altitude using the U.S. Standard atmosphere.

Line 3 contains the coordinate of the 1-D profile. If two values are given, the coordinate system is assumed to be that of the computational grid (either in projected or lon/lat grids). If more than two parameters are given, then the parameters beyond 2 are interpreted to be the list of parameters required by `libprojection.a`. If the `LLflag` is given as a third parameter with a value of 1, then the coordinate is interpreted to be in longitude and latitude. If the value is 0, then the coordinate system is projected with specification defined by subsequent values on line 3.

Line 4 contains the start of `nlev` rows of data  
 Here is an example of the 3-column format.

```

Hanford
2 5 #wind time, number of levels
-120.0 46.20 #Longitude, latitude
0 10.000 90.00
5000 10.000 90.00
10000 10.000 90.00
15000 10.000 90.00
20000 10.000 90.00

```

Here is an example of a user-specified number of columns with data on a projected grid.

```

Hanford
0 5 5 1 2 3 0 5 #wind time, number of levels, nvar ivars(1:nvar)
-2062.26 2606.69 0 4 265.0 25.0 25.0 25.0 6371.229 # (-120.0 46.20) in LCC grid 212 coords
0 -10.000 0.00 1000.0 10.0
5000 -10.000 0.00 560.0 -5.5
10000 -10.000 0.00 262.0 -47.3
15000 -10.000 0.00 126.0 -58.1
20000 -10.000 0.00 58.0 -54.9

```

#### 4.1.2 1-D radiosonde profiles (single or multiple times)

Instead of user-specified profiles, MetReader can also read radiosonde data. Both the global archive as well as the 00Z and 12Z radiosonde data are available from the NOAA Earth System Research Laboratory Rapid Update Cycle Radiosonde Database (<https://ruc.noaa.gov/raobs/>). These data are also archived at the University of Wyoming (<http://weather.uwyo.edu/upperair/sounding.html>). To use data in this format, set `iwf = 2`. From both these sites, radiosonde data are available in a variety of formats. The default format for the UW site is "Text: List"

72694 SLE Salem Observations at 00Z 20 Jun 2018

PRES	HGHT	TEMP	DWPT	RELH	MIXR	DRCT	SKNT	THTA	THTE	THTV
hPa	m	C	C	%	g/kg	deg	knot	K	K	K
1007.0	61	30.6	8.6	25	7.01	345	7	303.1	324.5	304.4
1002.0	110	29.4	9.4	29	7.44	349	8	302.4	324.9	303.7
1000.0	129	29.2	9.2	29	7.35	350	8	302.4	324.6	303.7
980.2	305	27.5	8.5	30	7.17	355	9	302.4	324.1	303.7
946.7	610	24.6	7.4	33	6.85	0	7	302.4	323.2	303.7
925.0	814	22.6	6.6	36	6.65	350	6	302.4	322.6	303.6
914.4	914	21.7	5.7	35	6.33	345	7	302.5	321.8	303.7
908.0	975	21.2	5.2	35	6.14	352	7	302.6	321.3	303.7
:										
:										
:										
13.3	29602	-39.7	-76.7	1	0.08	66	18	802.1	803.1	802.2
11.7	30480	-39.0	-76.4	1	0.10	90	30	834.6	835.8	834.6
10.2	31394	-38.2	-76.2	1	0.12	80	19	869.7	871.2	869.8
10.0	31557	-38.1	-76.1	1	0.12			876.2	877.8	876.2

Station information and sounding indices

```

Station identifier: SLE
Station number: 72694
Observation time: 180620/0000
Station latitude: 44.91
Station longitude: -123.00
Station elevation: 61.0
Showalter index: 6.44

```

```

:
:
:

```

This file contains many pressure levels, but the radiosonde standard include a limited set of "required" pressure levels (1000, 925, 850, 700, 500, 400, 300, 250, 200, 150, and 100 mbar; and supplementally 70, 50, 30, 20, 10 if the probe reaches those heights.). When MetReader reads a file of this format, only these mandatory pressure levels are read. Only the pressure, height,

temperature, wind direction and wind speed are read. After the columnar data are read, the **Station number** is read. From this number (72694 in the above example), the station longitude, latitude and elevation are automatically populated from internally stored station metadata available from [https://ruc.noaa.gov/raobs/General\\_Information.html](https://ruc.noaa.gov/raobs/General_Information.html). The station list is truncated to just what is currently operational and to just current coordinates (some stations have been moved or deactivated). Next the **Observation time** is read.

An alternative format is the FAA604 WMO/GTS format from <https://ruc.noaa.gov/raobs/> or the nearly identical "Text:Raw" format from the University of Wyoming.

```

TTAA 70001 72694 99007 30672 34507 00129 29270 35008 92814 22666 35006 85542
17466 03511 70163 04859 09008 50582 10178 20015 40750 22974 23023 30953 39945
24033 25075 49164 24541 20220 52974 25050 15405 55180 26035 10660 60376 23011
88188 54573 25052 77999 31313 58208 82346 51515 10164 00006 10194 00508 09506
TTBB 70008 72694 00007 30672 11002 29470 22908 21266 33890 20469 44814 15067
55762 10261 66693 04259 77655 00456 88615 02757 99592 05350 11580 05756 22573
05764 33566 06368 44551 07765 55547 07762 66536 08563 77526 08166 88518 08766
99509 09378 11494 10782 22468 14374 33440 16981 44381 25773 55346 31962 66330
34346 77323 35738 88302 39743 99278 44357 11261 47757 22245 49568 33240 49977
44231 49578 55220 51375 66194 53373 77188 54573 88175 54774 99172 53376 11170
52777 22160 54379 33153 54180 44143 56979 55134 56379 66129 57578 77120 57778
88110 60177 99106 59977 11104 60976 22102 59977 31313 58208 82346 41414 12761
PPBB 70008 72694 90012 34507 35509 36007 90346 34507 02009 03008 90789 07002
16003 12508 91246 12505 12011 20008 9179/ 19506 19015 92025 21516 23018 23523
93046 23537 24524 25049 940// 25053 950// 26022
TTCC 70002 72694 70883 60377 11004 50095 57778 11020 30421 52181 07020 20686
45783 08022 88999 77999 31313 58208 82346

```

This format contains nearly the same information as the previous example, but requires a bit of translation. Appendix F of <https://www.ofcm.gov/publications/fmh/FMH3/00-entire-FMH3.pdf> gives the full specification. If the string "TTAA" is found in the ASCII file, this format is attempted first, otherwise the "Text:List" is assumed. This WMO/GTS format is intended for the Rapid Update Cycle data and does not contain the year or month (only the day-of-month and hour) of the data. If this format is detected, the year and month of the at execution time is assumed unless otherwise specified by the calling program.

## 4.2 Network of 1-D radiosonde profiles

In most of the world, the radiosonde launch sites are too dispers to form a meaningful network across which we can interpolate values. The radiosonde data is most useful for the regions of interest proximal to launch sites. Currently, the use of multiple sonde locations is not implemented, but simple interpolation using a small number of sonde points is planned for future versions of this library.

## 4.3 3-D forecast or reanalysis data

MetReader's preferred format for 3-D forecast and reanalysis files is NetCDF. Many products are available directly in NetCDF. format such as NCEP/NCAR Reanalysis 1 or NASA

GEOS-5. These products generally require NetCDF version 4. Most of the NCEP forecast products are provided in GRIB2 format. To convert these forecast products to NetCDF format, MetReader relies on the NetCDF. java package (<https://www.unidata.ucar.edu/software/thredds/current/netcdf-java/>). The conversion script in the `autorun_scripts` folder are linked to `netcdfA11-4.5` however any recent version should work. Older versions create NetCDF files with different variable names.

Many of the forecast and reanalysis products are built in to MetReader where the grid geometry, variable list, and pressure levels are preset. These include all the `iwf` numbers listed in section 3.1. If the products change such that the pressure levels differ or if variable names change, then the source code will need to be modified to account for the changes. Alternatively, a template file could be used as described in section 5 below.

MetReader can also read GRIB2 files directly for certain forecast products, including the NAM 196 grid over HI (`iwf=11`), the NAM 91 grid over AK (`iwf=13`) and the GFS 0.25° and 0.5° products (`iwf=22,20`). To read these GRIB2 files, an index file containing the locations of all the records must be created. This index file can be created by the utility `gen_GRIB2_index`, or if the file does not exist at run-time, can be generated through the subroutine call `MR_Set_Gen_Index_GRIB(filename)`. In practice, if windfiles will be used more than a few times, it is generally worthwhile to convert them to NetCDF prior to usage by MetReader. The current implementation of the GRIB2 reader will read each of the required GRIB2 records into memory, decompress the full gridded data, then extract the needed sub-grid; whereas the NetCDF interface allows a more natural direct access to sub-grids. For high resolution, global grids where only a small sub-grid is needed, the impact of reading and uncompressing full grids will be noticeable.

## 5 Custom meteorological file specification

As an alternative to using a particular forecast or reanalysis product with hard-coded file structure, a template file can be used that allows a custom specification of a NetCDF file. The custom template is indicated by `iwf=0` and setting `MR_iwf_template` to the name of the template file. The template file has the following format.

Line 1: The line of projection parameters as expected by `libprojection.a`. e.g.

```
Polar-stereographic  Grid 91   0 1 -150.0 90.0 0.933 6371.229
Polar-stereographic  Grid 216  0 1 -135.0 90.0 0.933 6371.229
Mercator             Grid 196  0 5 198.475 20.0 0.933 6371.229
Lambert Conformal    Grid 212  0 4 265.0 25.0 25.0 25.0 6371.229
Lambert Conformal    Grid 221  0 4 -107.0 50.0 50.0 50.0 6371.229
```

Line 2: `StepInterval [T/F] ::` time interval in hours, and flag indicating whether or not to use leap years (Assumed T if not provided)

Line 3: `ndims nvars ::` number of dimensions and variable that will be listed

Line 4-: list of dimensions where each row specifies the information about the dimension. For example

```
d      1  1.6667e-2 time           hours since
d      2      100.0 lev            Pa
```

In this example, the `d` signifies that the row describes a dimension, `1` is the dimension number, `1.6667e-2` is a scaling factor, and `time` is the name of the dimension. The dimensions should be listed in the following order: time, pressure used for primary state variables (velocity, temperature), y (or longitude), x (or latitude), pressure used for vertical velocities, and pressure used for moisture variables. Additional dimensions can be listed to support other specific variables. The scaling factor is the factor needed to convert the dimension from the units provided to the units required. In this example, the time values are supplied in minutes, and must be multiplied by `1.6667e-2` to be in the expected units of hours. Similarly, pressure is supplied in hPa but is expected in Pa.

Line `4+ndims-` : list of variables where each row specifies the information about the variable.

v	4	2	1	1.0	HGT	H	gpm
v	4	2	2	1.0	UGRD	U	m/s
v	4	2	3	1.0	VGRD	V	m/s
v	4	5	4	1.0	VVEL	OMEGA	Pa/s
v	4	2	5	1.0	TMP	T	K
v	4	6	30	1.0	RH	RH	%
v	4	6	31	1.0	SPFH	SH	kg/kg

In this example, the first character, `v`, signifies that the row describes a variable. The next integer is the dimensionality of the variable (3 or 4). It is assumed that the variable is a function of x, y, and t, but if also a function of pressure, the vertical dimension needs to be specified. The next integer indicated which of the vertical dimensions listed earlier in the file are used for this variable. The fourth item on the line (third integer) is the variable ID according to Table 8.1. Next is the scaling factor to convert the provided values to the units expected for the variable. The next two fields are the WMO variable name and the variable name given in the NetCDF file.

There are several limitations of the template for the custom NWP reader. Firstly, only the NetCDF reader is currently implemented. Secondly, currently all variables are assumed to be transient; so land-use or topography cannot be read with this scheme. Variables that cannot be simply scaled to the expected units cannot be described in this template. For example, WRF files (`iwf=50`) uses a staggered grid with pressure and geopotential given both as a base value plus a perturbation. Lastly, to determine the start time of the data, a reference time must be read. First the time variable is checked for the `GRIB.orgReferenceTime` attribute. If this is not found, then the variable `reftime` is checked, if it exists.

Several examples of these template files used with NAM and GFS forecast products are shown in Appendix 8.3.

## 6 Additional tools/Examples

In the `tools` folder, three programs are included the illustrate the use of the library.

## 6.1 MetCheck

This utility is used for checking the validity of the NWP files. In some cases, the downloading of files might be interrupted or the files on the NCEP server might be incomplete or corrupt. This utility opens each of the files and reads the full grid for the variables: geopotential height,  $V_x$ ,  $V_y$ ,  $V_z$ , and temperature. If any values are outside of expected ranges, then the program stops returning a non-zero stop-code (`stop 1`). Usage:

```
MetCheck iwf idf filename [year]
```

## 6.2 MetSonde

This utility is used for probing the NWP files and returning the temperature as a function of height. Usage:

```
MetSonde lon lat YYYY MM DD HH [WIND_ROOT]
```

The program assumes that the GFS and NCEP files are placed in the default directory, or at least in default subdirectories off of `WIND_ROOT`. Values are interpolated onto the requested point and time and are written to a file (`GFS_prof.dat`) in the current working directory. This 3-column file contains the height (in km), pressure (in Pa) and temperature (in °C).

## 6.3 MetTraj

This utility takes a similar argument list as `MetSonde`, but instead of probing a single point and time, this utility calculates trajectories within either a GFS forecast package or the NCEP reanalysis files. This utility must be compiled with either a `-DFORWARD` or `-DBACKWARD` pre-processor flag to generate separate executables for forward or backward trajectories. The location of the windfiles is assumed to be the default for GFS and NCEP. For forward trajectories, forecast package beginning most closely prior to the requested start time is used unless the start time is greater than two weeks from present. If so, the NCEP files are used. For the backward trajectories, the forecast package chosen is that which can accommodate the default 24-hours of backward integration.

For each of the trajectory calculations, files are written to the current working directory with the longitude and latitude of the trajectory in 1-hour increments at various elevations. The default elevations are 5000, 10000, 20000, 30000, 40000, and 50000 ft. Files for forward trajectories are named `ftraj[1-6].dat` while the backward trajectories are `btraj[1-6].dat`.

Optionally, the length of time to integrate and a user-specified list of output elevations can be given. Elevations are given on the command line must be in km. Usage:

```
MetTraj_F lon lat YYYY MM DD HH [FC_hours nlev lev1 lev2 ...]
```

## 7 Windfile managment scripts

Scripts used to manage GFS and NCEP Reanalysis files are provided in the folder `autorun_scripts`. The script `autorun_gfs0.5deg.00` (and the dependent scripts `get_gfs0.5deg.sh` and `convert_gfs0.5deg.sh`) is used for downloading and converting the 0.5° GFS files from NCEP. This script is intended to be run from a cron job and take as a parameter, the hour of the forecast package. For example `autorun_gfs0.5deg.sh 0` will download for 00Z for the



current date. These scripts use a default root download directory of `/data/WindFiles`, but this can be modified within the scripts. The GFS files will be downloaded to `/data/WindFiles/gfs/gfs.YYYYMMDDHH` where `HH` is the forecast hour (00, 06, 12, or 18). The forecast files from hours 0-99 are downloaded then converted using `netcdf-java 4.5`. The path this package is set to `/ncj/netcdfAll-4.5.jar` but can be modified in `convert_gfs0.5deg.sh`.

Similarly, the script `get_NCEP_50YearReanalysis.sh` manages the downloading of the NCEP reanalysis files. These are placed in `/data/WindFiles/NCEP/YYYY/`

These scripts are intended to be run as cron jobs. The following line can be added to the crontab to download the files once they are available.

```
01 22 * * * /opt/USGS/Ash3d/bin/autorun_scripts/autorun_gfs0.5deg.sh 0 > gfs00_log 2>&1
01 11 * * * /opt/USGS/Ash3d/bin/autorun_scripts/autorun_gfs0.5deg.sh 12 > gfs12_log 2>&1
01 01 * * Sun /opt/USGS/Ash3d/bin/autorun_scripts/autorun_NCEP_50YearReanalysis.sh > NCEP_50yr_log 2>&1
```

Also included in `autorun_scripts` is a simple stand-alone script (`grib2nc.sh`) for converting GRIB2 files to NetCDF using `netcdf-java`.

## 8 Appendix

### 8.1 Variable codes

Table 1: Variable codes

Variable type	ID	Name	dimensions
3d Mechanical	1	Geopotential Height	gmp or $m^2/s^2$
	2	Vx	m/s
	3	Vy	m/s
	4	Vz	Pa/s
	5	Temperature	K
	6	Wind speed	m/s
	7	Wind direction	deg East of North
Surface	10	Planetary Boundary Layer Height	m
	11	U @ 10m	m/s
	12	V @ 10m	m/s
	13	Friction velocity	m/s
	14	Displacement Height	m
	15	Snow cover	%
	16	Soil moisture	kg/m <sup>2</sup>
	17	Surface Roughness	m
	18	Wind gust speed	m/s
19	surface temperature	K	
Atmospheric Structure	20	pressure at lower cloud base	Pa
	21	pressure at lower cloud top	Pa
	22	temperature at lower cloud top	K
	23	Total Cloud cover	%
	24	Cloud cover (low)	%
25	Cloud cover (convective)	%	
Moisture	30	Rel. Hum	%
	31	QV (specific humidity)	kg/kg
	32	QL (liquid)	kg/kg
	33	QI (ice)	kg/kg
Precipitation	40	Categorical rain	0 or 1
	41	Categorical snow	0 or 1
	42	Categorical frozen rain	0 or 1
	43	Categorical ice	0 or 1
	44	Precipitation rate large-scale (liquid)	kg/m <sup>2</sup> s
	45	Precipitation rate convective (liquid)	kg/m <sup>2</sup> s
	46	Precipitation rate large-scale (ice)	kg/m <sup>2</sup> s
47	Precipitation rate convective (ice)	kg/m <sup>2</sup> s	

## 8.2 NWP product ID (iwf)

Several NWP product (both reanalysis and forecast) are recognized and do not need template files. These are listed in the table below by the format code. `iwf=1,2` are ASCII files. `iwf=3-19` are reserved for products with projected grids. `iwf=20-49` are reserved for global (lon/lat) grids. `iwf=50` is for NetCDF output from Weather Research and Forecasting (WRF) simulations.

Table 2: NWP product ID

iwf	Product	NCEP Grid	resolution
0	Custom format based on template		
1	ASCII profile		
2	Radiosonde data		
3	North American Regional Reanalysis NARR-A	221	32 km
4	North American Regional Reanalysis NARR-A+B	221	32 km
5	NAM Regional Alaska	216	45 km
6	NAM N. Hemisphere	104	90 km
7	NAM Regional CONUS	212	40 km
8	NAM Regional CONUS	218	12 km
10	NAM Regional Alaska	242	11.25 km
11	NAM Regional Hawaii	196	2.5 km
12	NAM Regional Alaska	198	5.953 km
13	NAM Regional Alaska	91	2.976 km
20	GFS	4	0.5°
21	GFS	3	1.0°
22	GFS	193	0.25°
23	NCEP-DOE Reanalysis 2	2	2.5°
24	NASA-MERRA-2 Reanalysis	None	0.625 × 0.5°
25	NCEP/NCAR Reanalysis 1	2	2.5°
27	NOAA-CIRES 20th Century Reanalysis	2	2.5°
28	ECMWF ERA-Interim Reanalysis	170	~ 0.7°-Gaussian
32	Air Force Weather Agency subcenter = 0	None	0.25°
33	CCSM3.0 Community Atmosphere Model	None	3.75 × 3.7°
40	NASA GEOS-5 Cp	None	0.625 × 0.5°
41	NASA GEOS-5 Np	None	0.3125 × 0.25°
50	WRF - output	None	

### 8.3 Example Template files

## NCEP-Grids




Figure 1: Outlines of North America Mesoscale (NAM) and Global Forecast System (GFS) models from NCEP that are compatible with MetReader template files.

#### 8.4 NAM 221 Regional N.America (32.5 km)

nam.t00z.awip3200.tm00

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>




Figure 2: NAM 221, Regional N. America 32.5 km Lambert Conformal Conical

## n221\_template.txt

0 4	-107.0	50.0	50.0	50.0	6371.229		Projection params line
3.0							Time interval
9 29							ndims, nvars
d	1		1.0	time			hours since
d	2		1.0	isobaric1			Pa
d	3		1.0	y			km or deg
d	4		1.0	x			km or deg
d	5		1.0	isobaric1			Pa
d	6		1.0	isobaric1			Pa
d	7		1.0	height_above_ground3			m
d	8		1.0	depth_below_surface_layer			m
d	9		1.0	isobaric4			Pa
v 4	2 1		1.0	HGT	Geopotential_height_isobaric		gpm
v 4	2 2		1.0	UGRD	u-component_of_wind_isobaric		m/s
v 4	2 3		1.0	VGRD	v-component_of_wind_isobaric		m/s
v 4	5 4		1.0	VVEL	Vertical_velocity_pressure_isobaric		Pa/s
v 4	2 5		1.0	TMP	Temperature_isobaric		K
v 3	0 10		1.0	HPBL	Planetary_Boundary_Layer_Height_surface		m
v 4	7 11		1.0	UGRD	u-component_of_wind_height_above_ground		m/s
v 4	7 12		1.0	VGRD	v-component_of_wind_height_above_ground		m/s
v 3	0 13		1.0	FRICV	Frictional_Velocity_surface		m.s-1
v 3	0 15		1.0	SNOD	Snow_depth_surface		m
v 4	2 16		1.0	SOILW	Volumetric_Soil_Moisture_Content_depth_below_surface_layer		Fraction
v 3	0 17		1.0	SFCR	Surface_roughness_surface		m
v 3	0 18		1.0	GUST	Wind_speed_gust_surface		m/s
v 3	0 20		1.0	PRES	Pressure_cloud_base		Pa
v 3	0 21		1.0	PRES	Pressure_cloud_tops		Pa
v 3	0 23		1.0	TCDC	Total_cloud_cover_entire_atmosphere		%
v 4	6 30		1.0	RH	Relative_humidity_isobaric		%
v 4	6 31		1.0	SPFH	Specific_humidity_isobaric		kg/kg
v 4	9 32		1.0	CLWMR	Cloud_mixing_ratio_isobaric		kg/kg
v 4	9 33		1.0	SNMR	Snow_mixing_ratio_isobaric		kg/kg
v 3	0 40		1.0	CRAIN	Categorical_Rain_surface		0 or 1
v 3	0 41		1.0	CSNOW	Categorical_Snow_surface		0 or 1
v 3	0 42		1.0	CFRZR	Categorical_Freezing_Rain_surface		0 or 1
v 3	0 43		1.0	CICEP	Categorical_Ice_Pellets_surface		0 or 1
v 3	0 44		1.0	PRATE	Precipitation_rate_surface		kg.m-2.s-1
v 3	0 45		1.0	CPRAT	Convective_Precipitation_Rate_surface		kg.m-2.s-1
v 3	0 45		1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation		kg.m-2
v 3	0 46		1.0	ACPCP	Convective_precipitation_surface_0_Hour_Accumulation		kg.m-2
v 3	0 47		1.0	NCPCP	Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation		kg.m-2

## 8.5 NAM 216 AK (45.0 km); NAM 242 AK (11.25 km)

nam.t00z.awipak00.tm00

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>

<http://motherlode.ucar.edu/native/conduit/data/nccf/com/nam/prod/nam.YYYYMMDD/>




Figure 3: NAM 216, AK 45.0 km Polar Stereographic

## n216\_template.txt

0 1	-135.0	90.0	0.933	6371.229		Projection params line
3.0						Time interval
9 27						ndims, nvars
d	1		1.0	time		hours since
d	2		1.0	isobaric1		Pa
d	3		1.0	y		km or deg
d	4		1.0	x		km or deg
d	5		1.0	isobaric1		Pa
d	6		1.0	isobaric1		Pa
d	7		1.0	height_above_ground3		m
d	8		1.0	depth_below_surface_layer		m
d	9		1.0	isobaric4		Pa
v 4	2 1		1.0	HGT Geopotential_height_isobaric		gpm
v 4	2 2		1.0	UGRD u-component_of_wind_isobaric		m/s
v 4	2 3		1.0	VGRD v-component_of_wind_isobaric		m/s
v 4	5 4		1.0	VVEL Vertical_velocity_pressure_isobaric		Pa/s
v 4	2 5		1.0	TMP Temperature_isobaric		K
v 3	0 10		1.0	HPBL Planetary_Boundary_Layer_Height_surface		m
v 4	7 11		1.0	UGRD u-component_of_wind_height_above_ground		m/s
v 4	7 12		1.0	VGRD v-component_of_wind_height_above_ground		m/s
v 3	0 13		1.0	FRICV Frictional_Velocity_surface		m.s-1
v 3	0 15		1.0	SNOD Snow_depth_surface		m
v 4	2 16		1.0	SOILW Volumetric_Soil_Moisture_Content_depth_below_surface_layer		Fraction
v 3	0 17		1.0	SFCR Surface_roughness_surface		m
v 3	0 18		1.0	GUST Wind_speed_gust_surface		m/s
v 3	0 20		1.0	PRES Pressure_cloud_base		Pa
v 3	0 21		1.0	PRES Pressure_cloud_tops		Pa
v 3	0 23		1.0	TCDC Total_cloud_cover_entire_atmosphere		%
v 4	6 30		1.0	RH Relative_humidity_isobaric		%
v 4	9 32		1.0	CLWMR Cloud_mixing_ratio_isobaric		kg/kg
v 3	0 40		1.0	CRAIN Categorical_Rain_surface		0 or 1
v 3	0 41		1.0	CSNOW Categorical_Snow_surface		0 or 1
v 3	0 42		1.0	CFRZR Categorical_Freezing_Rain_surface		0 or 1
v 3	0 43		1.0	CICEP Categorical_Ice_Pellets_surface		0 or 1
v 3	0 44		1.0	PRATE Precipitation_rate_surface		kg.m-2.s-1
v 3	0 45		1.0	CPRAT Convective_Precipitation_Rate_surface		kg.m-2.s-1
v 3	0 45		1.0	APCP Total_precipitation_surface_0_Hour_Accumulation		kg.m-2
v 3	0 46		1.0	ACPCP Convective_precipitation_surface_0_Hour_Accumulation		kg.m-2
v 3	0 47		1.0	NCPCP Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation		kg.m-2



Grid 242 uses the same projection and domain as Grid 216, but with 11.25 km grid spacing.

nam.t00z.awak3d00.grb2.tm00

Available from:

ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/  
n242\_template.txt

0 1	-135.0	90.0	0.933	6371.229		Projection params line
3.0						Time interval
9 27						ndims, nvars
d	1		1.0	time		hours since
d	2		1.0	isobaric2		Pa
d	3		1.0	y		km or deg
d	4		1.0	x		km or deg
d	5		1.0	isobaric		Pa
d	6		1.0	isobaric2		Pa
d	7		1.0	height_above_ground3		m
d	8		1.0	depth_below_surface_layer		m
d	9		1.0	isobaric6		Pa
v 4	2 1		1.0	HGT	Geopotential_height_isobaric	gpm
v 4	2 2		1.0	UGRD	u-component_of_wind_isobaric	m/s
v 4	2 3		1.0	VGRD	v-component_of_wind_isobaric	m/s
v 4	5 4		1.0	VVEL	Vertical_velocity_pressure_isobaric	Pa/s
v 4	2 5		1.0	TMP	Temperature_isobaric	K
v 3	0 10		1.0	HPBL	Planetary_Boundary_Layer_Height_surface	m
v 4	7 11		1.0	UGRD	u-component_of_wind_height_above_ground	m/s
v 4	7 12		1.0	VGRD	v-component_of_wind_height_above_ground	m/s
v 3	0 13		1.0	FRICV	Frictional_Velocity_surface	m.s-1
v 3	0 15		1.0	SNOD	Snow_depth_surface	m
v 4	2 16		1.0	SOILW	Volumetric_Soil_Moisture_Content_depth_below_surface_layer	Fraction
v 3	0 17		1.0	SFCR	Surface_roughness_surface	m
v 3	0 18		1.0	GUST	Wind_speed_gust_surface	m/s
v 3	0 20		1.0	PRES	Pressure_cloud_base	Pa
v 3	0 21		1.0	PRES	Pressure_cloud_tops	Pa
v 3	0 23		1.0	TCDC	Total_cloud_cover_entire_atmosphere	%
v 4	6 30		1.0	RH	Relative_humidity_isobaric	%
v 4	9 32		1.0	CLWMR	Cloud_mixing_ratio_isobaric	kg/kg
v 3	0 40		1.0	CRAIN	Categorical_Rain_surface	0 or 1
v 3	0 41		1.0	CSNOW	Categorical_Snow_surface	0 or 1
v 3	0 42		1.0	CFRZR	Categorical_Freezing_Rain_surface	0 or 1
v 3	0 43		1.0	CICEP	Categorical_Ice_Pellets_surface	0 or 1
v 3	0 44		1.0	PRATE	Precipitation_rate_surface	kg.m-2.s-1
v 3	0 45		1.0	CPRAT	Convective_Precipitation_Rate_surface	kg.m-2.s-1
v 3	0 45		1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v 3	0 46		1.0	ACPCP	Convective_precipitation_surface_0_Hour_Accumulation	kg.m-2
v 3	0 47		1.0	NPCCP	Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation	kg.m-2

## 8.6 NAM 91 AK (2.95 km); NAM 198 AK (5.9 km)

nam.t00z.alaskanest.hiresf00.tm00

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>



Figure 4: NAM 91, AK 2.95 km Polar Stereographic (previously NAM 198, AK 5.9 km)

n091\_template.txt

0 1	-150.0	90.0	0.933	6371.229		Projection params line
1.0						Time interval
9 29						ndims, nvars
d	1		1.0	time		hours since
d	2		1.0	isobaric2		Pa
d	3		1.0	y		km or deg
d	4		1.0	x		km or deg
d	5		1.0	isobaric2		Pa
d	6		1.0	isobaric2		Pa
d	7		1.0	height_above_ground3		m
d	8		1.0	depth_below_surface_layer		m
d	9		1.0	isobaric3		Pa
v 4	2 1		1.0	HGT	Geopotential_height_isobaric	gpm
v 4	2 2		1.0	UGRD	u-component_of_wind_isobaric	m/s
v 4	2 3		1.0	VGRD	v-component_of_wind_isobaric	m/s
v 4	5 4		1.0	VVEL	Vertical_velocity_pressure_isobaric	Pa/s
v 4	2 5		1.0	TMP	Temperature_isobaric	K
v 3	0 10		1.0	HPBL	Planetary_Boundary_Layer_Height_surface	m
v 4	7 11		1.0	UGRD	u-component_of_wind_height_above_ground	m/s
v 4	7 12		1.0	VGRD	v-component_of_wind_height_above_ground	m/s
v 3	0 13		1.0	FRICV	Frictional_Velocity_surface	m.s-1
v 3	0 15		1.0	SNOD	Snow_depth_surface	m
v 4	2 16		1.0	SOILW	Volumetric_Soil_Moisture_Content_depth_below_surface_layer	Fraction
v 3	0 17		1.0	SFCR	Surface_roughness_surface	m
v 3	0 18		1.0	GUST	Wind_speed_gust_surface	m/s
v 3	0 20		1.0	PRES	Pressure_cloud_base	Pa
v 3	0 21		1.0	PRES	Pressure_cloud_tops	Pa
v 3	0 23		1.0	TCDC	Total_cloud_cover_entire_atmosphere	%
v 4	6 30		1.0	RH	Relative_humidity_isobaric	%
v 4	6 31		1.0	SPFH	Specific_humidity_isobaric	kg/kg
v 4	9 32		1.0	CLWMR	Cloud_mixing_ratio_isobaric	kg/kg
v 4	9 33		1.0	SNMR	Snow_mixing_ratio_isobaric	kg/kg
v 3	0 40		1.0	CRAIN	Categorical_Rain_surface	0 or 1
v 3	0 41		1.0	CSNOW	Categorical_Snow_surface	0 or 1
v 3	0 42		1.0	CFRZR	Categorical_Freezing_Rain_surface	0 or 1
v 3	0 43		1.0	CICEP	Categorical_Ice_Pellets_surface	0 or 1
v 3	0 44		1.0	PRATE	Precipitation_rate_surface	kg.m-2.s-1
v 3	0 45		1.0	CPRAT	Convective_Precipitation_Rate_surface	kg.m-2.s-1
v 3	0 45		1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v 3	0 46		1.0	ACPCP	Convective_precipitation_surface_0_Hour_Accumulation	kg.m-2
v 3	0 47		1.0	NCPCP	Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation	kg.m-2

## 8.7 NAM 243 Eastern N. Pac./HI (0.40°)

nam.t00z.awiphi00.tm00

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>




Figure 5: NAM 243, Eastern N. Pac./HI 0.4°

## n243\_template.txt

1 1	-150.0	90.0	0.933	6371.229		Projection params line	
3.0						Time interval	
9	24					ndims, nvars	
d	1	1.0	time			hours since	
d	2	1.0	isobaric2			Pa	
d	3	1.0	lat			km or deg	
d	4	1.0	lon			km or deg	
d	5	1.0	isobaric1			Pa	
d	6	1.0	isobaric2			Pa	
d	7	1.0	height_above_ground			m	
d	8	1.0	depth_below_surface_layer			m	
d	9	1.0	isobaric4			Pa	
v	4	2	1.0	HGT	Geopotential_height_isobaric	gpm	
v	4	2	2	1.0	UGRD	u-component_of_wind_isobaric	m/s
v	4	2	3	1.0	VGRD	v-component_of_wind_isobaric	m/s
v	4	5	4	1.0	VVEL	Vertical_velocity_pressure_isobaric	Pa/s
v	4	2	5	1.0	TMP	Temperature_isobaric	K
v	4	7	11	1.0	UGRD	u-component_of_wind_height_above_ground	m/s
v	4	7	12	1.0	VGRD	v-component_of_wind_height_above_ground	m/s
v	3	0	13	1.0	FRICV	Frictional_Velocity_surface	m.s-1
v	3	0	17	1.0	SFCR	Surface_roughness_surface	m
v	3	0	20	1.0	PRES	Pressure_cloud_base	Pa
v	3	0	21	1.0	PRES	Pressure_cloud_tops	Pa
v	3	0	23	1.0	TCDC	Total_cloud_cover_entire_atmosphere	%
v	4	6	30	1.0	RH	Relative_humidity_isobaric	%
v	4	6	31	1.0	SPFH	Specific_humidity_isobaric	kg/kg
v	4	9	32	1.0	CLWMR	Cloud_mixing_ratio_isobaric	kg/kg
v	4	9	33	1.0	SNMR	Snow_mixing_ratio_isobaric	kg/kg
v	3	0	40	1.0	CRAIN	Categorical_Rain_surface	0 or 1
v	3	0	41	1.0	CSNOW	Categorical_Snow_surface	0 or 1
v	3	0	42	1.0	CFRZR	Categorical_Freezing_Rain_surface	0 or 1
v	3	0	43	1.0	CICEP	Categorical_Ice_Pellets_surface	0 or 1
v	3	0	44	1.0	PRATE	Precipitation_rate_surface	kg.m-2.s-1
v	3	0	45	1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v	3	0	46	1.0	ACPCP	Convective_precipitation_surface_0_Hour_Accumulation	kg.m-2
v	3	0	47	1.0	NCPCP	Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation	kg.m-2

## 8.8 NAM 182 HI (0.108°)

nam.t00z.afwahi00.grb2.tm00

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>




Figure 6: NAM 182, HI 0.108°

n182\_template.txt

1 1	-150.0	90.0	0.933	6371.229		Projection params line	
3.0						Time interval	
8	22					ndims, nvars	
d	1	1.0	time1			hours since	
d	2	1.0	isobaric			Pa	
d	3	1.0	lat			km or deg	
d	4	1.0	lon			km or deg	
d	5	1.0	isobaric			Pa	
d	6	1.0	isobaric			Pa	
d	7	1.0	height_above_ground2			m	
d	8	1.0	depth_below_surface_layer			m	
v	4	2	1	1.0	HGT	Geopotential_height_isobaric	gpm
v	4	2	2	1.0	UGRD	u-component_of_wind_isobaric	m/s
v	4	2	3	1.0	VGRD	v-component_of_wind_isobaric	m/s
v	4	5	4	1.0	VVEL	Vertical_velocity_pressure_isobaric	Pa/s
v	4	2	5	1.0	TMP	Temperature_isobaric	K
v	3	0	10	1.0	HPBL	Planetary_Boundary_Layer_Height_surface	m
v	4	7	11	1.0	UGRD	u-component_of_wind_height_above_ground	m/s
v	4	7	12	1.0	VGRD	v-component_of_wind_height_above_ground	m/s
v	3	0	13	1.0	FRICV	Frictional_Velocity_surface	m.s-1
v	3	0	15	1.0	SNOD	Snow_depth_surface	m
v	4	2	16	1.0	SOILW	Volumetric_Soil_Moisture_Content_depth_below_surface_layer	Fraction
v	3	0	23	1.0	TCDC	Total_cloud_cover_entire_atmosphere	%
v	4	6	30	1.0	RH	Relative_humidity_isobaric	%
v	4	6	31	1.0	SPFH	Specific_humidity_isobaric	kg/kg
v	4	9	32	1.0	CLWMR	Cloud_mixing_ratio_isobaric	kg/kg
v	4	9	33	1.0	SNMR	Snow_mixing_ratio_isobaric	kg/kg
v	3	0	40	1.0	CRAIN	Categorical_Rain_surface	0 or 1
v	3	0	41	1.0	CSNOW	Categorical_Snow_surface	0 or 1
v	3	0	42	1.0	CFRZR	Categorical_Freezing_Rain_surface	0 or 1
v	3	0	43	1.0	CICEP	Categorical_Ice_Pellets_surface	0 or 1
v	3	0	45	1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v	3	0	46	1.0	ACPCP	Convective_precipitation_surface_0_Hour_Accumulation	kg.m-2

## 8.9 NAM 196 HI (2.5 km)

nam.t00z.hawaiiinst.hiresf00.tm0

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>




Figure 7: NAM 196, AK 2.5 km Mercator



## n196\_template.txt

0 5	198.475	20.0	6371.229				Projection params line
1.0							Time interval
9 27							ndims, nvars
d	1	1.0	time				hours since
d	2	1.0	isobaric2				Pa
d	3	1.0	y				km or deg
d	4	1.0	x				km or deg
d	5	1.0	isobaric2				Pa
d	6	1.0	isobaric2				Pa
d	7	1.0	height_above_ground3				m
d	8	1.0	depth_below_surface_layer				m
d	9	1.0	isobaric4				Pa
v 4	2 1	1.0	HGT	Geopotential_height_isobaric			gpm
v 4	2 2	1.0	UGRD	u-component_of_wind_isobaric			m/s
v 4	2 3	1.0	VGRD	v-component_of_wind_isobaric			m/s
v 4	5 4	1.0	VVEL	Vertical_velocity_pressure_isobaric			Pa/s
v 4	2 5	1.0	TMP	Temperature_isobaric			K
v 3	0 10	1.0	HPBL	Planetary_Boundary_Layer_Height_surface			m
v 4	7 11	1.0	UGRD	u-component_of_wind_height_above_ground			m/s
v 4	7 12	1.0	VGRD	v-component_of_wind_height_above_ground			m/s
v 3	0 13	1.0	FRICV	Frictional_Velocity_surface			m.s-1
v 3	0 15	1.0	SNOD	Snow_depth_surface			m
v 4	2 16	1.0	SOILW	Volumetric_Soil_Moisture_Content_depth_below_surface_layer			Fraction
v 3	0 17	1.0	SFCR	Surface_roughness_surface			m
v 3	0 18	1.0	GUST	Wind_speed_gust_surface			m/s
v 3	0 20	1.0	PRES	Pressure_cloud_base			Pa
v 3	0 21	1.0	PRES	Pressure_cloud_tops			Pa
v 3	0 23	1.0	TCDC	Total_cloud_cover_entire_atmosphere			%
v 4	6 30	1.0	RH	Relative_humidity_isobaric			%
v 4	6 31	1.0	SPFH	Specific_humidity_isobaric			kg/kg
v 4	9 32	1.0	CLWMR	Cloud_mixing_ratio_isobaric			kg/kg
v 4	9 33	1.0	SNMR	Snow_mixing_ratio_isobaric			kg/kg
v 3	0 40	1.0	CRAIN	Categorical_Rain_surface			0 or 1
v 3	0 41	1.0	CSNOW	Categorical_Snow_surface			0 or 1
v 3	0 42	1.0	CFRZR	Categorical_Freezing_Rain_surface			0 or 1
v 3	0 43	1.0	CICEP	Categorical_Ice_Pellets_surface			0 or 1
v 3	0 44	1.0	PRATE	Precipitation_rate_surface			kg.m-2.s-1
v 3	0 45	1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation			kg.m-2
v 3	0 47	1.0	NCPCP	Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation			kg.m-2

## 8.10 NAM 211/212/218/227 CONUS




Figure 8: NAM 211/212/218/227 CONUS Lambert Conformal Conical (dots correspond to the 81.3 km 211 grid)

Available from:

ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/  
NAM 211 CONUS (81.3 km) :: nam.t00z.awp21100.tm00  
n211\_template.txt

```
0 4 265.0 25.0 25.0 25.0 6371.229
6.0
9 18
d 1 1.0 time
d 2 1.0 isobaric
d 3 1.0 y
d 4 1.0 x
d 5 1.0 isobaric
d 6 1.0 isobaric
d 7 1.0 height_above_ground1
d 8 1.0 depth_below_surface_layer
d 9 1.0 isobaric
v 4 2 1 1.0 HGT Geopotential_height_isobaric
v 4 2 2 1.0 UGRD u-component_of_wind_isobaric
v 4 2 3 1.0 VGRD v-component_of_wind_isobaric
v 4 5 4 1.0 VVEL Vertical_velocity_pressure_isobaric
v 4 2 5 1.0 TMP Temperature_isobaric
v 3 0 10 1.0 HPBL Planetary_Boundary_Layer_Height_surface
v 4 7 11 1.0 UGRD u-component_of_wind_height_above_ground
v 4 7 12 1.0 VGRD v-component_of_wind_height_above_ground
v 3 0 18 1.0 GUST Wind_speed_gust_surface
v 3 0 20 1.0 PRES Pressure_cloud_base
v 3 0 21 1.0 PRES Pressure_cloud_tops
v 4 6 30 1.0 RH Relative_humidity_isobaric
v 3 0 40 1.0 CRAIN Categorical_Rain_surface
v 3 0 41 1.0 CSNOW Categorical_Snow_surface
v 3 0 42 1.0 CFRZR Categorical_Freezing_Rain_surface
v 3 0 43 1.0 CICEP Categorical_Ice_Pellets_surface
v 3 0 45 1.0 APCP Total_precipitation_surface_0_Hour_Accumulation
v 3 0 46 1.0 ACPCP Convective_precipitation_surface_0_Hour_Accumulation

Projection params line
Time interval
ndims, nvars
hours since
Pa
km or deg
km or deg
Pa
Pa
m
m
Pa
gpm
m/s
m/s
Pa/s
K
m
m/s
m/s
m/s
Pa
Pa
%
0 or 1
0 or 1
0 or 1
0 or 1
kg.m-2
kg.m-2
```

NAM 212 CONUS (40.6 km) :: nam.t00z.awip3d00.tm00  
n212\_template.txt

```
0 4 265.0 25.0 25.0 25.0 6371.229
3.0
9 29
d 1 1.0 time
d 2 1.0 isobaric1
d 3 1.0 y
d 4 1.0 x
d 5 1.0 isobaric1
d 6 1.0 isobaric1
d 7 1.0 height_above_ground2
d 8 1.0 depth_below_surface_layer
d 9 1.0 isobaric1
v 4 2 1 1.0 HGT Geopotential_height_isobaric
v 4 2 2 1.0 UGRD u-component_of_wind_isobaric
v 4 2 3 1.0 VGRD v-component_of_wind_isobaric
v 4 5 4 1.0 VVEL Vertical_velocity_pressure_isobaric
v 4 2 5 1.0 TMP Temperature_isobaric
v 3 0 10 1.0 HPBL Planetary_Boundary_Layer_Height_surface
v 4 7 11 1.0 UGRD u-component_of_wind_height_above_ground
v 4 7 12 1.0 VGRD v-component_of_wind_height_above_ground
v 3 0 13 1.0 FRICV Frictional_Velocity_surface
v 3 0 15 1.0 SNOD Snow_depth_surface
v 4 2 16 1.0 SOILW Volumetric_Soil_Moisture_Content_depth_below_surface_layer
v 3 0 17 1.0 SFCR Surface_roughness_surface
v 3 0 18 1.0 GUST Wind_speed_gust_surface
v 3 0 20 1.0 PRES Pressure_cloud_base
v 3 0 21 1.0 PRES Pressure_cloud_tops
v 3 0 23 1.0 TCDC Total_cloud_cover_entire_atmosphere
v 4 6 30 1.0 RH Relative_humidity_isobaric
v 4 6 31 1.0 SPFH Specific_humidity_isobaric
v 4 9 32 1.0 CLWMR Cloud_mixing_ratio_isobaric
v 4 9 33 1.0 SNMR Snow_mixing_ratio_isobaric
v 3 0 40 1.0 CRAIN Categorical_Rain_surface
v 3 0 41 1.0 CSNOW Categorical_Snow_surface
v 3 0 42 1.0 CFRZR Categorical_Freezing_Rain_surface
v 3 0 43 1.0 CICEP Categorical_Ice_Pellets_surface
v 3 0 44 1.0 PRATE Precipitation_rate_surface
v 3 0 45 1.0 CPRAT Convective_Precipitation_Rate_surface
v 3 0 45 1.0 APCP Total_precipitation_surface_0_Hour_Accumulation
v 3 0 46 1.0 ACPCP Convective_precipitation_surface_0_Hour_Accumulation
v 3 0 47 1.0 NCPCP Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation

Projection params line
Time interval
ndims, nvars
hours since
Pa
km or deg
km or deg
Pa
Pa
m
m
Pa
gpm
m/s
m/s
Pa/s
K
m
m/s
m/s
m.s-1
m
Fraction
m/s
Pa
Pa
%
%
kg/kg
kg/kg
kg/kg
0 or 1
0 or 1
0 or 1
0 or 1
0 or 1
kg.m-2.s-1
kg.m-2.s-1
kg.m-2
kg.m-2
kg.m-2
```

NAM 218 CONUS (12.2 km) :: nam.t00z.awphys00.grb2.tm00  
n218\_template.txt

0 4 265.0 25.0 25.0 25.0 6371.229		Projection params line
1.0		Time interval
9 21		ndims, nvars
d 1	1.0 time	hours since
d 2	1.0 isobaric1	Pa
d 3	1.0 y	km or deg
d 4	1.0 x	km or deg
d 5	1.0 isobaric1	Pa
d 6	1.0 isobaric	Pa
d 7	1.0 height_above_ground4	m
d 8	1.0 depth_below_surface_layer	m
d 9	1.0 isobaric4	Pa
v 4 2 1	1.0 HGT Geopotential_height_isobaric	gpm
v 4 2 2	1.0 UGRD u-component_of_wind_isobaric	m/s
v 4 2 3	1.0 VGRD v-component_of_wind_isobaric	m/s
v 4 5 4	1.0 VVEL Vertical_velocity_pressure_isobaric	Pa/s
v 4 2 5	1.0 TMP Temperature_isobaric	K
v 3 0 10	1.0 HPBL Planetary_Boundary_Layer_Height_surface	m
v 4 7 11	1.0 UGRD u-component_of_wind_height_above_ground	m/s
v 4 7 12	1.0 VGRD v-component_of_wind_height_above_ground	m/s
v 3 0 13	1.0 FRICV Frictional_Velocity_surface	m.s-1
v 3 0 15	1.0 SNOD Snow_depth_surface	m
v 4 2 16	1.0 SOILW Volumetric_Soil_Moisture_Content_depth_below_surface_layer	Fraction
v 3 0 17	1.0 SFCR Surface_roughness_surface	m
v 3 0 18	1.0 GUST Wind_speed_gust_surface	m/s
v 3 0 23	1.0 TCDC Total_cloud_cover_entire_atmosphere	%
v 4 6 30	1.0 RH Relative_humidity_isobaric	%
v 3 0 40	1.0 CRAIN Categorical_Rain_surface	0 or 1
v 3 0 41	1.0 CSNOW Categorical_Snow_surface	0 or 1
v 3 0 42	1.0 CFRZR Categorical_Freezing_Rain_surface	0 or 1
v 3 0 43	1.0 CICEP Categorical_Ice_Pellets_surface	0 or 1
v 3 0 45	1.0 APCP Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v 3 0 46	1.0 ACPCP Convective_precipitation_surface_0_Hour_Accumulation	kg.m-2

NAM 227 CONUS (5.1 km) :: nam.t00z.conusnest.hiresf00.tm00  
n227\_template.txt

0 4 265.0 25.0 25.0 25.0 6371.229		Projection params line
1.0		Time interval
9 27		ndims, nvars
d 1	1.0 time	hours since
d 2	1.0 isobaric2	Pa
d 3	1.0 y	km or deg
d 4	1.0 x	km or deg
d 5	1.0 isobaric2	Pa
d 6	1.0 isobaric2	Pa
d 7	1.0 height_above_ground3	m
d 8	1.0 depth_below_surface_layer	m
d 9	1.0 isobaric4	Pa
v 4 2 1	1.0 HGT Geopotential_height_isobaric	gpm
v 4 2 2	1.0 UGRD u-component_of_wind_isobaric	m/s
v 4 2 3	1.0 VGRD v-component_of_wind_isobaric	m/s
v 4 5 4	1.0 VVEL Vertical_velocity_pressure_isobaric	Pa/s
v 4 2 5	1.0 TMP Temperature_isobaric	K
v 3 0 10	1.0 HPBL Planetary_Boundary_Layer_Height_surface	m
v 4 7 11	1.0 UGRD u-component_of_wind_height_above_ground	m/s
v 4 7 12	1.0 VGRD v-component_of_wind_height_above_ground	m/s
v 3 0 13	1.0 FRICV Frictional_Velocity_surface	m.s-1
v 3 0 15	1.0 SNOD Snow_depth_surface	m
v 4 2 16	1.0 SOILW Volumetric_Soil_Moisture_Content_depth_below_surface_layer	Fraction
v 3 0 17	1.0 SFCR Surface_roughness_surface	m
v 3 0 18	1.0 GUST Wind_speed_gust_surface	m/s
v 3 0 20	1.0 PRES Pressure_cloud_base	Pa
v 3 0 21	1.0 PRES Pressure_cloud_tops	Pa
v 3 0 23	1.0 TCDC Total_cloud_cover_entire_atmosphere	%
v 4 6 30	1.0 RH Relative_humidity_isobaric	%
v 4 6 31	1.0 SPFH Specific_humidity_isobaric	kg/kg
v 4 9 32	1.0 CLWMR Cloud_mixing_ratio_isobaric	kg/kg
v 4 9 33	1.0 SNMR Snow_mixing_ratio_isobaric	kg/kg
v 3 0 40	1.0 CRAIN Categorical_Rain_surface	0 or 1
v 3 0 41	1.0 CSNOW Categorical_Snow_surface	0 or 1
v 3 0 42	1.0 CFRZR Categorical_Freezing_Rain_surface	0 or 1
v 3 0 43	1.0 CICEP Categorical_Ice_Pellets_surface	0 or 1
v 3 0 44	1.0 PRATE Precipitation_rate_surface	kg.m-2.s-1
v 3 0 45	1.0 APCP Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v 3 0 47	1.0 NCPCP Large-scale_precipitation_non-convective_surface_0_Hour_Accumulation	kg.m-2

## 8.11 NAM 181 Caribbean (0.108°)

nam.t00z.afwaca00.grb2.tm00

Available from:

<ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/nam/prod/nam.YYYYMMDD/>




Figure 9: NAM 181, Caribbean 0.108°

n181\_template.txt

1 1	-150.0	90.0	0.933	6371.229		Projection params line	
3.0						Time interval	
8	22					ndims, nvars	
d	1	1.0	time1			hours since	
d	2	1.0	isobaric			Pa	
d	3	1.0	lat			km or deg	
d	4	1.0	lon			km or deg	
d	5	1.0	isobaric			Pa	
d	6	1.0	isobaric			Pa	
d	7	1.0	height_above_ground2			m	
d	8	1.0	depth_below_surface_layer			m	
v	4	2	1	1.0	HGT	Geopotential_height_isobaric	gpm
v	4	2	2	1.0	UGRD	u-component_of_wind_isobaric	m/s
v	4	2	3	1.0	VGRD	v-component_of_wind_isobaric	m/s
v	4	5	4	1.0	VVEL	Vertical_velocity_pressure_isobaric	Pa/s
v	4	2	5	1.0	TMP	Temperature_isobaric	K
v	3	0	10	1.0	HPBL	Planetary_Boundary_Layer_Height_surface	m
v	4	7	11	1.0	UGRD	u-component_of_wind_height_above_ground	m/s
v	4	7	12	1.0	VGRD	v-component_of_wind_height_above_ground	m/s
v	3	0	13	1.0	FRICV	Frictional_Velocity_surface	m.s-1
v	3	0	15	1.0	SNOD	Snow_depth_surface	m
v	4	2	16	1.0	SOILW	Volumetric_Soil_Moisture_Content_depth_below_surface_layer	Fraction
v	3	0	23	1.0	TCDC	Total_cloud_cover_entire_atmosphere	%
v	4	6	30	1.0	RH	Relative_humidity_isobaric	%
v	4	6	31	1.0	SPFH	Specific_humidity_isobaric	kg/kg
v	4	9	32	1.0	CLWMR	Cloud_mixing_ratio_isobaric	kg/kg
v	4	9	33	1.0	SNMR	Snow_mixing_ratio_isobaric	kg/kg
v	3	0	40	1.0	CRAIN	Categorical_Rain_surface	0 or 1
v	3	0	41	1.0	CSNOW	Categorical_Snow_surface	0 or 1
v	3	0	42	1.0	CFRZR	Categorical_Freezing_Rain_surface	0 or 1
v	3	0	43	1.0	CICEP	Categorical_Ice_Pellets_surface	0 or 1
v	3	0	45	1.0	APCP	Total_precipitation_surface_0_Hour_Accumulation	kg.m-2
v	3	0	46	1.0	ACPCP	Convective_precipitation_surface_0_Hour_Accumulation	kg.m-2

8.12 GFS 3/4/193




Figure 10: GFS 3,4,193, (dots correspond to the 1.0° NAM 3 grid)

Available from:  
 ftp://ftp.ncep.noaa.gov/pub/data/nccf/com/gfs/prod/gfs.YYYYMMDD/  
 GFS NCEP Grid 3 (1.0°) :: gfs.t00z.pgrb2.1p00.f000.nc  
 n003\_template.txt

GFS NCEP grid 4 (0.5°) :: n004\_template.txt

1 1 -150.0 90.0 0.933 6371.229		Projection params line
3.0		Time interval
9 13		ndims, nvars
d 1 1.0 time		hours since
d 2 1.0 isobaric3		Pa
d 3 1.0 lat		km or deg
d 4 1.0 lon		km or deg
d 5 1.0 isobaric2		Pa
d 6 1.0 isobaric3		Pa
d 7 1.0 height_above_ground3		m
d 8 1.0 depth_below_surface_layer		m
d 9 1.0 isobaric4		Pa
v 4 2 1 1.0 HGT Geopotential_height_isobaric		gpm
v 4 2 2 1.0 UGRD u-component_of_wind_isobaric		m/s
v 4 2 3 1.0 VGRD v-component_of_wind_isobaric		m/s
v 4 5 4 1.0 VVEL Vertical_velocity_pressure_isobaric		Pa/s
v 4 2 5 1.0 TMP Temperature_isobaric		K
v 3 0 10 1.0 HPBL Planetary_Boundary_Layer_Height_surface		m
v 4 7 11 1.0 UGRD u-component_of_wind_height_above_ground		m/s
v 4 7 12 1.0 VGRD v-component_of_wind_height_above_ground		m/s
v 3 0 15 1.0 SNOD Snow_depth_surface		m
v 4 2 16 1.0 SOILW Volumetric_Soil_Moisture_Content_depth_below_surface_layer		Fraction
v 3 0 18 1.0 GUST Wind_speed_gust_surface		m/s
v 4 6 30 1.0 RH Relative_humidity_isobaric		%
v 4 5 32 1.0 CLWMR Cloud_mixing_ratio_isobaric		kg/kg

GFS NCEP grid 193 (0.25°) :: n193\_template.txt

1 1 -150.0 90.0 0.933 6371.229		Projection params line
3.0		Time interval
9 13		ndims, nvars
d 1 1.0 time		hours since
d 2 1.0 isobaric3		Pa
d 3 1.0 lat		km or deg
d 4 1.0 lon		km or deg
d 5 1.0 isobaric2		Pa
d 6 1.0 isobaric3		Pa
d 7 1.0 height_above_ground3		m
d 8 1.0 depth_below_surface_layer		m
d 9 1.0 isobaric4		Pa
v 4 2 1 1.0 HGT Geopotential_height_isobaric		gpm
v 4 2 2 1.0 UGRD u-component_of_wind_isobaric		m/s
v 4 2 3 1.0 VGRD v-component_of_wind_isobaric		m/s
v 4 5 4 1.0 VVEL Vertical_velocity_pressure_isobaric		Pa/s
v 4 2 5 1.0 TMP Temperature_isobaric		K
v 3 0 10 1.0 HPBL Planetary_Boundary_Layer_Height_surface		m
v 4 7 11 1.0 UGRD u-component_of_wind_height_above_ground		m/s
v 4 7 12 1.0 VGRD v-component_of_wind_height_above_ground		m/s
v 3 0 15 1.0 SNOD Snow_depth_surface		m
v 4 2 16 1.0 SOILW Volumetric_Soil_Moisture_Content_depth_below_surface_layer		Fraction
v 3 0 18 1.0 GUST Wind_speed_gust_surface		m/s
v 4 6 30 1.0 RH Relative_humidity_isobaric		%
v 4 5 32 1.0 CLWMR Cloud_mixing_ratio_isobaric		kg/kg



### 8.13 NASA GEOS-5




Figure 11: NASA GEOS-5, 0.625/0.5°

Available from:

[ftp://gmao\\_ops@ftp.nccs.nasa.gov/fp/forecast//](ftp://gmao_ops@ftp.nccs.nasa.gov/fp/forecast//)

NASA GEOS-5 Cp (0.625/0.5°) ::

GEOS.fp.fcst.inst3\_3d\_asm\_Cp.YYYYYMDD\_00+YYYYMMDD\_mmmm.V01.nc4

nGCp\_template.txt

```
1 1 -150.0 90.0 0.933 6371.229
3.0
6 7
d 1 1.6667e-2 time
d 2 100.0 lev
d 3 1.0 lat
d 4 1.0 lon
d 5 100.0 lev
d 6 100.0 lev
v 4 2 1 1.0 HGT H
v 4 2 2 1.0 UGRD U
v 4 2 3 1.0 VGRD V
v 4 5 4 1.0 VVEL OMEGA
v 4 2 5 1.0 TMP T
v 4 6 30 1.0 RH RH
v 4 6 31 1.0 SPFH SH
```

Projection params line  
Time interval  
ndims, nvars  
hours since  
Pa  
km or deg  
km or deg  
Pa  
Pa  
gpm  
m/s  
m/s  
Pa/s  
K  
%  
kg/kg

NASA GEOS-5 Np (0.25/0.3125°) ::

GEOS.fp.fcst.inst3\_3d\_asm\_Np.YYYYYMDD\_00+YYYYMMDD\_mmmm.V01.nc4

nGNp\_template.txt

```
1 1 -150.0 90.0 0.933 6371.229
3.0
6 7
d 1 1.6667e-2 time
d 2 100.0 lev
d 3 1.0 lat
d 4 1.0 lon
d 5 100.0 lev
d 6 100.0 lev
v 4 2 1 1.0 HGT H
v 4 2 2 1.0 UGRD U
v 4 2 3 1.0 VGRD V
v 4 5 4 1.0 VVEL OMEGA
v 4 2 5 1.0 TMP T
v 4 6 30 1.0 RH RH
v 4 6 31 1.0 SPFH SH
```

Projection params line  
Time interval  
ndims, nvars  
hours since  
Pa  
km or deg  
km or deg  
Pa  
Pa  
gpm  
m/s  
m/s  
Pa/s  
K  
%  
kg/kg