

TI-30X IIS:

A Guide for Teachers

Developed by
Texas Instruments Incorporated

Activities developed by
Gary Hanson and Aletha Paskett

Illustrated by
Jay Garrison

About the Authors

Gary Hanson and **Aletha Paskett** are math teachers in the Jordan Independent School District in Sandy, Utah. They developed the Activities section and assisted in evaluating the appropriateness of the examples in the How to Use the TI-30X IIS section of this guide.

Important Notice Regarding Book Materials

Texas Instruments makes no warranty, either expressed or implied, including but not limited to any implied warranties of merchantability and fitness for a particular purpose, regarding any programs or book materials and makes such materials available **solely** on an “as-is” basis. In no event shall Texas Instruments be liable to anyone for special, collateral, incidental, or consequential damages in connection with or arising out of the purchase or use of these materials, and the sole and exclusive liability of Texas Instruments, regardless of the form of action, shall not exceed the purchase price of this book. Moreover, Texas Instruments shall not be liable for any claim of any kind whatsoever against the use of these materials by any other party.

Note: Using calculators other than the TI-30X IIS may produce results different from those described in these materials.

Permission To Reprint or Photocopy

Permission is hereby granted to teachers to reprint or photocopy in classroom, workshop, or seminar quantities, the pages or sheets in this book that carry a Texas Instruments copyright notice. These pages are designed to be reproduced by teachers for use in classes, workshops, or seminars, provided each copy made shows the copyright notice. Such copies may not be sold, and further distribution is expressly prohibited. Except as authorized above, prior written permission must be obtained from Texas Instruments Incorporated to reproduce or transmit this work or portions thereof in any other form or by any other electronic or mechanical means, including any information storage or retrieval system, unless expressly permitted by federal copyright law.

Send inquiries to this address:
Texas Instruments Incorporated
7800 Banner Drive, M/S 3918
Dallas, TX 75251
Attention: Manager, Business Services

Note: If you request photocopies of all or portions of this book from others, you must include this page (with the permission statement above) to the supplier of the photocopying services.



www.ti.com/calc
ti-cares@ti.com

Copyright © 1999 Texas Instruments Incorporated.
Except for the specific rights granted herein, all rights are reserved.
Printed in the United States of America.

Automatic Power Down, APD, and EOS are trademarks of Texas Instruments Incorporated.

About the Teacher Guide



How the Teacher Guide is Organized

This guide consists of two sections: Activities and How to Use the TI-30X IIS. The Activities section is a collection of activities for integrating the TI-30X IIS into mathematics instruction. The How To Use the TI-30X IIS section is designed to help you teach students how to use the calculator.

Activities

The activities are designed to be teacher-directed. They are intended to help develop mathematical concepts while incorporating the TI-30X IIS as a teaching tool. Each activity is self-contained and includes the following:

- An overview of the mathematical purpose of the activity.
- The mathematical concepts being developed.
- The materials needed to perform the activity.
- The detailed procedure, including step-by-step TI-30X IIS key presses.
- A student activity sheet.

How to Use the TI-30X IIS

This section contains examples on transparency masters. Chapters are numbered and include the following.

- An introductory page describing the calculator keys presented in the example, the location of those keys on the TI-30X IIS, and any pertinent notes about their functions.
- Transparency masters following the introductory page and providing examples of practical applications of the key(s) being discussed. The key(s) being discussed are circled on the TI-30X IIS keyboard.

Things to Keep in Mind

- While many of the examples on the transparency masters may be used to develop mathematical concepts, they were not designed specifically for that purpose.
- For maximum flexibility, each example and activity is independent of the others. Select the transparency master appropriate for the key you are teaching, or select the activity appropriate for the mathematical concept you are teaching.
- If an example does not seem appropriate for your curriculum or grade level, use it to teach the function of a key (or keys), and then provide relevant examples of your own.
- To ensure that everyone starts at the same point, have students reset the calculator by pressing **ON** and **CLEAR** simultaneously or by pressing **2nd** **[RESET]** and then selecting **Y** (yes).

Conventions Used in the Teacher Guide

- In the text, brackets [] around a key's symbol/name indicate that the key is a second, or alternate, function.

For example: **[SIN⁻¹]**

- On the transparency masters, second functions are shown just as they appear on the keyboard.

For example: **SIN⁻¹**
[SIN]

How to Order Additional Teacher Guides

To place an order or to request information about Texas Instruments (TI) calculators, use our e-mail address: **ti-cares@ti.com** visit our TI calculator home page: **www.ti.com/calc** or, call our toll-free number:

1.800.TI.CARES (1.800.842.2737)

Table of Contents



<i>About the Authors</i>	<i>ii</i>	How to Use the TI-30X IIS <i>(continued)</i>	
<i>About the Teacher Guide</i>	<i>iii</i>	14 Notation	88
<i>About the TI-30X IIS</i>	<i>iv</i>	15 Logarithms and Antilogarithms	91
Activities		16 Angle Settings and Conversions	94
The Better Batter — Using the FIX Key	2	17 Polar and Rectangular Conversions	98
Star Voyage — Using Scientific Notation	6	18 Hyperbolics	100
Trig Functions	10	Appendix A Quick Reference to Keys	A-1
What's My Score — 1-Variable Statistics	14	Appendix B Display Indicators	B-1
Heart Rates — 1-Variable Statistics	17	Appendix C Error Messages	C-1
WNBA Stats — 2-Variable Statistics	23	Appendix D Support and Service Information	D-1
How to Use the TI-30X IIS		Appendix E Warranty Information	E-1
1 TI-30X IIS Basic Operations	29		
2 Clear, Insert, and Delete	33		
3 Basic Math	36		
4 Order of Operations and Parentheses	40		
5 Constant	43		
6 Decimals and Decimal Places	45		
7 Memory	47		
8 Fractions	52		
9 Pi	58		
10 Powers, Roots, and Reciprocals	61		
11 Probability	68		
12 Statistics	75		
13 Trigonometry	81		



Activities

The Better Batter — The FIX Key	2
Star Voyage — Scientific Notation	6
Trig Functions	10
What's My Score? — 1-Variable Statistics	14
Heart Rates — 1-Variable Statistics	17
WNBA Stats — 2-Variable Statistics	23

The Better Batter — The FIX Key

Overview

Students use **2nd** [FIX] on the TI-30X IIS to change numbers to different place values. Students calculate batting averages using the TI-30X IIS and then round their answers to three decimal places.

Math Concepts

- rounding
- place value
- division
- comparing and ordering decimals

Materials

- TI-30X IIS
- pencil
- student activity

Introduction

1. Have students practice rounding the following numbers to 3 decimal places using pencil and paper.

- | | |
|------------|--------|
| a. 2.35647 | 2.356 |
| b. 15.3633 | 15.363 |
| c. 0.02698 | 0.027 |

2. Have students round the following numbers to 4 decimal places using the TI-30X IIS.

- | | |
|--------------|---------|
| a. 4.39865 | 4.3987 |
| b. 72.965912 | 72.9659 |
| c. 0.29516 | 0.2952 |
| d. 0.00395 | 0.0040 |

Activity

Present the following problem to students:

You are going to play Virtual Baseball. You need to select 9 players from the list to be on your team. Choose the players with the best batting averages. Find the batting averages (number of hits ÷ number of times at bat) rounded to 3 decimal places for each player. Make a list of your players in order, from highest to lowest.

See the table on the next page for solutions.

1. Enter the first number.
4.39865
2. Press **2nd** [FIX] to display the menu that lets you set the number of decimal places.
F0123456789
3. Press **4** to select 4 decimal places.
4.39865
4. Press **ENTER**.
4.39865
4.3987

The Better Batter — The FIX Key (Continued)

Player	Number of Hits	Number of Times at Bat	Batting Average
C. Ripken	122	368	0.332
Puckett	119	363	0.328
Molitor	119	364	0.327
Greenwell	104	334	0.311
Tartabull	103	311	0.331
Palmeiro	120	366	0.328
Franco	109	344	0.317
Joyner	105	338	0.311
Boggs	106	329	0.322
Baines	91	290	0.314
Sax	113	388	0.291
Williams	20	74	0.270
Sheridan	15	63	0.238
Barfield	64	284	0.225
Mattingly	109	367	0.297
Hall	87	280	0.311

The Better Batter — Name _____

The FIX Key Date _____




Problems

1. Round the following numbers to 3 decimal places.

- a. 2.35647 _____
- b. 15.3633 _____
- c. 0.02698 _____

2. Using the TI-30X IIS, round the following numbers to 4 decimal places.

- a. 4.39865 _____
- b. 72.965912 _____
- c. 0.29516 _____
- d. 0.00395 _____



The Better Batter — Name _____

The FIX Key Date _____



Problem

You are going to play Virtual Baseball. You need to select 9 players from the list to be on your team. Choose the players with the best batting averages.

Procedure

1. Find the batting averages (number of hits ÷ number of times at bat) rounded to 3 decimal places for each player.

Player	Number of Hits	Number of Times at Bat	Batting Average (rounded to 3 decimal places)
C. Ripken	122	368	
Puckett	119	363	
Molitor	119	364	
Greenwell	104	334	
Tartabull	103	311	
Palmeiro	120	366	
Franco	109	344	
Joyner	105	338	
Boggs	106	329	
Baines	91	290	
Sax	113	388	
Williams	20	74	
Sheridan	15	63	
Barfield	64	284	
Mattingly	109	367	
Hall	87	280	

2. Make a list of your players in order, from highest to lowest.

Player 1 _____	Player 6 _____
Player 2 _____	Player 7 _____
Player 3 _____	Player 8 _____
Player 4 _____	Player 9 _____
Player 5 _____	

Star Voyage — Scientific Notation

Overview

Students investigate scientific notation by changing numbers into scientific notation, and then using them in calculations.

Math Concepts

- scientific notation
- addition
- division

Materials

- TI-30X IIS
- pencil
- student activity

Introduction

Set up the activity by telling your students:

The standard form for scientific notation is $a \times 10^n$, where a is greater than or equal to 1 and less than 10, and n is an integer.

1. Have students practice writing the following numbers in scientific notation using pencil and paper.

- | | |
|---------------------|------------------------|
| a. 93 000 000 | 9.3×10^7 |
| b. 384 000 000 000 | 3.84×10^{11} |
| c. 0.00000000000234 | 2.34×10^{-12} |
| d. 0.0000000157 | 1.57×10^{-8} |

2. Have students change the following numbers into scientific notation using the TI-30X IIS.

- | | |
|----------------|----------------------|
| a. 12 000 000 | 1.2×10^7 |
| b. 974 000 000 | 9.74×10^8 |
| c. 0.0000034 | 3.4×10^{-6} |
| d. 0.000000004 | 4×10^{-9} |

Note: Answers assume the default floating decimal setting.

3. Have students change the following numbers into floating decimal (standard notation).

- | | |
|-------------------------|------------|
| a. 5.8×10^7 | 58 000 000 |
| b. 7.32×10^5 | 732 000 |
| c. 6.2×10^{-6} | 0.0000062 |
| d. 3×10^{-8} | 0.00000003 |

Note: To enter a negative number, press $\boxed{-}$ and then enter the number.

1. Enter the first number.
12000000

2. Press $\boxed{2nd}$ $\boxed{[SCI/ENG]}$.
FLO SCI ENG

3. Press $\boxed{\rightarrow}$ $\boxed{ENTER}$ $\boxed{ENTER}$.
12000000
1.2x10⁰⁷

4. Now, just type the next number and press $\boxed{ENTER}$.

1. Enter **5.8**; press $\boxed{2nd}$ $\boxed{[EE]}$.
5.8E

2. Enter **7**; press $\boxed{2nd}$ $\boxed{[SCI/ENG]}$.
FLO SCI ENG

3. Press $\boxed{\rightarrow}$.
FLO SCI ENG

4. Press $\boxed{ENTER}$ $\boxed{ENTER}$.
5.8E7
58000000.

5. Type the next number and press $\boxed{ENTER}$.

Star Voyage — Scientific Notation (Continued)

Activity

Present the following problem to students:

You are a captain of a starship. You have been assigned to go to Alpha Centauri and you have 5 years to get there. The distance from the sun to Alpha Centauri is 2.5×10^{13} miles. The distance from the earth to the sun is approximately 9.3×10^7 miles. Your ship can travel at the speed of light. You know that light can travel a distance of 6×10^{12} miles in 1 light year. Will you be able to get to Alpha Centauri on time?

Procedure

1. Using the TI-30X IIS, find the total distance you need to travel.

$$2.5 \times 10^{13} + 9.3 \times 10^7 = 2.5000093 \times 10^{13} \text{ miles}$$

2. Next, find out how long it will take you to travel the distance. (distance traveled $\div$ 1 light year)

$$2.5000093 \times 10^{13} \div 6 \times 10^{12} = 4.166682167 \text{ years}$$

3. Can you make the trip in the given time?

Yes

Extension

Now that you have been successful, you have been asked to make another trip. The distance from the Sun to Delta Centauri is 9×10^{13} miles. How long will it take you to get there from Earth?

≈ 15 years

Hint: Make sure your calculator is in scientific notation mode before you beginning addition.

Hint: The Earth is approximately 9.3×10^7 miles from the Sun.

Star Voyage — Scientific Notation

Name _____

Date _____



Problems

1. Write the following numbers in scientific notation.

Standard Notation

Scientific Notation

a. 93 000 000

b. 384 000 000 000

c. 0.000000000000234

d. 0.0000000157

2. Using the TI-30X IIS, change the following numbers into scientific notation.

Standard Notation

Scientific Notation

a. 12 000 000 000 000

b. 974 000 000 000

c. 0.00000000000034

d. 0.0000000004

3. Using the TI-30X IIS, change the following numbers into floating decimal notation (standard).

Scientific Notation

Standard Notation

a. 5.8×10^7

b. 7.32×10^5

c. 6.2×10^{-6}

d. 3×10^{-8}

Star Voyage — Scientific Notation

Name _____

Date _____



Problem

You are a captain of a starship. You have been assigned to go to Alpha Centauri and you have 5 years to get there. The distance from the Sun to Alpha Centauri is 2.5×10^{13} miles. The distance from the Earth to the Sun is approximately 9.3×10^7 miles. Your ship can travel at the speed of light. You know that light can travel a distance of 6×10^{12} miles in 1 light year. Will you be able to get to Alpha Centauri on time?

Procedure

1. Using the TI-30X IIS, find the total distance that you need to travel. _____

Hint: Make sure your calculator is in scientific notation mode before you begin addition.


2. Next, find out how long it will take you to travel the distance. (distance traveled $\div$ 1 light year) _____

3. Can you make the trip in the given time? _____

Extension

Now that you have been successful, you have been asked to make another trip. The distance from the Sun to Delta Centauri is 9×10^{13} miles. How long will it take you to get there from Earth?

Hint: The Earth is approximately 9.3×10^7 miles from the Sun.



Trig Functions

Overview

Students practice solving sine, cosine, and tangent ratios, and solve problems involving trigonometric ratios.

Math Concepts

- multiplication
- division
- trigonometric ratios

Materials

- TI-30X IIS
- pencil
- student activity


Introduction

Introduce the trigonometric ratios to students.

$\sin = \text{opposite leg} \div \text{hypotenuse}$

$\cos = \text{adjacent leg} \div \text{hypotenuse}$

$\tan = \text{opposite leg} \div \text{adjacent leg}$



- Have students find the trigonometric ratios for the triangle using the above definitions. Round to the nearest hundredth if necessary. (Use $\boxed{2\text{nd}} \boxed{[FIX]}$ for rounding.)

- | | |
|-------------|-------------------|
| a. $\sin C$ | $3 \div 5 = 0.60$ |
| b. $\cos C$ | $4 \div 5 = 0.80$ |
| c. $\tan C$ | $3 \div 4 = 0.75$ |
| d. $\sin A$ | $4 \div 5 = 0.80$ |
| e. $\cos A$ | $3 \div 5 = 0.60$ |
| f. $\tan A$ | $4 \div 3 = 1.33$ |

- Have students find the value of each ratio using the TI-30X IIS. Round to the nearest 10 thousandth.

- | | |
|--------------------|----------|
| a. $\sin 71^\circ$ | 0.9455 |
| b. $\tan 31^\circ$ | 0.6009 |
| c. $\cos 25^\circ$ | 0.9063 |

- Have students find the measure of each angle using the TI-30X IIS. Round to the nearest degree.

- | | |
|----------------------|----------------------|
| a. $\sin B = 0.4567$ | 27 degrees |
| b. $\cos A = 0.6758$ | 47 degrees |
| c. $\tan C = 5.83$ | 80 degrees |

- To set 2 decimal places:

- Press $\boxed{2\text{nd}} \boxed{[FIX]}$.
F0123456789
- Press **2** to select 2 decimal places.

- To find $\sin 71^\circ$:

- Press $\boxed{[SIN]}$.
sin(
- Enter **71**; press $\boxed{)} \boxed{[ENTER]}$.
sin(71)
0.945518576
- Press $\boxed{2\text{nd}} \boxed{[FIX]}$ **4**.
sin(71)
0.9455

- To find B when $\sin B = 0.4567$:

- Press $\boxed{2\text{nd}} \boxed{[SIN^{-1}]}$.
sin⁻¹(
- Enter **.4567**; press $\boxed{)} \boxed{[ENTER]}$.
sin⁻¹(.4567)
27.1744
- Press $\boxed{2\text{nd}} \boxed{[FIX]}$ **0**.
sin⁻¹(.4567)
27.

Trig Functions (Continued)


Activity

Present the following problem to students:

You need to build a ramp to your front door. The distance from the ground to the bottom of the door is 1.5 feet. You don't want the angle of incline to be more than 6 degrees. The distance from the street to the door is 20 feet. Is there enough room to build the ramp?

Procedure

1. Make a drawing of the problem.



2. Use the trigonometric ratio

$$\tan = \text{opposite leg} \div \text{adjacent leg}$$

to find angle A.

Angle A is 4.3 degrees (rounded to the nearest tenth). Yes, there is enough room to build the ramp.

Extension

Present the following problem to students:

You want to start the ramp 15 feet away from the door. Can you do that and still have the angle of incline be less than 6 degrees?

Yes, angle A is 5.7°.

1. Press $\boxed{2\text{nd}} \boxed{[\text{TAN}^{-1}]}$.
 $\tan^{-1}()$

2. Enter $1.5 \div 20$ and press $\boxed{)} \boxed{[\text{ENTER}]}$.
 $\tan^{-1}(1.5/20)$
4.3

1. Press $\boxed{2\text{nd}} \boxed{[\text{TAN}^{-1}]}$.
 $\tan^{-1}()$

2. Enter $1.5 \div 15$ and press $\boxed{)} \boxed{[\text{ENTER}]}$.
 $\tan^{-1}(1.5/15)$
5.7

Trig Functions

Name _____


Date _____



Problems

1. Find the trigonometric ratios for the triangle. Round to the nearest hundredth. (Use $\boxed{2nd}$ $\boxed{FIX}$ for rounding.)

- a. $\sin C$ _____
- b. $\cos C$ _____
- c. $\tan C$ _____
- d. $\sin A$ _____
- e. $\cos A$ _____
- f. $\tan A$ _____



2. Using the TI-30X IIS, find the value of each ratio. Round to the nearest ten thousandth.

- a. $\sin 71^\circ$ _____
- b. $\tan 31^\circ$ _____
- c. $\cos 25^\circ$ _____

3. Using the TI-30X IIS, find the measure of each angle. Round to the nearest degree.

- a. $\sin B = 0.4567$ _____
- b. $\cos A = 0.6758$ _____
- c. $\tan C = 5.83$ _____

Trig Functions

Name _____

Date _____



Problem

You need to build a ramp to your front door. The distance from the ground to the bottom of the door is 1.5 feet. You don't want the angle of incline to be more than 6 degrees. The distance from the street to the door is 20 feet. Is there enough room to build the ramp?

Procedure

1. Make a drawing of the problem.

2. Use the trig ratio $\tan = \text{opposite leg} \div \text{adjacent leg}$ to find angle A . (Round your answer to the nearest tenth.) _____

3. Is there room to build the ramp? _____

Extension

You want to start the ramp 15 feet away from the door. Can you do that and still have the angle of incline be less than 6 degrees?

What's My Score? — 1-Variable Statistics

Overview

Students use the given test scores to find averages.

Math Concepts

- averages

Materials

- TI-30X IIS
- pencil
- student activity

Introduction

Discuss finding averages with your students.

Activity

Present the following problem to students:

You and your friend are having a contest. The one gets the highest average on their math tests for one quarter wins. Your scores are 98, 89, 78, 98, and 100. Your friend's scores are 89, 89, 97, 90, and 100. Who is the winner?

Procedure

1. Have students find the average of their scores using the TI-30X IIS. Remember to enter 2 as the frequency for 98 and 1 for all others.
 1. Press $\boxed{2\text{nd}} \boxed{[\text{STAT}]} \boxed{[\text{ENTER}]}$ to select **1-VAR** mode.
 2. Press $\boxed{[\text{DATA}]}$ and enter your first score.
X1 = 98
 3. Press $\odot$ and enter 2 as the frequency for 98.
FRQ = 2
 4. Press $\odot$. Continue entering your scores and frequencies, pressing $\odot$ after each score and frequency.
 5. When finished, press $\boxed{[\text{STATVAR}]} \boxed{\triangleright}$ to select $\bar{x}$, the average. Write it down.
n $\bar{x}$ Sx $\sigma x \rightarrow$
92.6

What's My Score? — 1-Variable Statistics (Cont.)

2. Now find the average of your friend's scores.
Remember to put 2 as the frequency for 89 and 1 for all others.
3. Who won?

Your friend: 93 (You had 92.6.)

Extension

Present the following problem to students:

Your friend took a test on the day you were absent and scored 95. What score do you need to get so that you are the winner?

The score you need: 98

Note: Make sure you exit the **STAT** mode before going on to another problem.

1. Press $\boxed{2\text{nd}} \boxed{[\text{STAT}]} \downarrow \downarrow$ to select **CLRDATA**. Press $\boxed{ENTER}$.
2. Press $\boxed{DATA}$ and enter the friend's first score.
X1 = 89
3. Continue entering the friend's scores and frequencies, following steps 3 and 4 on the previous page.
4. When finished, press $\boxed{STATVAR} \downarrow$ to select $\bar{x}$, the average. Write it down.
n $\bar{x}$ Sx $\sigma_x \rightarrow$
93.0

1. Press $\boxed{2\text{nd}} \boxed{[\text{STAT}]} \downarrow \downarrow$ to **CLRDATA**. Press $\boxed{ENTER}$.
2. Recalculate your friend's average, making sure to include the new score.
3. Use guess and check to figure out what score you need to get.
4. To exit **STAT** mode, press $\boxed{2\text{nd}} \boxed{[\text{EXIT STAT}]} \boxed{ENTER}$.

What's My Score? — Name _____

1-Variable Statistics Date _____



Problems

1. You and your friend are having a contest. Whoever gets the highest average on their math tests for one quarter wins. Your scores are 98, 89, 78, 98, and 100. Your friend's scores are 89, 89, 97, 90, and 100. Who is the winner?


Your average _____

Your friend's average _____

2. Your friend took a test on the day you were absent and scored 95. What score do you need to get so that you are the winner?

Your friend's new average _____

Your new score _____



Heart Rates — 1-Variable Statistics

Overview

Students use the statistics functions of the TI-30XIIS calculator to investigate the effect of exercise on heart rate.

Math Concepts

- mean, minimum, maximum, and range

Materials

- TI-30XIIS
- stopwatch or a watch with a second hand
- student activity

Introduction

Students may be placed in smaller groups for this activity to minimize the amount of data to be entered. Ask students:

- *What do you think the average heart rate is for someone your age?*
- *What about after exercising?*

Activity

Have students complete the following investigation to check their estimations.

1. Have students check their resting heart rates by timing their pulse for 1 minute. (You could have them time for 10 seconds and then multiply by 6, but this could be the quietest minute of your day!)
2. Collect data on the chart. Enter each student's heart rate and a mark in the frequency column. As other students have the same heart rate, add another tally mark in the frequency column.
3. Enter the heart rate data into the TI-30XIIS.
 - a. Enter the first heart rate on the chart as the first **X** value, and the number of tallies for that heart rate as the frequency.
 - b. You must press $\odot$ between entries. For example, enter the first heart rate, and then press $\odot$. Enter the first frequency, and then press $\odot$.

For example, assume a class of 22 students:

Rate	Students	Rate	Students
60	3	63	3
61	5	64	1
62	6	65	4

1. Press $\boxed{2nd}$ $\boxed{[STAT]}$ $\boxed{ENTER}$.
2. Press $\boxed{DATA}$ to enter the heart rates and frequencies.
X1=
3. Enter first heart rate and press $\odot$.
FRQ=
4. Continue entering until you have entered all the heart rates and frequencies.

Heart Rates — 1-Variable Statistics (Continued)

4. Check the statistics calculations. After students display Σx (Sigma x), explain that Σx is the sum of all the heart rates. Ask students:

- *How many heartbeats were there in one minute?*
- *Is the average heart rate higher or lower than you expected?*

5. Now we will see the effect of some exercise on heart rate. Tell students:

If at any point during this portion of the activity you experience pain, weakness, or shortness of breath, stop immediately.

6. Have the students run in place for 2 minutes and then give them these instructions:

- Time your pulse for 1 minute.*
- Record your heart rate as before.*
- Enter the data into the calculator.*
- Compare the average heart rate after running with the resting heart rate.*

7. Now have the students do jumping jacks for 2 minutes. Instruct them to time their pulse for 1 minute again and record as before. Have them enter the data into the calculator again and calculate the average heart rate after jumping jacks. Compare to the other 2 averages.

8. How fit is the class? If the class (or individual) heart rate after jumping jacks is less than 90, then you are in great shape. If it is higher than 125, then you are in poor shape.

9. Instruct students to make a histogram of the 3 sets of data they collected. Ask students:

- *How are the histograms the same?*
- *How are they different?*
- *Is the data grouped the same or is it more spread out in one graph compared to another?*

1. Press $\boxed{\text{STATVAR}}$.

n $\bar{x}$ Sx σx
22.

n should equal the total number of student sampled.

2. Press $\blacktriangleright$ to $\bar{x}$ to see the average heart rate.

n $\bar{x}$ Sx σx
62.

3. Press $\blacktriangleright \blacktriangleright \blacktriangleright$ to Σx .

Σx Σx^2
1370.

Note: The numbers show the results of the example described above. Your students' results will vary depending on the size of group and the heart rate readings.

Heart Rates —

1-Variable Statistics

Name _____

Date _____



Problem

What do you think the average heart rate is for someone your age? What about after exercising?

Procedure

1. Use this table to record your class or group data (resting).

Heartbeats per minute (resting)	Frequency

2. What is the class (group) average? _____
3. What is the total number of heartbeats for the minute? _____

Heart Rates — 1-Variable Statistics

Name _____

Date _____



4. Use this table to record your class or group data (running).

Heartbeats per minute (running)	Frequency

5. What is the class (group) average? _____

6. What is the total number of heartbeats for the minute? _____



Heart Rates —

1-Variable Statistics

Name _____

Date _____



7. Use this table to record your class or group data (jumping).

Heartbeats per minute (jumping)	Frequency

8. What is the class (group) average? _____

9. What is the total number of heartbeats for the minute? _____

10. How fit is the class? _____

Note: If the class (or individual) heart rate after jumping jacks is less than 90, then you are in great shape. If it is higher than 125, then you are in poor shape.

Heart Rates —

1-Variable Statistics

Name _____

Date _____



11. Now make a histogram for each of the 3 sets of data you collected.

Resting

Running

Jumping

12. How are the histograms the same? How are they different? _____

13. Is the data grouped the same or is it more spread out in one graph compared to another? _____

WNBA Stats — 2-Variable Statistics

Overview

Students use WNBA statistics to explore the relationship between 2 variables. They use the TI-30X IIS to compute the regression equation and evaluate some values.

Math Concepts

- 2-variable statistics

Materials

- TI-30X IIS
- pencils
- student activity

Activity

Present the following problem to students:

Do you think WNBA (Women's National Basketball Association) playing time (in minutes per game) is related to how many points a player scores? Do you think it is related to how many rebounds a player gets? Or is it related to the player's field-goal percentage?

Procedure

1. Put the calculator in **STAT** mode and choose **2-VAR** statistics.

2. Using the table in the activity, enter the data. Enter points per game as the **X**-variable and minutes per game (playing time) as the **Y**-variable.

1. Press **[2nd]** **[STAT]** and then **[↓]**.

1-VAR 2-VAR

2. Press **[ENTER]** to select **2-VAR**.

1. Press **[DATA]**.
X1=

2. Enter **10.1** (points per game for the first player, Rhonda Mapp).
X1=10.1

3. Press **[→]**.
Y1=

4. Enter **21.7** (minutes per game for Rhonda Mapp).
Y1=21.7

5. Press **[→]** to enter data for the second player.

6. Enter data for each player in the table. Press **[→]** after entering each data point.

WNBA Stats — 2-Variable Statistics (Continued)

3. Calculate the statistical data.

You may want to fix the decimal to 2 places before doing the statistical calculations.

Ask students:

- *What is the average points scored for the players shown?*
- *What is the average playing time?*
- *What is the total number of points scored per game for all the given players?*

You may want to discuss the other statistical variables and what they mean.

4. The form of the equation is $y = ax + b$. Write the equation for the line of best fit (round to the nearest hundredth).

$$1.56x + 7.02$$

5. The closer the correlation coefficient value is to 1 (or -1), the better the correlation between the two variables. Write the correlation coefficient.

$$r = .91$$

6. Now calculate how many minutes you would expect a player to play if she averages 15 points per game.

1. Press $\boxed{2nd} \boxed{[FIX]}$.
F0123456789

2. Press 2.

1. Press $\boxed{STATVAR}$.
 $\bar{n} \bar{x} Sx \sigma x \bar{y}$
12.00

2. Press $\boxed{\downarrow}$ to $\bar{x}$.
 $\bar{n} \bar{x} Sx \sigma x \bar{y}$
9.33

3. Press $\boxed{\downarrow} \boxed{\downarrow} \boxed{\downarrow}$ to $\bar{y}$.
 $\bar{n} \bar{x} Sx \sigma x \bar{y}$
21.59

4. Press $\boxed{\downarrow} \boxed{\downarrow} \boxed{\downarrow}$ to Σx .
 $Sy \sigma y \Sigma x$
112.00

1. Press $\boxed{\downarrow}$ until you get to **a**. This is the slope of the line of best fit.
 $\Sigma XY \ a \ b \ r$
1.56

2. Press $\boxed{\downarrow}$ to **b**. This is the y-intercept of the line.
 $\Sigma XY \ a \ b \ r$
7.02

3. Press $\boxed{\downarrow}$ to **r**. This is the correlation coefficient.
 $\Sigma XY \ a \ b \ r$
0.91

1. Press $\boxed{\downarrow} \boxed{\downarrow}$ to y' .
 $x' \ y'$

2. Press $\boxed{ENTER}$.

3. Type **15** $\boxed{)}$ and press $\boxed{ENTER}$.
 $y'(15)$
30.44

WNBA Stats — 2-Variable Statistics (Continued)

7. Now calculate how many points you would expect a player to score if she plays 35 minutes a game.
8. Discuss the correlation as a class. Ask students:
 - *Are there other factors affecting the players minutes per game besides points scored?*
 - *What about defense, rebounding, etc.?*

Extension

Now have students use the calculator to investigate the correlation of the other data in the chart such as the relation of field-goal percentage to minutes per game, or rebounds per game to minutes per game. (Remember, since you have already entered the minutes in Y , you only need to enter the new data in X .)

Ask students:

Which 2 variables have the closest correlations? (That is, which have the correlation coefficient closest to 1 or -1?)

1. Press $\boxed{\text{STATVAR}}$.
 $n \quad \bar{x} \quad Sx \quad \sigma x \quad \bar{y}$
12.00
2. Press $\boxed{\text{2ND}} \boxed{\text{2ND}}$ to x' .
 $x' \quad y'$
3. Press $\boxed{\text{ENTER}}$.
4. Type 35 $\boxed{\text{ENTER}}$ and press $\boxed{\text{ENTER}}$.
 $x'(35)$
17.92

WNBA Stats —

2-Variable Statistics

Name _____

Date _____



Problem

Do you think WNBA playing time (in minutes per game) is related to how many points a player scores? Do you think it is related to how many rebounds a player gets? Or is it related to the player's field goal percentage?

Procedure

Use the following table of data to explore the relationships of different pairs of data. Begin by entering the points per game as the x -variable and the minutes per game as the y -variable.

Player	Field-Goal Percentage	Points per Game	Rebounds per Game	Minutes per Game
1. Rhonda Mapp	.506	10.1	4.3	21.7
2. Vicky Bullet	.441	13.3	6.5	31.6
3. Janeth Arcain	.426	6.8	3.6	21.9
4. Cynthia Cooper	.446	22.7	3.7	35
5. Elena Baranova	.420	12.9	9.3	33.6
6. Malgozata Dydek	.482	12.9	7.6	28
7. Heidi Burge	.509	6.7	3.3	16.7
8. Keri Chaconas	.297	4.8	.8	13.2
9. Rebecca Lobo	.484	11.7	6.9	29.2
10. Coquese Washington	.294	1.9	.9	8.1
11. Toni Foster	.467	4.9	1.9	13.6
12. Maria Stepanova	.426	3.3	1.9	6.5

WNBA Stats —

2-Variable Statistics

Name _____

Date _____



Extension

Use the calculator to investigate the correlation of the other data in the table such as the relation of field-goal percentage to minutes per game, or rebounds per game to minutes per game. (Remember, since you have already entered the minutes per game in **Y**, you only need to enter the new data in **X**.)

1. What is the average field-goal percentage?

2. Write the equation for the line of best fit.

3. Write the correlation coefficient.

4. What is the average number of rebounds per game?


5. Write the equation for the line of best fit.


6. What is the total number of rebounds per game for all the given players?

7. Write the equation for the line of best fit.

8. Write the correlation coefficient.

9. Which 2 variables have the closest correlation?
(That is, which have the correlation coefficient closest to 1 or -1?)





How to Use the TI-30X IIS

TI-30X IIS Basic Operations	29
Clear, Insert, and Delete	33
Basic Math	36
Order of Operations and Parentheses	40
Constant	43
Decimals and Decimal Places	45
Memory	47
Fractions	52
Pi	58
Powers, Roots, and Reciprocals	61
Probability	68
Statistics	75
Trigonometry	81
Notation	88
Logarithms and Antilogarithms	91
Angle Settings and Conversions	94
Polar and Rectangular Conversions	98
Hyperbolics	100

TI-30X IIS Basic Operations

1

Keys

1. **[ON]** turns on the calculator.
2. **[2nd]** turns on the **2nd** indicator and accesses the function shown above the next key you press.
3. **[2nd] [OFF]** turns off the calculator and clears the display.
4. **[ENTER]** completes the operation or executes the command.
5. **[2nd] [ANS]** recalls the most recently calculated result and displays it as **Ans**.
6. **[←]** and **[→]** move the cursor left and right to scroll the entry line. Press **[2nd] [←]** or **[2nd] [→]** to scroll to the beginning or end of the entry line.
[↑] and **[↓]** move the cursor up and down through previous entries. **[2nd] [↑]** or **[2nd] [↓]** scroll to the beginning or end of history.

7. **[2nd] [RESET]** displays the **RESET** menu.


RESET: N Y

- Press **[ENTER]** when **N** (no) is underlined to return to the previous screen without resetting the calculator.
- Press **[ENTER]** when **Y** (yes) is underlined to reset the calculator. The message **MEM CLEARED** is displayed.


Note: Pressing **[ON]** and **[CLEAR]** **simultaneously** resets the calculator immediately. No menu or message is displayed.

Notes

- The examples on the transparency masters assume all default settings.
- Resetting the calculator:
 - Returns settings to their defaults: floating decimal (standard) notation and degree (**DEG**) mode.
 - Clears memory variables, pending operations, entries in history, statistical data, constants, and **Ans** (Last Answer).
- The entry line can contain up to 88 characters. When **←** or **→** appear in the display, the entry line contains more characters to the left or right. When **↑** or **↓** appear, more characters are located above or below the entry line.
- Press **[ON]** after Automatic Power Down™ (APD™). The display, pending operations, settings, and memory are retained.



Second, Off, Arrows, Equals



Enter $46 - 23$. Change 46 to 41.
Change 23 to 26 and complete the
operation. Enter $81 + 57$ and
complete the operation. Scroll to
see your previous entries.

Press	Display
$46 \square 23$	$46-23$ DEG
$\leftarrow \leftarrow \leftarrow \leftarrow 1$ $\rightarrow \rightarrow 6 \text{ ENTER}$	$41-26$ 15. DEG
$81 \text{ + } 57 \text{ ENTER}$	$81+57$ 138. DEG
2nd OFF ON ON	$\blacksquare$ DEG
$\uparrow \uparrow \downarrow$	$81+57$ DEG



Reset

2nd ^{RESET}
0

Reset the calculator.

Press	Display
^{RESET} 2nd 0	RESET: <u>N</u> Y DEG
	RESET: N <u>Y</u> DEG
ENTER	MEM CLEARED DEG
CLEAR	 DEG

Pressing **ON** and **CLEAR** at the same time also resets the calculator immediately. No menu or message is displayed.

Using ^{RESET}
2nd **0** or **ON** and **CLEAR** returns all settings to their defaults and clears the memory.



Last Answer (Ans)

2nd ^{ANS}
(-)

Use Last Answer (**Ans**) to calculate $(2+2)^2$.

Press

Display

2 **+** 2 **=**

2+2

4.
DEG

^{ANS}
2nd **(-)** **x²**
=

Ans²

16.
DEG



Clear, Insert, and Delete


2

Keys

1. **CLEAR** clears characters and error messages. Once the display is clear, it moves the cursor to the most recent entry.
2. **2nd** **INS** lets you insert a character at the cursor.
3. **DEL** deletes the character at the cursor.
Hold **DEL** down to delete all characters to the right. Then, each time you press **DEL**, it deletes 1 character to the left of the cursor.

Notes

- The examples on the transparency masters assume all default settings.
- Pressing **CLEAR** does not affect the memories, statistical registers, angle units, or numeric notation.



Delete, Insert

DEL **2nd** **INS** **DEL**

Enter $4569 + 285$, and then change it to $459 + 2865$. Complete the problem.

Press	Display
4569 + 285	4569+285 DEG
← ← ← ← ← ← DEL	459+285 DEG
→ → → → 2nd INS DEL 6	459+2865 DEG
ENTER	459+2865 3324. DEG



Clear

CLEAR

Enter 21595.

Clear the 95.

Clear the entry.

Press

Display

21595

21595

DEG

◀ ▶ **CLEAR**

(Clear to right)

215

DEG

CLEAR

(Clear entry)

█

DEG




Keys


1. $+$ adds.
2. $-$ subtracts.
3. $\times$ multiplies.
4. $\div$ divides.
5. ENTER completes the operation or executes the command.
6. $(-)$ lets you enter a negative number.
7. 2^{nd} $\%$ changes a real number to a percent.

Notes

- The examples on the transparency masters assume all default settings.
- The TI-30X IIS allows implied multiplication.
Example: $3 (4+3) = 21$
- Do not confuse $(-)$ with $-$. $-$ allows subtraction.
- Results of percent calculations display according to the decimal notation mode setting.




Add, Subtract, Multiply, Divide, Equals



Find: $2 + 54 - 6 =$
 $16 \times 21 =$
 $78 \div 2 =$
 $12 \times (5 + 6) =$

Press	Display
2 $\boxed{+}$ 54 $\boxed{-}$ 6 $\boxed{\text{ENTER}}$	2+54-6 50. DEG
16 $\boxed{\times}$ 21 $\boxed{\text{ENTER}}$	16*21 336. DEG
78 $\boxed{\div}$ 2 $\boxed{\text{ENTER}}$	78/2 39. DEG
12 $\boxed{\times}$ $\boxed{(}$ 5 $\boxed{+}$ 6 $\boxed{)}$ $\boxed{\text{ENTER}}$	12*(5+6) 132. DEG



Negative Numbers

(-)

The temperature in Utah was -3°C at 6:00 a.m. By 10:00 a.m. the temperature had risen 12°C . What was the temperature at 10:00 a.m.?

Press

Display

(-) 3 + 12

ENTER

-3+12
9.
DEG



Percent

2nd (%)

Mike makes \$80 per week. He saves 15% of his earnings. How much does Mike save per week?

Press

Display

15

15

DEG

2nd (%) × 80
ENTER

15%*80

12.
DEG



Order of Operations and Parentheses

4


Keys

1. $\left[\right]$ opens a parenthetical expression.
2. $\left) \right]$ closes a parenthetical expression.

Notes

- The examples on the transparency masters assume all default settings.
- The transparency master showing the Equation Operating System (EOS™) demonstrates the order in which the TI-30X IIS completes calculations.
- Operations inside parentheses are performed first. Use $\left[\right]$ $\left) \right]$ to change the order of operations and, therefore, change the result.

Example: $1 + 2 \times 3 = 7$
 $(1 + 2) \times 3 = 9$



Equation Operating System		EOS
1 (first)	Expressions inside $\boxed{() \quad ()}$.	
2	Functions that need a $\boxed{)}$ and precede the expression, such as the $\boxed{\text{SIN}}$, $\boxed{\text{LOG}}$, or $\boxed{2^{\text{nd}}}\boxed{\text{R}\leftrightarrow\text{P}}$ menu items.	
3	Functions entered after the expression, such as $\boxed{x^2}$ and angle unit modifiers ($^\circ$, $'$, $''$, r , g).	
4	Fractions.	
5	Exponentiation ($\boxed{\wedge}$) and roots ($\boxed{2^{\text{nd}}}\boxed{\wedge}^{\sqrt{x}}$).	
6	Negation ($\boxed{(-)}$).	
7	Permutations (nPr), and combinations (nCr).	
8	Multiplication, implied multiplication, and division.	
9	Addition and subtraction.	
10	Conversions ($\boxed{2^{\text{nd}}}\boxed{\text{A}\frac{\text{b}}{\text{c}}\leftrightarrow\text{d}/\text{e}}$, $\boxed{2^{\text{nd}}}\boxed{\text{PRB}}$, and $\blacktriangleright\text{DMS}$).	
11 (last)	$\boxed{\text{ENTER}}$ completes all operations and closes all open parentheses.	

Order of Operations



$$1 + 2 \times 3 =$$

Press

Display

1 $\boxed{+}$ 2 $\boxed{\times}$ 3
 $\boxed{\text{ENTER}}$

1+2*3[↑]
 7.
 DEG

$$(1 + 2) \times 3 =$$

Press

Display

$\boxed{(}$ 1 $\boxed{+}$ 2 $\boxed{)}$ $\boxed{\times}$ 3 $\boxed{\text{ENTER}}$

(1+2)*3[↑]
 9.
 DEG




Keys

1. **[2nd] [K]** turns on the constant mode and lets you define a constant. A **K** displays when the constant mode is on.
2. **[ENTER]** places the contents of **K** at the end of the expression in the display.

Notes

- The examples on the transparency masters assume all default settings.
- All functions, except statistics, work in constant mode.
- To enter a constant:
 1. Press **[2nd] [K]**. If a constant is already stored, press **[CLEAR]** to clear it.
 2. Enter your constant (any set of operations, functions, and values).
 3. Press **[ENTER]** to turn on the constant mode. **K** appears in the display.
 4. Press **[CLEAR]** to clear the display.
 5. Enter an initial value. If you do not enter a value, 0 is assumed, and **Ans** will appear in the display.
 6. Press **[ENTER]** to place the contents of **K** at the end of the expression and evaluate it.
 7. Continue pressing **[ENTER]** to repeat the constant.

The result is stored in **Ans**, which is displayed, and the constant is used to evaluate the new expression.



Constant

2nd ^K ÷

Three people babysit for \$3.25 each per hour. First person works 16 hours. Second person works 12 hours. Third person works 17 hours. How much did each person earn?

Press

Display

2nd ^K ÷

K =
DEG

× 3.25 ENTER

K = *3.25
DEG K

CLEAR

DEG K

16 ENTER

16*3.25 ↑
52.
DEG K

12 ENTER

12*3.25 ↑
39.
DEG K

17 ENTER

17*3.25 ↑
55.25
DEG K

2nd ^K ÷

(Constant mode is off.)

DEG




Keys

1. $\square$ enters a decimal point.
2. 2^{nd} [FIX] displays the following menu that lets you set the number of decimal places.

F 0 1 2 3 4 5 6 7 8 9

F Sets floating decimal (standard) notation.

0-9 Sets number of decimal places.



Notes

- The examples on the transparency masters assume all default settings.
- 2^{nd} [FIX] $\square$ removes the setting and returns to standard notation (floating decimal).
- The **FIX** setting affects all decimal results and the mantissa of Scientific and Engineering notation results.
- The TI-30X IIS automatically rounds the result to the number of decimal places selected. For example, when the decimal is set to 2 places, 0.147 becomes 0.15 when you press $\square$. The TI-30X IIS also rounds or pads resulting values with trailing zeros to fit the selected setting. For example, when the decimal is set to 5 places, 0.147 becomes 0.14700 when you press $\square$.
- All results are displayed to the **FIX** setting until you clear the setting by either pressing 2^{nd} [FIX] $\square$ or selecting **F**(floating) on the decimal notation menu. Resetting the calculator also clears the **FIX** setting.
- After pressing 2^{nd} [FIX], you can select the number of decimal places in 2 ways:
 - Press $\uparrow$ or $\downarrow$ to move to the number of decimal places you want, and then press $\square$, or
 - Press the number key that corresponds to the number of decimal places you want.
- **FIX** affects only the results, not the entry.

Decimal, FIX

• 2nd • ^{FIX}

Round 12.345 to the hundredths place, to the tenths place, and then cancel the **FIX** setting.

Press

Display

12 • 345

12.345

DEG

^{FIX}
2nd •

F0123456789

DEG

▶ ▶ ▶

F0123456789

DEG

ENTER

12.345

FIX

DEG

ENTER

12.345

12.35

FIX

DEG

^{FIX}
2nd • 1

12.345

12.3

FIX

DEG

^{FIX}
2nd • •

12.345

12.345

DEG



Memory

7

Keys

1. **[STO▶]** displays the following menu of variables.

A B C D E Lets you select a variable in which to store the displayed value. The new variable replaces any previously stored value.

rand Lets you set a seed value for random integers.

2. **[MEMVAR]** displays the following menu of variables.

A B C D E Lets you view the stored value before pasting it in variable form to the display.


3. **[2nd] [CLRVAR]** clears all variables.

4. **[2nd] [RCL]** displays the following menu of variables.

A B C D E Lets you view the stored value before pasting it to the display.

Notes

- The examples on the transparency masters assume all default settings.
- You can store a real number or an expression that results in a real number to a memory variable.
- When you select a variable using **[MEMVAR]**, the variable name (**A**, **B**, **C**, **D**, or **E**) is displayed on the entry line.
- When you select a variable using **[2nd] [RCL]**, the value of the stored variable is displayed on the entry line.
- Resetting the calculator clears all memory variables.
- For more about **rand**, see Chapter 11, Probability.



Store, Memory Variable, Clear Variable

STO► **MEMVAR**

2nd CLRVAR
MEMVAR

Test scores: 96, 76, 85.

Weekly scores: 92, 83, 97, and 86.

Find the average of test and weekly scores. Find the final average.

Press

Display

96 **+** 76 **+**
85 **ENTER**

96+76+85
257.
DEG

÷ 3 **ENTER**

Ans/3
85.66666667
DEG

STO► **ENTER**

Ans→A
85.66666667
DEG

92 **+** 83 **+**
97 **+** 86 **ENTER**

92+83+97+86
358.
DEG

÷ 4 **ENTER**


Ans/4
89.5
DEG

+ **MEMVAR**
ENTER **ENTER**


Ans+A
175.1666667
DEG

÷ 2 **ENTER**

Ans/2
87.58333333
DEG



Store, Recall



Which would be the better buy: 3 cassette tapes for \$7.98, or 4 cassette tapes for \$9.48?


Press	Display
7 $\square$ 98 $\div$ 3 $\square$ ENTER	7.98/3 2.66 DEG
STO $\square$ ENTER	Ans $\rightarrow$ A 2.66 DEG
9 $\square$ 48 $\div$ 4 $\square$ ENTER	9.48 / 4 2.37 DEG
STO $\square$ $\rightarrow$ ENTER	Ans $\rightarrow$ B 2.37 DEG

View the first price again.

RCL 2nd STO $\square$	A B C D E 2.66 DEG
--------------------------	--------------------------

View the second price again.

$\rightarrow$	A B C D E 2.37 DEG
---------------	--------------------------



Store, Recall

STO▶

2nd ^{RCL} **STO▶**

<u>Shop</u>	<u>Purchases</u>	<u>Qty</u>	<u>Cost</u>
A	shirts	2	\$13.98 ea.
B	ties	3	\$7.98 ea.
C	belt	1	\$6.98
	suspenders	1	\$9.98

How much did you spend at each shop, and how much did you spend altogether?

Press

Display

2 **×** 13 **.** 98

ENTER

2*13.98

27.96

DEG

STO▶

A B C D E → ↑

27.96

DEG

ENTER

Ans→A ↑

27.96

DEG

3 **×** 7 **.** 98


ENTER

3*7.98 ↑

23.94

DEG

Continued



Store, Recall (Continued)

STO►

RCL
2nd **STO►**

Press

Display

STO► **►** **ENTER**

Ans→B ↑
23.94
DEG

6 **□** 98 **+**
9 **□** 98 **ENTER**

6.98+9.98 ↑
16.96
DEG

STO► **►** **►**
ENTER

Ans→C ↑
16.96
DEG

RCL
2nd **STO►**
ENTER **+**


27.96+ ↑
DEG

RCL
2nd **STO►** **►**
ENTER **+**

←.96+23.94+ → ↑
DEG


RCL
2nd **STO►** **►** **►**
ENTER **ENTER**

27.96+23.94 → ↑
68.86
DEG



Keys

1. **[Ab/c]** lets you enter mixed numbers and fractions.
2. **[2nd] [Ab/c ↔ d/e]** converts a simple fraction to a mixed number or a mixed number to a simple fraction.
3. **[2nd] [F ↔ D]** converts a fraction to its decimal equivalent or changes a decimal to its fractional equivalent, if possible.




Notes

- The examples on the transparency masters assume all default settings.
- To enter a mixed number or a fraction, press **[Ab/c]** between the whole number and the numerator and between the numerator and the denominator.
- You can enter a fraction or mixed number anywhere you can enter a decimal value.
- You can use fractions and decimals together in a calculation.
- Fractional results and entries are automatically reduced to their lowest terms.
- Fractional calculations can show fractional or decimal results.
 - When possible, calculations involving 2 fractions or a fraction and any integer will display fractional results.
 - Calculations involving a fraction and a decimal will always display results as decimals.
- For a mixed number, the whole number can be up to 3 digits, the numerator can be up to 3 digits, and the denominator can be any number through 1000.
- For a simple fraction, the numerator can be up to 6 digits and the denominator can be any number through 1000.

Fractions

A^{b/c}

At the party, you ate $\frac{5}{6}$ of the pepperoni pizza and $\frac{1}{10}$ of the sausage pizza. How much pizza did you eat?




Press

5 **A^{b/c}** 6 **+** 1
A^{b/c} 10 **=**

Display

5,6+1,10[↑]
 14 / 15
 DEG




Mixed Numbers

$\boxed{a\frac{b}{c}}$

A baby weighed $4\frac{3}{8}$ kilograms at birth. In the next 6 months, she gained $2\frac{3}{4}$ kilograms.

How much does she weigh?



Press

Display

4 $\boxed{a\frac{b}{c}}$ 3 $\boxed{a\frac{b}{c}}$
 8 $\boxed{+}$ 2 $\boxed{a\frac{b}{c}}$ 3
 $\boxed{a\frac{b}{c}}$ 4 $\boxed{\text{ENTER}}$

4 $\frac{3}{8}$ + 2 $\frac{3}{4}$ $\uparrow$
 7 $\frac{1}{8}$
 DEG



Mixed Number to Fraction, Fraction to Mixed Number

2nd $A^{b/c} \leftrightarrow d/e$
A^{b/c}

Sam is making his birthday cake.
The recipe calls for $3 \frac{1}{2}$ cups of
flour. He has only a $\frac{1}{2}$ -cup
measuring cup. To find out how
many times Sam must use his
measuring cup, change the mixed
number to a fraction.

$$3 \frac{1}{2} \div \frac{1}{2} = 7$$



Press

Display

3 **A^{b/c}** **1** **A^{b/c}** **2**

3 1 2

DEG

$A^{b/c} \leftrightarrow d/e$
2nd **A^{b/c}**

3 1 2 $A^{b/c} \leftrightarrow d/e$

DEG

ENTER

3 1 2 $A^{b/c} \leftrightarrow d/e$
7 / 2

DEG


Show the mixed number again.

$A^{b/c} \leftrightarrow d/e$
2nd **A^{b/c}**

Ans $A^{b/c} \leftrightarrow d/e$
3 1 2

DEG

ENTER



Fraction to Decimal

2nd **F $\leftrightarrow$ D**
PRB

Juan swims 20 laps in 5.72 minutes.
Mary swims 20 laps in $5 \frac{3}{4}$ minutes. Change Mary's time to a decimal to determine who swims faster.

Press

Display

5 **Ab/c** 3 **Ab/c**

5.3.4 **F $\leftrightarrow$ D**

4 **2nd** **F $\leftrightarrow$ D**
PRB

DEG

ENTER

5.3.4 **F $\leftrightarrow$ D**

5.75
DEG



Decimal to Fraction

2nd **F $\leftrightarrow$ D**
PRB

Change 2.25 to its fractional equivalent.

Press


Display

2 **.** 25

F $\leftrightarrow$ D
2nd **PRB** **ENTER**

2.25 **F $\leftrightarrow$ D**

2 **1**/**4**
DEG




Keys

1. π displays the value of pi rounded to 10 digits (3.141592654).

Notes

- The examples on the transparency masters assume all default settings.
- Internally, pi is stored to 13 digits (3.141592653590).
- After pressing 2nd [FIX] , you can select the number of decimal places in 2 ways:
 - Press ◀ or ▶ to move to the number of decimal places you want, and then press ENTER , or
 - Press the number key that corresponds to the number of decimal places you want.

The transparency masters show both ways.




Circumference



Use this formula to find the amount of border you need if you want to put a circular border all the way around the tree.

$$C = 2\pi r = 2 \times \pi \times 1.5\text{m}$$




Press

Display

2 $\times$ π $\times$ 1.5
 ENTER

2* π *1.5
 9.424777961
 DEG




Area



Use this formula to find how much of a lawn would be covered by the sprinkler. Round your answer to the nearest whole number, and then return to Floating Decimal mode.

$$A = \pi r^2 = \pi \times 4^2$$



Press

Display

π $\times$ 4 x^2
 ENTER

$\pi * 4^2$
 50.26548246
 DEG

FIX
 2nd $\square$ $\rightarrow$


F0123456789
 DEG

ENTER

$\pi * 4^2$ ↑
 50.
 FIX DEG

FIX
 2nd $\square$ $\square$

$\pi * 4^2$ ↑
 50.26548246
 DEG



Powers, Roots, and Reciprocals

10


Keys

1. x^2 squares the value.
2. 2^{nd} $\sqrt{}$ calculates the square root.
3. 2^{nd} $\sqrt[x]{}$ calculates the specified root (x) of the value.
4. x^{-1} calculates the reciprocal.
5. $\wedge$ raises a value to a specified power.

Notes

- The examples on the transparency masters assume all default settings.
- To use $\wedge$, enter the base, press $\wedge$, and then enter the exponent.
- The base (or mantissa) and the exponent may be either positive or negative. Refer to Domain under Error Messages in Appendix C for restrictions.
- The result of calculations with $\wedge$ must be within the range of the TI-30X IIS.
- A sign change takes precedence over exponents.

Example: $-5^2 = -25$
 $(-5)^2 = 25$




Squares

x^2 $\wedge$

Use this formula to find the size of the tarp needed to cover the entire baseball infield.

$$A = x^2 = 27.4^2$$



Press

Display

27.4 x^2 ENTER


or


27.4 $\wedge$ 2

ENTER

27.4²
750.76
DEG

90^{^2}
750.76
DEG




$$\boxed{2\text{nd}} \boxed{x^2}$$
$$L = \sqrt{x} = \sqrt{3}$$


Cubes



Use this formula to find the volume of a cube with sides 2.3 meters long. Change your answer to a fraction.

$$V = L^3 = 2.3^3$$



Press


Display

2 $\square$ 3 $\square$ 3
 $\square$ ENTER

2.3^3
 12.167
 DEG

F $\leftrightarrow$ D
 $\square$ 2nd $\square$ PRB
 $\square$ ENTER

Ans $\square$ F $\leftrightarrow$ D
 12.167/1000
 DEG



Powers



Fold a piece of paper in half, in half again, and so on until you cannot physically fold it in half again. How many sections would there be after 10 folds? After 15 folds?

Press


Display

2 $\wedge$ 10 ENTER


2^10
1024.
DEG

2 $\wedge$ 15 ENTER

2^15
32768.
DEG



Roots




If the volume of a cube is 125 cm^3 ,
what is the length of each side?

Press

Display

3 $\sqrt[3]{}$ 125
ENTER

3 $\sqrt[3]{}$ 125.
5.
DEG



Reciprocals

$$x^{-1}$$

The chart below shows the amount of time spent building model ships.

<u>Ships</u>	<u>Time Spent Building</u>	<u>Portion Completed Per Hour</u>
Sailing	10 hrs.	?
Steam	5 hrs.	?
Luxury	5 $\frac{1}{3}$ hrs.	?

How much of each model was completed per hour?

Press

Display

Sailing ship:

10 x^{-1} 2nd PRB
ENTER

10⁻¹ F4 D 1 / 10
DEG

Steam ship:


5 x^{-1} 2nd PRB
ENTER

5⁻¹ F4 D 1 / 5
DEG

Luxury liner:

5 $\frac{1}{3}$ 1 $\frac{1}{3}$ 3
 x^{-1} ENTER

5 1 3⁻¹ 1 / 16
DEG



Keys

1. **[PRB]** displays the following menu of functions.

nPr Calculates the number of possible permutations.

nCr Calculates the number of possible combinations.


! Calculates the factorial.

RAND Generates a random 10-digit real number between 0 and 1.

RANDI Generates a random integer between 2 numbers that you specify.

Notes

- The examples on the transparency masters assume all default settings.
- A *combination* is an arrangement of objects in which the order is not important, as in a hand of cards.
- A *permutation* is an arrangement of objects in which the order is important, as in a race.
- A *factorial* is the product of all the positive integers from 1 to n , where n is a positive whole number ≤ 69 .
- To control a sequence of random numbers, you can store (**[STO▶]**) an integer to **RAND** just as you would store values to memory variables. The seed value changes randomly when a random number is generated.
- For **RANDI**, use a comma to separate the 2 numbers that you specify.



Combination (nCr)

PRB

You have space for 2 books on your bookshelf. You have 4 books to put on the shelf. Use this formula to find how many ways you could place the 4 books in the 2 spaces.

$$4 \text{ nCr } 2 = x$$



A B C D

AB and BA count as only 1 combination.

AB	AC	AD
BA	BC	BD
CA	CB	CD
DA	DB	DC

Press

Display

4 **PRB** **►**

nPr nCr ! →
DEG

2 **ENTER**

4 nCr 2
6.
DEG




Permutation (nPr)

PRB

Four different people are running in a race. Use this formula to find how many different ways they can place 1st and 2nd.

$$4 \text{ nPr } 2 = x$$



AB and BA ——— count as 2 permutations.

AB	AC	AD
BA	BC	BD
CA	CB	CD
DA	DB	DC

Press


Display

4 **PRB**

nPr nCr ! →
DEG

2 **ENTER**

4 nPr 2 ↑
12.
DEG



Factorial (!)

PRB

Using the digits 1, 3, 7, and 9 only one time each, how many 4-digit numbers can you form?

$$4! = x$$

1 3 7 9
A B C D

ABCD	ABDC	ACBD	ACDB	ADBC	ADCB
BACD	BADC	BCAD	BCDA	BDCA	BDAC
CABD	CADB	CBAD	CBDA	CDAB	CDBA
DABC	DACB	DBAC	DBCA	DCAB	DCBA

Press

Display

4 **PRB** **►** **►**

nPr nCr **!** →
DEG

ENTER **ENTER**

4!
24.
DEG



Random (RAND)

PRB

Generate a sequence of random numbers.

Press

Display

PRB → → →

← RAN RANDI
DEG

ENTER ENTER

RAND ↑
0.839588694
DEG

ENTER ENTER

RAND ↑
0.482688185
DEG

Results will vary.



Random (RAND)

PRB

Set 1 as the current seed and generate a sequence of random numbers.

Press

Display

1 **STO►** **◀**

← rand
1083958869.
DEG

ENTER **ENTER**

1→rand ↑
1.
DEG

PRB **▶** **▶** **▶**
ENTER **ENTER**

RAND ↑
0.000018633
DEG

ENTER

RAND ↑
0.745579721
DEG



Random Integer (RANDI)

PRB

Generate a random integer from 2 through 10.

Press

Display

PRB $\leftarrow$

$\leftarrow$ RAND RANDI
DEG

ENTER 2 2nd)

$\leftarrow$ RANDI(2, 10) $\uparrow$
DEG

10)

ENTER

RANDI(2, 10) $\rightarrow$ $\uparrow$
3.
DEG

Results will vary.




Keys

1. **[2nd] [STAT]** displays a menu from which you can select **1-VAR**, **2-VAR** or **CLRDATA**.

1-VAR	Analyzes data from 1 set of data with 1 measured variable— x .
2-VAR	Analyzes paired data from 2 sets of data with 2 measured variables— x , the independent variable, and y , the dependent variable.
CLRDATA	Clears data values without exiting STAT mode.

2. **[DATA]** lets you enter data points (x for **1-VAR** stats; x and y for **2-VAR** stats).



3. **[2nd] [EXIT STAT]** displays the following menu that lets you clear data values and exit **STAT** mode.

EXIT ST: Y N

- Press **[ENTER]** when **Y** (yes) is underlined to clear data values and exit **STAT** mode.
- Press **[ENTER]** when **N** (no) is underlined to return to the previous screen without exiting **STAT** mode.

4. **[STATVAR]** displays the menu of variables with their current values.

n	Number of x (or x,y) data points.
$\bar{x}$ or $\bar{y}$	Mean of all x or y values.
S_x or S_y	Sample standard deviation of x or y .
σ_x or σ_y	Population standard deviation of x or y .
Σx or Σy	Sum of all x values or y values.
Σx^2 or Σy^2	Sum of all x^2 values or y^2 values.
Σxy	Sum of $(x \times y)$ for all xy pairs in 2 lists.
a	Linear regression slope.
b	Linear regression y -intercept.
r	Correlation coefficient.

Notes

- The examples on the transparency masters assume all default settings.
- To save the last data point or frequency value entered, you must press **[ENTER]** or **[∇]**.
- You can change data points once they are entered.

Entering 1-VAR Stat Data

2nd ^{STAT}**DATA** **DATA**

Five students took a math test.
Using their scores, enter the data
points—85, 85, 97, 53, 77.

Press

Display

2nd ^{STAT}**DATA**

1-VAR_ 2-VAR →
DEG

ENTER **DATA**

X₁=
STAT DEG

85

X₁=85
STAT DEG

⏴

FRQ=1
STAT DEG

2

FRQ=2
STAT DEG

⏴ 97

X₂=97
STAT DEG


⏴ ⏴ 53

X₃=53
STAT DEG

⏴ ⏴ 77 **ENTER**

X₄=77
STAT DEG 77.

Continued



Viewing the Data (Cont.)

STATVAR

Find the number of data points (**n**), the mean ($\bar{x}$), the sample standard deviation (**Sx**), the population standard deviation (σx), the sum of the scores (Σx), and the sum of the squares (Σx^2).

Press

Display

STATVAR

$\bar{n}$ $\bar{x}$ **Sx** σx →
STAT DEG
5.



n $\bar{x}$ **Sx** σx →
STAT DEG
79.4



n $\bar{x}$ **Sx** σx →
STAT DEG
16.39512123




n $\bar{x}$ **Sx** σx →
STAT DEG
14.66424222



$\leftarrow \Sigma x$ Σx^2
STAT DEG
397.



$\leftarrow \Sigma x$ Σx^2
STAT DEG
32597.




Continued

Removing Data Points (Cont.)

2nd EXIT STAT
STATVAR

Return to the first data point.
Display the lowest score, drop it,
and then find the new mean ($\bar{X}$).
Clear all data by exiting **STAT** mode.

Press	Display
DATA	$X_1=85$ STAT DEG
$\downarrow \downarrow \downarrow \downarrow$	$X_3=53$ STAT DEG
$\downarrow 0$ ENTER	$FRQ=0$ STAT DEG
STATVAR $\rightarrow$	$n \quad \bar{X} \quad Sx \quad \sigma X$ STAT DEG 86.
2nd EXIT STAT STATVAR	EXIT ST: $\underline{Y}$ N STAT DEG
ENTER	 DEG



Entering 2-VAR Stat Data

STAT
2nd DATA DATA

The table below shows the number of pairs of athletic shoes sold by a small shoe store. Enter this data as the data points.

Month	Total No.(x)	Brand A (y)
April	58 (x1)	35 (y1)
May	47 (x2)	28 (y2)

Press

Display

2nd STAT DATA

1-VAR 2-VAR→
DEG

ENTER DATA

X1=
STAT DEG

58

X1=58
STAT DEG

35

Y1=35
STAT DEG

47


X2=47
STAT DEG

28

Y2=28
STAT DEG

ENTER

Y2=28
STAT DEG 28.



Continued

Viewing the Data (Cont.)

STATVAR

2nd EXIT STAT
STATVAR

If the store sells 32 pairs of shoes in June, predict the June sales of Brand A. When finished, exit **STAT** mode and clear all data points.

Press

Display

STATVAR 

$\leftarrow x' \quad y'$
[STAT] DEG

ENTER 32 **)**

$y' (32)$
18.45454545
[STAT] DEG


ENTER

EXIT STAT
2nd **STATVAR**

EXIT ST: y' N
[STAT] DEG

ENTER


DEG




Keys

1. **[TAN]** calculates the tangent.
2. **[2nd] [TAN⁻¹]** calculates the inverse tangent.
3. **[COS]** calculates the cosine.
4. **[2nd] [COS⁻¹]** calculates the inverse cosine.
5. **[SIN]** calculates the sine.
6. **[2nd] [SIN⁻¹]** calculates the inverse sine.

Notes


- The examples on the transparency masters assume all default settings.
- Before starting a trigonometric calculation, be sure to select the appropriate angle mode setting (**degree**, **radian**, or **gradient**—See Chapter 16, Angle Settings and Conversions). The calculator interprets values according to the current angle-unit mode setting.
- **[)]** ends a trig function.



Tangent

TAN

Use this formula to find the distance from the lighthouse to the boat. Round your answer to the nearest whole number, and then return to floating decimal mode.



Press

Display

78 $\div$ TAN
27 $)$ ENTER

78/tan (27)
153.0836194
DEG

FIX
2nd $\square$ $\rightarrow$

F0123456789
DEG

ENTER

78/tan (27)
153.
FIX DEG

FIX
2nd $\square$ $\square$

78/tan (27)
153.0836194
DEG




Inverse Tangent

2nd ^{TAN⁻¹}
TAN

Use this formula to find the angle of depression. Round your answer to the nearest tenth, and then return to floating decimal mode.

$$\text{TAN } x = 600/2500$$



Press

Display

^{TAN⁻¹}
2nd **TAN** 600 **÷**
2500 **)** **ENTER**

$\tan^{-1}(600/25 \rightarrow$
13.49573328
DEG

^{FIX}
2nd **.** **▶** **▶**


F0123456789
DEG

ENTER

$\tan^{-1}(600/25 \rightarrow^t$
13.5
FIX DEG

^{FIX}
2nd **.** **.**

$\tan^{-1}(600/25 \rightarrow^t$
13.49573328
DEG




Cosine

COS

Use this formula to find how far the base of the ladder is from the house. Round your answer to the nearest whole number, and then return to floating decimal mode.

$$D = 5 \times \cos 75$$



Press

Display

5 $\times$ COS
75) ENTER

5*cos (75) ↑
1.294095226
DEG

FIX
2nd . ►

F0123456789
FIX DEG


ENTER

5*cos (75) ↑
1.
FIX DEG

FIX
2nd . .

5*cos (75) ↑
1.294095226
DEG



$$\boxed{2\text{nd}} \boxed{\cos}^{-1}$$
$$\cos x = 453/500$$


Display

$\cos^{-1}(453/50) \rightarrow$
25.04169519
DEG

F	0	1	2	3	4	5	6	7	8	9
DEG										

$\cos^{-1}(453/50) \rightarrow$ [↑]
25.0
FIX DEG


$$\cos^{-1}(453/500) \rightarrow 25.04169519^\circ$$


Sine

SIN

Use this formula to find the length of the ramp. Round your answer to the nearest whole number, and then return to floating decimal mode.

$$D = 1.5 / \sin 12$$



Press

Display

1 $\square$. 5 $\square$ $\div$ $\square$ SIN
12 $\square$) $\square$ ENTER

1.5/sin (12) $\rightarrow$ $\uparrow$
7.214601517
DEG

FIX
2nd $\square$ $\square$ $\rightarrow$


F0123456789
DEG

ENTER

1.5/sin (12) $\rightarrow$ $\uparrow$
7.
FIX DEG

FIX
2nd $\square$. $\square$

1.5/sin (12) $\rightarrow$ $\uparrow$
7.214601517
DEG




Inverse Sine

2nd SIN^{-1} **SIN**

Use this formula to find the angle of the conveyor belt. Round your answer to the nearest tenth, and then return to floating decimal mode.

$$\text{SIN } x = 13/20$$



Press

Display

SIN^{-1}
2nd **SIN** 13 $\div$
20 **)** **ENTER**

$\text{sin}^{-1}(13/20) \rightarrow$ $\uparrow$
40.54160187
DEG

FIX
2nd **.** $\rightarrow$ $\rightarrow$

F0123456789
DEG

ENTER

$\text{sin}^{-1}(13/20) \rightarrow$ $\uparrow$
40.5
FIX DEG

FIX
2nd **.** **.**

$\text{sin}^{-1}(13/20) \rightarrow$ $\uparrow$
40.54160187
DEG



Keys


1. **[2nd] [SCI/ENG]** displays the following numeric notation mode menu.

FLO	Restores standard mode (floating decimal).
SCI	Turns on scientific mode and displays results as a number from 1 to 10 ($1 \leq n < 10$) times 10 to an integer power.
ENG	Turns on engineering mode and displays results as a number from 1 to 1000 ($1 \leq n < 1000$) times 10 to an integer power. The integer power is always a multiple of 3.

2. **[2nd] [EE]** lets you enter and calculate the exponent.

Notes

- The examples on the transparency masters assume all default settings.
- You can enter a value in scientific notation regardless of the numeric notation mode setting. For a negative exponent, press **[(-)]** before entering it.
- Results requiring more than 10 digits are automatically displayed in scientific notation.
- For the Decimal notation mode, refer to **[2nd] [FIX]** in Chapter 6, Decimals and Decimal Places.
- These modes (**FLO**, **SCI**, and **ENG**) affect only the display of results.



Engineering, Scientific, Floating Decimal

SCI/ENG
2nd DRG

Enter 12543, which will be in floating decimal notation (default), and alternate between scientific and engineering notations.

Press	Display
12543	FLO SCI ENG DEG
SCI/ENG 2nd DRG ►	
ENTER ENTER	12543 ↑ 1.2543 × 10 ⁰⁴ SCI DEG
SCI/ENG 2nd DRG ►	FLO SCI ENG SCI DEG
ENTER	12543 ↑ 12.543 × 10 ⁰³ ENG DEG
SCI/ENG 2nd DRG ►	FLO SCI ENG ENG DEG
ENTER	12543 ↑ 12543. DEG



$$\boxed{2\text{nd}} \boxed{\text{EE}} \boxed{x^{-1}}$$

Press

$$7 \square \cdot 783$$
$$\text{EE}$$
$$\boxed{2\text{nd}} \boxed{x^{-1}} 8$$

7.783E8-1.4 → [†]
628700000.
DEG


$$\boxed{-} 1 \boxed{\cdot} 496$$
$$\boxed{2\text{nd}} \boxed{x^{-1}} 8$$
$$\boxed{\text{ENTER}}$$


Keys

1. **[LOG]** calculates the common logarithm (base 10).
2. **[LN]** calculates the natural logarithm (base e , where $e = 2.718281828459$).
3. **[2nd] [10^x]** calculates the common antilogarithm (10 raised to the power of the value).
4. **[2nd] [e^x]** calculates the natural antilogarithm (e raised to the power of the value).

Note

- The examples on the transparency masters assume all default settings.
- **[)]** ends a logarithmic function.



Common Logarithm, Natural Logarithm

LOG **LN**

Find $\log 23$ rounded to 4 decimal places. Then find $\ln 23$ rounded to 4 decimal places and return to Floating Decimal notation.

Press

Display

LOG 23 **)**

ENTER

$\log (23)$
1.361727836
DEG

2nd ^{FIX} **.**

$\overline{\text{E}}0123456789$
DEG

4

$\log (23)$
1.3617
FIX DEG


LN 23 **)**

ENTER

$\ln (23)$
3.1355
FIX DEG

2nd ^{FIX} **.** **.**

$\ln (23)$
3.135494216
DEG




Common Antilogarithm, Natural Antilogarithm

$$\boxed{2\text{nd}} \boxed{\text{LOG}}^{10^x}$$

$$\boxed{2\text{nd}} \boxed{\text{LN}}^{e^x}$$

Find antilog 3.9824 rounded to 4 decimal places. Then find antiln 3.9824 rounded to 4 decimal places. When finished, return to Floating Decimal notation.

Press	Display
$\boxed{2\text{nd}} \boxed{\text{LOG}}^{10^x} 3 \boxed{\cdot}$ $9824 \boxed{)} \boxed{\text{ENTER}}$	$10^{\wedge} (3.9824) \uparrow$ 9602.846792 DEG
$\boxed{2\text{nd}} \boxed{\cdot}^{\text{FIX}}$	$\text{E} 0123456789$ DEG
4	$10^{\wedge} (3.9824) \uparrow$ 9602.8468 FIX DEG
$\boxed{2\text{nd}} \boxed{\text{LN}}^{e^x} 3 \boxed{\cdot}$ $9824 \boxed{)} \boxed{\text{ENTER}}$	$e^{\wedge} (3.9824) \uparrow$ 53.6456 FIX DEG
$\boxed{2\text{nd}} \boxed{\cdot}^{\text{FIX}} \boxed{\cdot}$	$e^{\wedge} (3.9824) \uparrow$ 53.64562936 DEG



Angle Settings and Conversions

16

Keys

1. **DRG** displays the following menu that lets you change the angle mode setting to **DEG**, **RAD**, and **GRD** without affecting the value in the display.

DEG Sets degree mode.

RAD Sets radian mode.

GRD Sets gradient mode.

When you turn on the TI-30X IIS, it is always in the **DEG** mode.

2. **° ' "** displays a menu that lets you specify the unit of an angle.

° Specifies degrees.


r Specifies radians.

g Specifies gradients.

DMS Specifies degrees ($^{\circ}$), minutes ($'$), and seconds ($''$). It also lets you convert an angle from decimal degrees to **DMS** notation.

Notes

- The examples on the transparency masters assume all default settings.
- Angles with a trig function ignore the angle mode setting and display results in the original unit. Otherwise, angles (without a trig function) are converted and displayed according to the angle mode setting.
- You enter decimal-degree angles the same as you would any other number.
- For decimal/**DMS** conversions, the calculator interprets all values as degrees, regardless of the angle-unit setting.
- **DMS** angles are entered as $^{\circ}$ (degrees), $'$ (minutes), and $''$ (seconds).




Degrees, Minutes, and Seconds to Decimal



You watched 2 videos that were 2:05 (2 hours and 5 minutes) and 1:46 (1 hour and 46 minutes) in length. How long did you watch videos?

Press	Display
2	 DEG
	 DEG
5	 DEG
1 	 DEG
46 	 DEG
	 DEG
	 DEG



Fraction to Degrees, Minutes, and Seconds



How much is $\frac{2}{3}$ of an hour in hours, minutes, and seconds?

Press

Display

2 $\boxed{\text{Ab/c}}$ 3


2 1/3 ↑
DEG

$\boxed{0 ///}$ $\boxed{\leftarrow}$

← DMS
DEG

$\boxed{\text{ENTER}}$ $\boxed{\text{ENTER}}$

2 1/3 DMS ↑
0° 40' 0"
DEG



Degrees, Radians, Gradients

DRG

Calculate the sine of 30 in degrees, radians, and gradients, and then return to degrees.

Press	Display
SIN 30) ENTER	sin(30) ↑ 0.5 DEG
DRG ▶	DEG RAD GRD DEG
ENTER ENTER	sin(30) ↑ -0.988031624 RAD
DRG ▶	DEG RAD GRD RAD
ENTER ENTER	sin(30) ↑ 0.4539905 GRAD
DRG ▶ ENTER ENTER	sin(30) ↑ 0.5 DEG



Polar and Rectangular Conversions

17

Keys

1. **[2nd] [R↔P]** displays the following menu that lets you convert rectangular coordinates (x,y) to polar coordinates (r,θ) or vice versa.

R↔Pr Converts rectangular coordinate to polar coordinate r .

R↔Pθ Converts rectangular coordinate to polar coordinate θ .


P↔Rx Converts polar coordinate to rectangular coordinate x .

P↔Ry Converts polar coordinate to rectangular coordinate y .

2. **[2nd] [,]** enters a comma.

Notes


- The example on the transparency master assumes all default settings.
- Before starting calculations, set angle mode as necessary.



Polar to Rectangular

2nd $R \leftrightarrow P$ $\circ / \prime \prime$

Convert the polar ordered pair (7,30) to rectangular using the **DEG** ($^{\circ}$) angle unit.



Press

Display

2nd $R \leftrightarrow P$ $\circ / \prime \prime$ $\rightarrow$ $\rightarrow$

$\leftarrow$ P $\rightarrow$ Rx P $\rightarrow$ Ry
DEG

$\underline{\text{ENTER}}$ 7 **2nd** $\circ / \prime \prime$
30 $\circ / \prime \prime$ $\underline{\text{ENTER}}$

P $\rightarrow$ Rx(7,30) $\uparrow$
6.062177826
DEG


2nd $R \leftrightarrow P$ $\circ / \prime \prime$ $\leftarrow$

$\leftarrow$ P $\rightarrow$ Rx P $\rightarrow$ Ry
DEG

$\underline{\text{ENTER}}$ 7 **2nd** $\circ / \prime \prime$
30 $\circ / \prime \prime$ $\underline{\text{ENTER}}$


P $\rightarrow$ Ry(7,30) $\uparrow$
3.5
DEG

The rectangular ordered pair is 6.062177826,3.5.



Notes

1. **2nd** **[HYP]** accesses the hyperbolic (**sinh**, **cosh**, **tanh**) function of the next trig key that you press.
- The example on the transparency master assumes all default settings.
 - Hyperbolic calculations are not affected by the angle mode setting—whether or not the calculator is in **RAD** (radian), **GRD** (gradient), or **DEG** (degree) modes.



Sine, Cosine, Tangent

HYP
2nd π

Find the hyperbolic sine (**sinh**),
cosine (**cosh**), and tangent (**tanh**)
of 5.

Press

Display

HYP
2nd π SIN 5
) ENTER


sinh(5) ↑
74.20321058
DEG

HYP
2nd π COS 5
) ENTER

cosh(5) ↑
74.20994852
DEG

HYP
2nd π TAN 5
) ENTER

tanh(5) ↑
0.999909204
DEG



Quick Reference to Keys

A

Key	Function
$\leftarrow \rightarrow$	Moves the cursor left and right so you can scroll the entry line. Press 2nd $\leftarrow$ or 2nd $\rightarrow$ to scroll to the beginning or end of the entry line.
$\uparrow \downarrow$	Moves the cursor up and down so you can see previous entries. Press 2nd $\uparrow$ or 2nd $\downarrow$ to scroll to the beginning or end of the history.
$+$ $-$ $\times$ $\div$	Adds, subtracts, multiplies, and divides.
0 $-$ 9	Enters the digits 0 through 9.
$($	Opens a parenthetical expression.
$)$	Closes a parenthetical expression.
x^{-1}	Calculates the reciprocal.
x^2	Squares the value.
π	Enters the value of pi rounded to 10 digits (3.141592654).
$.$	Enters a decimal point.
$(-)$	Indicates the value is negative.
$\wedge$	Raises a value to a specified power.
$\square^{\circ}'''$	Displays the following menu that lets you specify the unit of an angle. $^{\circ}$ Specifies degrees. r Specifies radians. g Specifies gradients. DMS Specifies degrees ($^{\circ}$), minutes ($'$), and seconds ($''$). It also lets you convert an angle from decimal degrees to DMS notation.
2nd	Turns on the 2nd indicator and accesses the function shown above the next key that you press.
2nd $[10^x]$	Calculates the common antilogarithm (10 raised to the power of the value).
2nd $[\sqrt{}]$	Calculates the square root.
2nd $[\%]$	Changes a real number to percent. Results display according to the Decimal Notation mode setting.
2nd $[,]$	Enters a comma.
2nd $[\sqrt[x]{}]$	Calculates the specified root (x) of the value.

Quick Reference to Keys (Continued)

A

Key	Function
$\boxed{A\frac{b}{c}}$	Lets you enter mixed numbers and fractions.
$\boxed{2nd} \boxed{A\frac{b}{c} \leftrightarrow d/e}$	Converts a simple fraction to a mixed number or a mixed number to a simple fraction.
$\boxed{2nd} \boxed{ANS}$	Recalls the most recently calculated result, displaying it as Ans .
$\boxed{CLEAR}$	Clears characters and error messages on the entry line. Once the display is clear, it moves the cursor to the last entry in history.
$\boxed{2nd} \boxed{CLRVAR}$	Clears all memory variables.
$\boxed{COS}$	Calculates the cosine.
$\boxed{2nd} \boxed{COS^{-1}}$	Calculates the inverse cosine.
$\boxed{DATA}$	Lets you enter the statistical data points (x for 1-VAR stats; x and y for 2-VAR stats).
$\boxed{DEL}$	Deletes the character at the cursor. If you hold $\boxed{DEL}$ down, it deletes all characters to the right. Then each time you press $\boxed{DEL}$, it deletes 1 character to the left of the cursor.
$\boxed{DRG}$	Displays the following menu that lets you change the Angle mode to degrees ($^{\circ}$), radians (r), or gradients (g), and then back to degrees without affecting the value in the display. DEG Sets degree mode. RAD Sets radian mode. GRD Sets gradient mode. When you turn on the TI-30X IIS, it is always in the DEG mode.
$\boxed{2nd} \boxed{e^x}$	Calculates the natural antilogarithm (e raised to the power of the value).
$\boxed{2nd} \boxed{EE}$	Lets you enter and calculate the exponent.
$\boxed{ENTER}$	Completes the operation or executes the command.
$\boxed{2nd} \boxed{EXIT STAT}$	Displays the following menu that lets you clear data values and exit STAT mode. EXIT ST: <u>Y</u> N Press $\boxed{ENTER}$ when Y (yes) is underlined to clear data values and exit STAT mode. Press $\boxed{ENTER}$ when N (no) is underlined to return to the previous screen without exiting STAT mode.

Quick Reference to Keys (Continued)

A

Key	Function
[2nd] [F↔D]	Converts a fraction to its decimal equivalent or converts a decimal to its fractional equivalent, if possible.
[2nd] [FIX]	Displays the following menu that lets you set the number of decimal places. F 0 1 2 3 4 5 6 7 8 9 F Sets floating decimal (standard) notation. 0-9 Sets number of decimal places.
[2nd] [HYP]	Accesses the hyperbolic (sinh , cosh , tanh) function of the next trig key that you press.
[2nd] [INS]	Lets you insert a character at the cursor.
[2nd] [K]	Turns on the constant mode and lets you define a constant.
[LN]	Calculates the natural logarithm (base e , where $e = 2.718281828459$).
[LOG]	Calculates the common logarithm (base 10).
[MEMVAR]	Displays the following menu of variables. A B C D E Lets you view the stored value before pasting it to the display.
[2nd] [OFF]	Turns off the calculator and clears the display.
[ON]	Turns on the calculator.
[PRB]	Displays the following menu of functions. nPr Calculates the number of possible permutations. nCr Calculates the number of possible combinations. ! Calculates the factorial. RAND Generates a random 10-digit real number between 0 and 1. RANDI Generates a random integer between 2 numbers that you specify. Separate the 2 numbers with a comma.
[2nd] [RCL]	Recalls the stored values to the display.

Quick Reference to Keys (Continued)

A

Key	Function
[2nd] [RESET]	Displays the RESET menu. RESET: <u>N</u> Y Press [ENTER] when N (no) is underlined to return to the previous screen without resetting the calculator. Press [ENTER] when Y (yes) is underlined to reset the calculator. The message MEM CLEARED is displayed. Also, press [ON] and [CLEAR] simultaneously to reset the calculator immediately. No menu or message is displayed.
[2nd] [R↔P]	Displays the following menu that lets you convert rectangular coordinates (x,y) to polar coordinates (r,θ) or vice versa. R→Pr Converts rectangular coordinate to polar coordinate r. R→Pθ Converts rectangular coordinate to polar coordinate θ. P→Rx Converts polar coordinate to rectangular coordinate x. P→Ry Converts polar coordinate to rectangular coordinate y.
[2nd] [SCI/ENG]	Displays the following numeric notation mode menu. FLO Restores standard mode (floating decimal). SCI Turns on scientific mode and displays results as a number from 1 to 10 ($1 \leq n < 10$) times 10 to an integer power. ENG Turns on engineering mode and displays results as a number from 1 to 1000 ($1 \leq n < 1000$) times 10 to an integer power. The integer power is always a multiple of 3.
[SIN]	Calculates the sine.
[2nd] [SIN⁻¹]	Calculates the inverse sine.
[2nd] [STAT]	Displays the following menu from which you can select 1-VAR, 2-VAR, or CLRDATA. 1-VAR Analyzes data from 1 set of data with 1 measured variable—x. 2-VAR Analyzes paired data from 2 sets of data with 2 measured variables—x, the independent variable, and y, the dependent variable. CLRDATA Clears data values without exiting STAT mode.

Quick Reference to Keys (Continued)

A

Key	Function
[STATVAR]	Displays the following menu of stat variables with their current values. n Number of x (or x,y) data points. $\bar{x}$ or $\bar{y}$ Mean of all x or y values. s_x or s_y Sample standard deviation of x or y. σ_x or σ_y Population standard deviation of x or y. Σx or Σy Sum of all x values or y values. Σx^2 or Σy^2 Sum of all x² values or y² values. Σxy Sum of (x × y) for all xy pairs in 2 lists. a Linear regression slope. b Linear regression y-intercept. r Correlation coefficient.
[STO▶]	Displays the following menu of variables. A B C D E Lets you select a variable in which to store the displayed value. The new variable replaces any previously stored value. rand Lets you set a seed value for random integers.
[TAN]	Calculates the tangent.
[2nd] [TAN⁻¹]	Calculates the inverse tangent.

Display Indicators

B

Indicator	Meaning
2nd	2nd function.
HYP	Hyperbolic function.
FIX	Fixed-decimal setting.
SCI, ENG	Scientific or engineering notation.
STAT	Statistical mode.
DEG, RAD, GRAD	Angle mode (degrees, radians, or gradients).
K	Constant mode.
x10	Precedes the exponent in scientific or engineering notation.
↑ ↓	An entry is stored in history before and/or after the active screen. Press  and  to scroll.
← →	An entry or menu displays beyond 11 digits. Press  or  to scroll.

Error Messages

C

Message	Meaning
ARGUMENT	A function does not have the correct number of arguments.
DIVIDE BY 0	- You attempted to divide by 0. - In statistics, $n = 1$.
DOMAIN	You specified an argument to a function outside the valid range. For example: - For $x\sqrt{}$: $x = 0$ or $y < 0$ and x is not an odd integer. - For y^x: y and $x = 0$; $y < 0$ and x is not an integer. - For $\sqrt{x}$: $x < 0$. - For LOG or LN: $x \leq 0$. - For TAN: $x = 90^\circ, -90^\circ, 270^\circ, -270^\circ, 450^\circ$, etc. - For SIN⁻¹ or COS⁻¹: $x > 1$. - For nCr or nPr: n or r are not integers ≥ 0. - For $x!$: x is not an integer between 0 and 69.
EQUATION LENGTH ERROR	An entry exceeds the digit limits (88 for entry line and 47 for statistics or constant entry lines); for example, combining an entry with a constant that exceeds the limit.
FRQ DOMAIN	FRQ value (in 1-variable statistics) < 0 or > 99 , or not an integer..
OVERFLOW	$ \theta \geq 1E10$, where θ is an angle in a trig, hyperbolic, or RPr function.
STAT	- You pressed [STATVAR] with no defined data points. - You pressed [DATA], [STATVAR], or [2nd] [EXIT STAT] when not in STAT mode. - Statistical analyses do not have at least 2 data points ($n > 1$).
SYNTAX	The command contains a syntax error—entering more than 23 pending operations, 8 pending values, or having misplaced functions, arguments, parentheses, or commas.

Support and Service Information

D

Product Support

Customers in the U.S., Canada, Puerto Rico, and the Virgin Islands

For general questions, contact Texas Instruments Customer Support:

phone: **1.800.TI.CARES (1.800.842.2737)**

e-mail: **ti-cares@ti.com**

For technical questions, call the Programming Assistance Group of Customer Support:

phone: **1.972.917.8324**

Customers outside the U.S., Canada, Puerto Rico, and the Virgin Islands

Contact TI by e-mail or visit the TI calculator home page on the World Wide Web.

e-mail: **ti-cares@ti.com**

internet: **www.ti.com/calc**

Product Service

Customers in the U.S. and Canada Only

Always contact Texas Instruments Customer Support before returning a product for service.

Customers outside the U.S. and Canada

Refer to the leaflet enclosed with this product or contact your local Texas Instruments retailer/distributor.

Other TI Products and Services

Visit the TI calculator home page on the World Wide Web.

www.ti.com/calc

Warranty Information

E

Customers in the U.S. and Canada Only

One-Year Limited Warranty for Electronic Product

This Texas Instruments ("TI") electronic product warranty extends only to the original purchaser and user of the product.

Warranty Duration. This TI electronic product is warranted to the original purchaser for a period of one (1) year from the original purchase date.

This TI electronic product is warranted against defective materials and construction. **THIS WARRANTY IS VOID IF THE PRODUCT HAS BEEN DAMAGED BY ACCIDENT OR UNREASONABLE USE, NEGLIGENCE, IMPROPER SERVICE, OR OTHER CAUSES NOT ARISING OUT OF DEFECTS IN MATERIALS OR CONSTRUCTION.**

Warranty Disclaimers. ANY IMPLIED WARRANTIES ARISING OUT OF THIS SALE, INCLUDING BUT NOT LIMITED TO THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE, ARE LIMITED IN DURATION TO THE ABOVE ONE-YEAR PERIOD. TEXAS INSTRUMENTS SHALL NOT BE LIABLE FOR LOSS OF USE OF THE PRODUCT OR OTHER INCIDENTAL OR CONSEQUENTIAL COSTS, EXPENSES, OR DAMAGES INCURRED BY THE CONSUMER OR ANY OTHER USER.

Some states/provinces do not allow the exclusion or limitation of implied warranties or consequential damages, so the above limitations or exclusions may not apply to you.

Legal Remedies. This warranty gives you specific legal rights, and you may also have other rights that vary from state to state or province to province.

Warranty Performance. During the above one (1) year warranty period, your defective product will be either repaired or replaced with a reconditioned model of an equivalent quality (at TI's option) when the product is returned, postage prepaid, to Texas Instruments Service Facility. The warranty of the repaired or replacement unit will continue for the warranty of the original unit or six (6) months, whichever is longer. Other than the postage requirement, no charge will be made for such repair and/or replacement. TI strongly recommends that you insure the product for value prior to mailing.

Software. Software is licensed, not sold. TI and its licensors do not warrant that the software will be free from errors or meet your specific requirements. **All software is provided "AS IS."**

Copyright. The software and any documentation supplied with this product are protected by copyright.

For information about the length and terms of the warranty, refer to your package and/or to the warranty statement enclosed with this product, or contact your local Texas Instruments retailer/distributor.

All Customers outside the U.S. and Canada