

## JHeimbach LLC



August 17, 2018

Paulette Gaynor, Ph.D.  
Senior Regulatory Project Manager  
Division of Biotechnology and GRAS Notice Review (HFS-255)  
Office of Food Additive Safety  
Center for Food Safety and Applied Nutrition  
Food and Drug Administration  
5100 Paint Branch Parkway  
College Park, MD 20740

Dear Dr. Gaynor:

Pursuant to 21 CFR Part 170, Subpart E, Arla Foods Ingredients Group P/S (Arla), through me as its agent, hereby provides notice of a claim that the addition of *Lactobacillus paracasei* ssp. *paracasei* strain F19 to powdered milk-based non-exempt term infant formula is exempt from the premarket approval requirement of the Federal Food, Drug, and Cosmetic Act because Arla has determined that the intended use is generally recognized as safe (GRAS) based on scientific procedures.

As required, one copy of the GRAS monograph and one signed copy of the conclusion from each member of the Expert Panel are provided. Additionally, I have enclosed a virus-free CD-ROM with the GRAS monograph and the signed statements of the Expert Panel.

If you have any questions regarding this notification, please feel free to contact me at 804-742-5543 or [jh@jheimbach.com](mailto:jh@jheimbach.com).

Sincerely, 

  
James T. Heimbach, Ph.D., F.A.C.N.  
President

Encl.

**Generally Recognized As Safe (GRAS) Determination for the  
Intended Use of *Lactobacillus paracasei* ssp. *paracasei* strain F-19**

**Prepared for:  
Arla Foods Ingredients Group P/S  
Basking Ridge NJ**

**Prepared by:  
JHeimbach LLC  
Port Royal Virginia**

**August, 2018**

## Table of Contents

|                                                                                                  |    |
|--------------------------------------------------------------------------------------------------|----|
| Table of Contents .....                                                                          | 1  |
| Part 1: Signed Statements and Certification .....                                                | 3  |
| 1.1. GRAS Notice Submission.....                                                                 | 3  |
| 1.2. Name and Address of Notifier .....                                                          | 3  |
| 1.3. Name of Notified Organism.....                                                              | 3  |
| 1.4. Intended Conditions of Use.....                                                             | 3  |
| 1.5 Statutory Basis for GRAS Status .....                                                        | 3  |
| 1.6. Premarket Exempt Status .....                                                               | 4  |
| 1.7. Data Availability .....                                                                     | 4  |
| 1.8. Freedom of Information Act Statement .....                                                  | 4  |
| 1.9. Certification .....                                                                         | 4  |
| 1.10 FSIS Statement.....                                                                         | 4  |
| 1.11. Name, Position and Signature of Notifier .....                                             | 4  |
| Part 2: Identity, Methods of Manufacture, Specifications, and Physical and Technical Effect..... | 5  |
| 2.1. Name of the GRAS Organism .....                                                             | 5  |
| 2.2. Source, Description, Manufacture, and Specifications of the GRAS Organism .....             | 5  |
| 2.2.1. Source.....                                                                               | 5  |
| 2.2.2. Description .....                                                                         | 5  |
| 2.2.3. Manufacture .....                                                                         | 6  |
| 2.2.4. Specifications .....                                                                      | 7  |
| 2.2.5. Genetic Stability.....                                                                    | 8  |
| 2.2.6. Storage Stability .....                                                                   | 8  |
| Part 3: Dietary Exposure .....                                                                   | 10 |
| Part 4: Self-limiting Levels of Use .....                                                        | 11 |
| Part 5: Experience Based on Common Use in Food .....                                             | 12 |
| Part 6: Narrative .....                                                                          | 13 |
| 6.1. Genome Annotation .....                                                                     | 13 |
| 6.1.1. Search Against Antibiotic Resistance Gene Databases .....                                 | 13 |
| 6.1.2. Search Against the Virulence Factor Database .....                                        | 13 |
| 6.1.3. Search of Gene Annotations .....                                                          | 13 |
| 6.1.4. Conclusion of the Genome Safety Assessment.....                                           | 14 |
| 6.2. <i>In Vitro</i> Studies of <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 .....      | 14 |
| 6.2.1. Adhesion .....                                                                            | 14 |

|                                                                                  |    |
|----------------------------------------------------------------------------------|----|
| 6.2.2. Antibiotic Resistance .....                                               | 15 |
| 6.2.3. Production of Biogenic Amines .....                                       | 15 |
| 6.2.4. Production of D-Lactate .....                                             | 15 |
| 6.3. Human Studies of <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19..... | 15 |
| 6.3.1. Studies in Infants.....                                                   | 15 |
| 6.3.2. Studies in Children.....                                                  | 17 |
| 6.3.3. Studies in Adults .....                                                   | 18 |
| 6.5. Studies of Other Strains of <i>L. paracasei</i> ssp. <i>paracasei</i> ..... | 37 |
| 6.6. Toxicity Studies of <i>L. paracasei</i> ssp. <i>paracasei</i> .....         | 48 |
| 6.6. Decision-Tree Analysis .....                                                | 50 |
| 6.7. Evaluations by Authoritative Bodies .....                                   | 50 |
| 6.8. Safety Assessment and GRAS Determination .....                              | 51 |
| 6.8.1. Evidence of Safety .....                                                  | 51 |
| 6.8.2. Conclusion of the Expert Panel.....                                       | 52 |
| Part 7: List of Supporting Data and Information .....                            | 54 |

## List of Tables

|                                                                                                  |    |
|--------------------------------------------------------------------------------------------------|----|
| Table 1. Specifications for <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19.....           | 8  |
| Table 2. Stability: Water Activity. ....                                                         | 9  |
| Table 3. Stability: Cell Count.....                                                              | 9  |
| Table 4. MIC Values for <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19.....               | 15 |
| Table 5. Studies of <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19.....                   | 26 |
| Table 6. Human Studies in <i>L. paracasei</i> ssp. <i>paracasei</i> Strains Other Than F-19..... | 38 |

## List of Figures

|                                                                                         |   |
|-----------------------------------------------------------------------------------------|---|
| Figure 1. <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 Production Process..... | 7 |
|-----------------------------------------------------------------------------------------|---|

## **Part 1: Signed Statements and Certification**

### **1.1. GRAS Notice Submission**

Arla Foods Ingredients (Arla) submits this GRAS notification through its agent James T. Heimbach, president of JHeimbach LLC, in accordance with the requirements of 21 CFR Part 170, Subpart E.

### **1.2. Name and Address of Notifier**

Arla Foods Ingredients Group P/S  
106 Allen Road  
Basking Ridge NJ 07920

#### Notifier Contact

Dr. Kal Ramanujam  
Senior Scientific Advisor  
Sonderhoj 10-12  
8260 DK-Viby J  
Denmark  
[kal.ramanujam@arlafoods.com](mailto:kal.ramanujam@arlafoods.com)  
+1 (484) 919-5759

#### Agent Contact

James T. Heimbach, Ph.D., F.A.C.N.  
President  
JHeimbach LLC  
P.O. Box 66  
Port Royal VA 22535  
[jh@jheimbach.com](mailto:jh@jheimbach.com)  
+1 (804) 742-5543

### **1.3. Name of Notified Organism**

The subject of this Generally Recognized as Safe (GRAS) notification is the probiotic bacterium *L. paracasei* ssp. *paracasei* strain F-19.

### **1.4. Intended Conditions of Use**

*L. paracasei* ssp. *paracasei* strain F-19 is intended to be added to powdered milk-based non-exempt infant formula intended for consumption by healthy term infants at a level consistent with current Good Manufacturing Practice to result in a daily intake of  $10^9$  cfu of the strain.

### **1.5 Statutory Basis for GRAS Status**

Arla's GRAS determination for the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is based on scientific procedures in accordance with 21 CFR §170.30(b).

### **1.6. Premarket Exempt Status**

The intended use of *L. paracasei* ssp. *paracasei* strain F-19 is not subject to the premarket approval requirements of the Federal Food, Drug and Cosmetic Act based on Arla's determination that it is GRAS.

### **1.7. Data Availability**

The data and information that serve as the basis for the conclusion that *L. paracasei* ssp. *paracasei* strain F-19 is GRAS for its intended use will be made available to the FDA upon request. At FDA's option, a complete copy of the information will be sent to FDA in either paper or electronic format, or the information will be available for review at the home office of JHeimbach LLC, located at 923 Water Street, Port Royal VA 22535, during normal business hours.

### **1.8. Freedom of Information Act Statement**

None of the information in the GRAS notice is exempt from disclosure under the Freedom of Information Act, USC 552.

### **1.9. Certification**

To the best of my knowledge, this GRAS notice is a complete, representative, and balanced submission that includes unfavorable information, as well as favorable information, known to me and pertinent to the evaluation of the safety and GRAS status of the intended use of *L. paracasei* ssp. *paracasei* strain F-19.

### **1.10 FSIS Statement**

Not applicable.

### **1.11. Name, Position and Signature of Notifier**



James T. Heimbach, Ph.D., F.A.C.N.  
President  
JHeimbach LLC  
Agent to Arla Foods Ingredients Group P/S

## Part 2: Identity, Methods of Manufacture, Specifications, and Physical and Technical Effect

### 2.1. Name of the GRAS Organism

The notified organism is the probiotic bacterium *L. paracasei* ssp. *paracasei* strain F-19, produced by Chr. Hansen A/S and sold in lyophilized form with a maltodextrin filler under the trade name Probio-Tec F-19 Blend-30 IF; the product contains a minimum of 30 billion ( $3.0 \times 10^{10}$  cfu/g powder) throughout its shelf life.

The strain was deposited in the Belgian Coordinated Collections of Microorganisms, Microbiology Laboratory (BCCM/LMG) under deposit number LMG-P-17806.

### 2.2. Source, Description, Manufacture, and Specifications of the GRAS Organism

#### 2.2.1. Source

The isolation, selection, and characteristics of *Lactobacillus paracasei* ssp. *paracasei* strain F-19 were described in a published paper by Ljungh et al. (2002). At autopsy of hospital patients without gastrointestinal disease, rinsed colons were biopsied and incubated. Over 400 *Lactobacillus* strains were isolated and the 40 strains exhibiting the highest binding of porcine mucin were selected, followed by testing of cell surface hydrophobicity; binding of human fibronectin, collagen, fibrinogen, heparin, and fetuin; and acid tolerance. *L. paracasei* ssp. *paracasei* strain F-19 and 7 other strains with good adhesive and survival ability were chosen for further study, including antibacterial activity and antioxidant activity. The taxonomic placement of *L. paracasei* ssp. *paracasei* strain F-19 was confirmed by ribotyping. (The identification of strain F-19 as *L. paracasei* ssp. *paracasei* was again confirmed by pulsed-field gel electrophoresis [PFGE] and amplified fragment length polymorphism [AFPL] during the EU-PROSAFE project [Vankerckhoven et al. 2008]).

The authors concluded that, “*Lactobacillus F-19* has a documented ability to survive during passage in the GI tract and exerts multiple determinants for colonization and establishment, also expressed during conditions resembling those in the gut. Properties like fiber degradation, immunomodulatory effect and production of antioxidants make *Lactobacillus F-19* a strain which is likely to exert further beneficial effects in the large intestine.”

#### 2.2.2. Description

The taxonomic identification of *L. paracasei* ssp. *paracasei* strain F-19 was confirmed in-house at Chr. Hansen by comparison of its 16S rDNA sequence with that of the type strain of *L. paracasei*.

Cells are non-motile rods with rounded ends, single or in pairs or short chains, non-sporing, Gram+, and catalase negative. They are nonaerobic but aerotolerant.

As noted above, the microorganism is sold in lyophilized form with a food-grade maltodextrin filler. The product also is cryoprotected by food-grade sucrose, maltodextrin, and sodium ascorbate.

##### 2.2.2.1. Sugar Fermentation

The strain is homofermentative and ferments ribose, adonitol, galactose, D-glucose, D-fructose, D-mannose, L-sorbose, mannitol, sorbitol, N-acetyl glucosamine, amygdaline, arbutine, esculine, salicine, cellobiose, maltose, lactose, saccharose, trehalose, inuline, melezitose, D-turanose, D-tagatose, and gluconate. It does not ferment glycerol, erythritol, D-arabinose, L-arabinose, D-xylose, L-xylose,  $\beta$ -methyl-xyloside, rhamnose, dulcitol, inositol,  $\alpha$ -methyl-D-mannoside,  $\alpha$ -methyl-D-glucoside, melibiose, D-raffinose, amidon, glycogen, xylitol,  $\beta$ -gentiobiose, D-lyxose, D-fucose, L-fucose, D-arabitol, L-arabitol, 2-keto-gluconate, or 5-keto-gluconate.

*In vitro* fermentation studies by Hedberg et al. (2008) showed that, in comparison with other tested probiotic bacteria (*L. plantarum* strains 299v and 931, *L. rhamnosus* strains GG and LB21, and *L. reuteri* strain PTA), fermentation by *L. paracasei* ssp. *paracasei* strain F-19 generally proceeds at a significantly slower rate.

#### 2.2.2.2. Genome Analysis

*Lactobacillus paracasei* ssp. *paracasei* strain F-19 was genome sequenced in the in-house genome sequencing facility at Chr. Hansen using published methods with an average sequencing coverage of 62. The assembly yielded seven contigs—i.e., a circular chromosome of 3.1 Mb, a linear plasmid of 93.7 Kb, three circular plasmids of 8.7 Kb, 6.3 Kb, and 2.2 Kb, and two linear fragments of 2.3 Kb and 3.4 Kb.

The genome sequence of *L. paracasei* ssp. *paracasei* strain TMW 1.1434, which is referred to as being isogenic to F-19, was published by Schott et al. (2016; Accession CP016355 and CP016356) and found to be 3.1 Mb, in line with the genome sequence obtained in-house at Chr. Hansen. Furthermore, Schott et al. (2016) published the sequence of a 106.4 Kb plasmid, but not the three small plasmids. The three linear contigs of 2.3 Kb, 3.4 Kb, and 93.7 Kb. were found to be located on the 106.4 Kb plasmid. The presence of three small plasmids was published by Morelli and Campominski (2002), who found them to be 9.0 Kb, 6.5 Kb, and 2.2 Kb as determined by gel electrophoresis.

#### **2.2.3. Manufacture**

Production of *L. paracasei* ssp. *paracasei* strain F-19 takes place under certification that systems are in full compliance with ISO 9001:2008, ISO 22000:2005, ISO/TS 2202-1:2009, and Food Safety System Certification 22000.

Following is a general description of the manufacturing process for *L. paracasei* ssp. *paracasei* strain F-19; a schematic of the process is shown in Figure 1.

##### 2.2.3.1. Production of Substrate

The individual ingredients are weighed and mixed with tap water, and the medium is subjected to ultra-heat treatment (UHT) to remove foreign contamination. The fermentation media employed by Chr. Hansen are primarily based on food-grade milk powder and yeast extract. Various raw materials are used to optimize the fermentation media. All raw materials are regulated processing aids used consistent with their regulatory status, meet food-grade specifications, and are suitable for human consumption.

##### 2.2.3.2. Inoculation and Fermentation

Pre-inoculation material is used to inoculate the milk yeast medium and the medium is incubated. The medium is then transferred to a pre-fermenter containing milk yeast medium. Fermentation in the pre-fermenter is carried out under anaerobic conditions. Then the content is transferred from the pre-fermenter to the fermenter.

The fermentation is conducted with milk yeast medium under anaerobic conditions.

##### 2.2.3.3. Concentration and Treatment

The bacterial cells are harvested and concentrated by centrifugation using a separator.

The harvested bacterial cells are mixed with cryoprotectants and frozen into pellets in liquid nitrogen. The frozen pellets are lyophilized, resulting in very low water activity, ensuring stability of the culture.

##### 2.2.3.4. Standardization

An authorized electronic worksheet is used to calculate the amount of culture that is needed to produce a specific batch. The cell count of the freeze-dried culture forms the basis of the calculation. When the exact weight of the culture is determined, the weight of excipients can be calculated.

#### 2.2.3.5. Milling and Mixing

The lyophilized bacteria concentrate is milled and sieved. Excipients are added to the concentrate in order to standardize the blends and confer the dry powder with desirable handling properties.

The standardized blend is packed into aluminum foil bags of 5 kg each and stored below -20°C, awaiting approval from Quality Control, and then the product is released and available for shipping.




Figure 1. *L. paracasei* ssp. *paracasei* strain F-19 Production Process<sup>1</sup>.

#### 2.2.4. Specifications

Release specifications for *L. paracasei* ssp. *paracasei* strain F-19 are shown in Table 1, along with the results of testing of 5 non-consecutive batches.

<sup>1</sup> UHT = Ultra-High Temperature; PIM = Pre-Inoculation Material; IM = Inoculation Material; PFM = Pre-Fermentation Medium; FM = Fermentation Medium

**Table 1. Specifications for *L. paracasei* ssp. *paracasei* Strain F-19.**

| Parameter                       | Specification                  | Test Batches                   |                                |                                |                                |                                |
|---------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
|                                 |                                |                                |                                |                                |                                |                                |
| Cell count                      | $\geq 3.0 \times 10^{10}$      | $4.9 \times 10^{10}$           | $4.7 \times 10^{10}$           | $4.1 \times 10^{10}$           | $4.5 \times 10^{10}$           | $5.0 \times 10^{10}$           |
| Appearance                      | Fine powder, white-light beige | Fine powder, white-light beige | Fine powder, white-light beige | Fine powder, white-light beige | Fine powder, white-light beige | Fine powder, white-light beige |
| Water Activity                  | $\leq 0.15$                    | 0.05                           | 0.05                           | 0.09                           | 0.09                           | 0.06                           |
| Purity                          |                                |                                |                                |                                |                                |                                |
| <i>Bacillus cereus</i> (cfu*/g) | <100                           | <10                            | <10                            | <10                            | <10                            | <10                            |
| <i>Enterobacteriaceae</i>       | Absent/10x10 g                 | Absent                         | Absent                         | Absent                         | Absent                         | Absent                         |
| <i>Cronobacter sakazakii</i>    | Absent/10x10 g                 | Absent                         | Absent                         | Absent                         | Absent                         | Absent                         |
| <i>Salmonella</i> spp.          | Absent/10x10 g                 | Absent                         | Absent                         | Absent                         | Absent                         | Absent                         |
| <i>S. aureus</i> (cfu/g)        | <10                            | <10                            | <10                            | <10                            | <10                            | <10                            |
| Total aerobic bacteria (cfu/g)  | $\leq 2000$                    | <50                            | <50                            | <50                            | <50                            | <50                            |
| Total yeasts & molds (cfu/g)    | $\leq 100$                     | <5                             | <5                             | <5                             | <25                            | <25                            |
| *cfu = colony-forming units     |                                |                                |                                |                                |                                |                                |

### 2.2.5. Genetic Stability

Morelli and Campominosi (2002) assessed the stability of the plasmid complement of *L. paracasei* ssp. *paracasei* strain F-19 under the stress of the production process. After examining about 300 colony-forming units for their plasmid profile, the experimenters reported that all of them retained the typical plasmid profile of the wild type, suggesting that industrial production processes do not affect the plasmid content of the strain.

In a follow-up experiment, novobiocin was added to the fermentation medium, producing cured derivatives missing one or two of the largest plasmids; it was not found possible to delete the smallest plasmid. Curing did not affect sugar fermentation patterns, indicating that these traits are not plasmid-linked. The cured strains exhibited resistance to vancomycin, again demonstrating that this resistance is intrinsic and is not plasmidally based. Finally, bile resistance was not affected by plasmid curing. The authors concluded:

“The plasmid complement of Lactobacillus F-19 does not seem to be altered by the industrial production processes. Also, the plasmid profile was unaltered when compared with the plasmid content determined in the same strain by our laboratory 6 year earlier. Curing is obtained in laboratory conditions only and the two curable plasmids do not seem to be involved in the checked phenotypes.”

Genetic stability is monitored during storage and production via DNA fingerprinting and plasmid profiling. The results are compared with the reference material for *L. paracasei* ssp. *paracasei* strain F-19.

### 2.2.6. Storage Stability

The stability of batch [REDACTED] *L. paracasei* ssp. *paracasei* strain F-19 was tested when stored at temperatures of  $-20^{\circ}\text{C}$  and  $5^{\circ}\text{C}$  for 2 years and at accelerated conditions of  $25^{\circ}\text{C}$  and 60% relative humidity for 6 months. The product was evaluated for appearance, water activity, and cell

count at 0, 3, 6, 12, 16, and 24 months. The color of the product remained in compliance with specifications at all test points; water activity and cell count are shown in Tables 2 and 3.

**Table 2. Stability: Water Activity.**

| Months               | Water Activity |         |              |
|----------------------|----------------|---------|--------------|
|                      | -20°C±6°C      | 5°C±3°C | 25°C, 60% RH |
| 0                    | 0.08           | 0.08    | 0.08         |
| 3                    | N/A*           | 0.09    | 0.09         |
| 6                    | 0.09           | 0.09    | 0.06         |
| 12                   | 0.07           | 0.08    | N/A          |
| 16                   | 0.09           | N/A     | N/A          |
| 24                   | 0.07           | 0.07    | N/A          |
| *N/A = not available |                |         |              |

**Table 3. Stability: Cell Count.**

| Months               | Cell Count (cfu/g)   |                      |                      |
|----------------------|----------------------|----------------------|----------------------|
|                      | -20°C±6°C            | 5°C±3°C              | 25°C, 60% RH         |
| 0                    | 4.6x10 <sup>10</sup> | 4.6x10 <sup>10</sup> | 4.6x10 <sup>10</sup> |
| 3                    | N/A*                 | 3.7x10 <sup>10</sup> | 3.0x10 <sup>10</sup> |
| 6                    | 5.9x10 <sup>10</sup> | 5.6x10 <sup>10</sup> | 4.8x10 <sup>10</sup> |
| 12                   | 8.2x10 <sup>10</sup> | 7.8x10 <sup>10</sup> | N/A                  |
| 16                   | 4.2x10 <sup>10</sup> | N/A                  | N/A                  |
| 24                   | 3.7x10 <sup>10</sup> | 3.0x10 <sup>10</sup> | N/A                  |
| *N/A = not available |                      |                      |                      |

The data from the stability study show that *L. paracasei* ssp. *paracasei* strain F-19 is stable with respect to color, water activity, and cell count for up to 2 years at temperatures of -20°C or 5°C and for up to 6 months at 25°C.

### Part 3: Dietary Exposure

*L. paracasei* ssp. *paracasei* strain F-19 is intended to be added to powdered milk-based non-exempt infant formula intended for consumption by healthy term infants. The target dietary intake of *L. paracasei* ssp. *paracasei* strain F-19 is  $10^9$  cfu/day. In order to provide  $10^9$  cfu of the probiotic in 800 ml of hydrated formula (an average daily intake), *L. paracasei* ssp. *paracasei* strain F-19 must be present in the powder at a concentration of  $10^7$  cfu/g powder, assuming a hydration rate of 12.5-13.5 g/100 ml ( $10^7$  cfu/g x 0.125 g/ml x 800 ml =  $10^9$  cfu). In order to assure that viable probiotic is present at a concentration of at least  $10^7$  cfu/g powder through its shelf life, it will be introduced at a concentration of  $5 \times 10^7$  cfu/g, leading to a potential daily intake of  $5 \times 10^9$  cfu ( $\pm 20\%$  to allow for production variability).

If the probiotic is added to a formula with a hydration rate different from 12.5-13.5 g/100 ml, the addition concentration will be adjusted as needed to retain the target intake level of  $10^9$  cfu/day.

According to tables of daily energy intake by formula-fed infants provided by Fomon (1993), the subpopulation of infants with the highest intake per kg body weight is boys age 14–27 days. The mean energy intake by this group is 121.1 kcal/kg bw/day and the 90th percentile is 141.3 kcal/kg bw/day. Among girls, the highest energy intake is found in the same age group, 14–27 days, and is nearly as high as boys: the mean and 90th energy intake percentiles are 117.8 and 138.9 kcal/kg bw/day. Most term infant formulas contain 67.6 kcal/100 ml when ready to consume. Therefore, to obtain 141.3 kcal energy/kg bw, an infant boy must consume 209.0 ml formula/kg bw. To reach her 90th percentile of energy consumption, 138.9 kcal/kg bw/day, an infant girl must consume 205.5 ml formula/kg bw/day. The 90th percentile of formula intake for the two sexes combined is about 207 ml/kg bw/day.

The 90<sup>th</sup> percentile exposure is then calculated as  $5 \times 10^7$  cfu/g x 0.125 g/ml x 207 ml/kg bw/day =  $1.3 \times 10^9$  cfu/kg bw/day. Thus, the EDI for *L. paracasei* ssp. *paracasei* strain F-19 added to powdered formula at a level of  $5 \times 10^7$  cfu/g, i.e., the 90<sup>th</sup> percentile of intake, is  $1.3 \times 10^9$  cfu/kg bw/day  $\pm 20\%$ .

The target population is healthy infants and toddlers aged 0-3 years of age. Since it is not expected that these infants and toddlers will have other dietary sources of *L. paracasei* ssp. *paracasei* strain F-19, this figure represents the total EDI from all sources.

#### **Part 4: Self-limiting Levels of Use**

There is no technological or organoleptic limitation to the concentration of *L. paracasei* ssp. *paracasei* strain F-19 in foods.

## **Part 5: Experience Based on Common Use in Food**

The conclusion that the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is GRAS is based on scientific procedures rather than experience based on common use in food prior to 1958. Since 2001, the strain has been present in consumer products such as Gaio Dofilus, Gaio Yoghurtdryck, and in products of the Cultura brand in Denmark and Sweden, in porridge and baby food by Semper in Sweden, and in follow-on formulas in Sweden and the Dominican Republic. No adverse events associated with *L. paracasei* ssp. *paracasei* strain F-19 have been reported.

## Part 6: Narrative

### 6.1. Genome Annotation

The genome sequence of *L. paracasei* ssp. *paracasei* strain F-19 obtained in-house at Chr. Hansen was used for the genome safety assessment (Chr. Hansen 2018). For the assessment, the genome sequence of *L. paracasei* ssp. *paracasei* strain F-19 was subjected to annotation using published methods. The F-19 genome contained 3,080 coding sequences (CDS) and 74 RNAs with 2,963 CDS and 74 RNAs on the chromosome, 110 CDS on the 106 Kb plasmid, 7 CDS on the 8.7 Kb plasmid, 5 CDS on the 6.3 Kb plasmid, and 2 CDS on the 2.2 Kb plasmid. The number of CDS on the published sequence is slightly lower than the in-house data.

#### 6.1.1. Search Against Antibiotic Resistance Gene Databases

To identify genes with high homology to previously published antibiotic resistance genes, the genome of *L. paracasei* ssp. *paracasei* strain F-19 was analyzed against different published databases of antibiotic resistance genes. The databases focused either on resistance genes of relevance to Gram-positive bacteria and in particular to lactic acid bacteria and bifidobacteria or antibiotic resistance genes identified primarily in pathogenic species as they are most abundantly characterized. The genome of strain F-19 was analyzed and no antibiotic resistance genes were detected.

#### 6.1.2. Search Against the Virulence Factor Database

The genome of *L. paracasei* ssp. *paracasei* strain F-19 was analyzed against a published database of virulence factors containing virulence factors from 30 different pathogens, including Gram positive pathogens such as *Enterococcus*, *Staphylococcus*, *Streptococcus*, and *Listeria*.

Most of the hits were associated with Clp and heat shock proteins, surface structures, and transport or secretion systems. None of the hits were assessed to be virulence factors and all hits could be regarded as ‘niche factors’ (Hill 2012) since they are also found in commensal bacteria.

Two hits were associated with genes that could be linked to adhesion to extracellular matrix or host cell surfaces. Homologs of these genes were observed in many *L. paracasei* strains and, in case they should have a role in adhesion, this could be regarded as a probiotic feature in *L. paracasei* ssp. *paracasei* strain F-19 rather than a safety issue.

One hit had low homology to a gene annotated as “Peptide methionine sulfoxide reductase MsrB (EC 1.8.4.12).” *In vivo*, MsrS proteins are essential for the protection of cells against oxidative damage to proteins and thereby have a role in the virulence of some pathogens; however, a MsrB mutant in *L. reuteri* also shows reduced performance in the murine gut. Thus, MsrB could be regarded as a niche factor important for the probiotic feature of the strain and, since protection against oxidative stress is important for all bacteria, is not a safety issue.

One hit had low homology to a gene annotated as “Hyaluronate lyase precursor (EC 4.2.2.1).” Hyaluronate lyases are reported to play important roles in pathogenesis by providing nutrients to the pathogen to enhance spreading of the pathogen by degrading hyaluronic acid. However, in other pathogens, such as Group A *Streptococci*, hyaluronate lyases are suggested to be anti-virulence factors, as the GAS-capsule mainly consists of hyaluronic acid, suggesting that hyaluronate lyase is not a critical virulence factor. A homolog of the gene is observed in a wide range of *L. casei* and *L. paracasei* strains, supporting the conclusion that it is not a safety issue in *L. paracasei* ssp. *paracasei* strain F-19.

#### 6.1.3. Search of Gene Annotations

The genome sequence of *L. paracasei* ssp. *paracasei* strain F-19 was subjected to annotation using published methods. The gene annotations were searched to identify terms that could be linked to *Lactobacillus paracasei* ssp. *paracasei* strain F-19

antibiotic resistance. A total of 61 genes included one or more of these words in its annotation. Many proved to be housekeeping genes or transporters, 18 annotated as “resistance” were associated with heavy metals, and the remainder were annotated with names of antibiotics to which the strain shows no resistance and homologs were observed in many *L. paracasei* strains. All were dismissed as safety concerns.

In a similar fashion, the gene annotations were searched for terms that could be linked to virulence. This search identified 29 genes; 18 were determined to be housekeeping genes of no safety concern. Those containing the word “toxin” in the annotation were found to be bacterial toxin-antitoxin systems, part of the cellular regulatory machinery. Three genes were transporters of unknown specificity, two were adhesion factors, and one gene identified as a hemolysin appeared rather to encode a membrane protein; in any case, the strain is not hemolytic.

Finally, because biogenic amines are produced by decarboxylation of amino acids, the gene annotations were searched for the word “decarboxylase.” Only one such gene was found, labeled “ornithine decarboxylase,” but it is not accompanied by a transporter required for putrescine, nor does *L. paracasei* ssp. *paracasei* strain F-19 show evidence of biogenic amine production when tested phenotypically.

#### **6.1.4. Conclusion of the Genome Safety Assessment**

The genome of *L. paracasei* ssp. *paracasei* strain F-19 was analyzed for the presence of antibiotic resistance genes and no antibiotic resistance genes were detected. No virulence genes were identified in the genome of *L. paracasei* ssp. *paracasei* strain F-19, indicating a very low virulence potential.

In conclusion, there are no indications that *L. paracasei* ssp. *paracasei* strain F-19 is a safety concern as based on the performed genome safety assessment.

## **6.2. In Vitro Studies of *L. paracasei* ssp. *paracasei* Strain F-19**

### **6.2.1. Adhesion**

Human mucus was isolated from fecal samples of newborns (n = 28), 2-month-old (n = 11) and 6-month-old (n = 17) infants, and adults (n = 14) to compare mucus binding of 7 strains of probiotic bacteria (Kirjavainen et al. 1998). As compared with *B. animalis* ssp. *lactis* strain Bb-12, *L. crispatus* strain Mu5, *L. crispatus* strain M247, *L. rhamnosus* strain GG, *L. johnsonii* strain LJ-1, and *L. salivarius* strain LM2-118, *L. paracasei* ssp. *paracasei* strain F-19 was of intermediate adhesion ability for all age groups. In newborns, it exhibited greater adhesion than 3 strains and inferior adhesion to 3 other strains; in the other 3 age groups it adhered better than 2 strains and less well than 4 strains. The authors suggested that “the age of the target group may be worthy of consideration when planning a a schedule for probiotic or functional food therapy.”.

Juntunen et al. (2001) studied adhesion to human intestinal mucus *in vitro* using mucus derived from fecal samples from 20 infants during and after rotavirus diarrhea and 10 healthy age-matched infants. Both monoculture inocula of *L. casei* strain Shirota, *L. paracasei* F-19, *L. rhamnosus* GG, *L. acidophilus* LA5, and *B. animalis* ssp. *lactis* Bb12 and combinations of these bacteria were tested. Adherence was highest for *L. rhamnosus* GG (34%) and *B. animalis* ssp. *lactis* Bb12 (31%) and lowest for *L. casei* strain Shirota (1%); *L. paracasei* F-19 was second lowest with 3% adhesion. Some probiotic combinations appeared to result in synergistically enhanced adherence. There were no significant differences in adhesion between the samples from infants with rotavirus diarrhea, infants after rotavirus diarrhea, or healthy infants.

### 6.2.2. Antibiotic Resistance

Minimum inhibitory concentrations (MIC) of 9 antibiotics were determined for *L. paracasei* ssp. *paracasei* strain F-19 according to the ISO 10932 | IDF 223 international standard. These MIC, shown in Table 4, were compared with the cut-off values established for the *Lactobacillus casei/paracasei* group by the European Food Safety Authority (EFSA; FEEDAP 2012).

**Table 4. MIC Values for *L. paracasei* ssp. *paracasei* Strain F-19.**

| Antibiotic Type                  | Antibiotic      | MIC (µg/ml) | EFSA Cut-Off Value (µg/ml) |
|----------------------------------|-----------------|-------------|----------------------------|
| Aminoglycoside                   | Gentamicin      | 1-2         | 32                         |
|                                  | Kanamycin       | 32          | 64                         |
|                                  | Streptomycin    | 16          | 64                         |
| Tetracycline                     | Tetracycline    | 2           | 4                          |
| Macrolide                        | Erythromycin    | 0.12-0.25   | 1                          |
| Lincosamide                      | Clindamycin     | 0.25        | 1                          |
| Chloramphenicol                  | Chloramphenicol | 4-8         | 4                          |
| β-lactam                         | Ampicillin      | 1           | 4                          |
| Glycopeptide                     | Vancomycin      | >128        | n.r.                       |
| n.r. = not required to be tested |                 |             |                            |

*L. paracasei* ssp. *paracasei* strain F-19 is sensitive to most of the antibiotics tested with MIC values that are below the EFSA 2012 cut-off values for *Lactobacillus casei/paracasei* group. Although the MIC value for chloramphenicol is one two-fold dilution above the EFSA cut-off value in some replicates, that is considered acceptable due to the technical variation of the phenotypic method.

### 6.2.3. Production of Biogenic Amines

*L. paracasei* ssp. *paracasei* strain F-19 was analyzed for production of histamine, tyramine, cadaverine, and putrescine using an in-house procedure based on published methods; no production of the four biogenic amines was detected (Chr. Hansen, 2018).

### 6.2.4. Production of D-Lactate

*L. paracasei* ssp. *paracasei* strain F-19 was analyzed for production of L- and D-lactate; the ratio between L- and D-lactic acid was detected and it was determined that over 95% of the lactate produced was the L-enantiomer (Chr. Hansen, 2018).

## 6.3. Human Studies of *L. paracasei* ssp. *paracasei* Strain F-19

The studies discussed in this section are summarized in Table 5 at the end of the section.

### 6.3.1. Studies in Infants

West and her colleagues (West et al. 2008; West et al. 2009) reported on a clinical trial of *L. paracasei* ssp. *paracasei* strain F-19 (referred to as LF-19) with follow-ups when the infants reached the age of 8-9 years (West et al. 2013, Chorell et al. 2013, Hasslof et al. 2013, Videhult et al. 2015a, Videhult et al. 2015b).

In a prospective, randomized, double-blind, placebo-controlled study reported by West et al. (2008), 179 healthy 4-month-old term infants (75 boys and 104 girls) were enrolled and assigned to receive cereal supplemented with 0 (n = 90) or 10<sup>8</sup> cfu (n = 89) of *L. paracasei* ssp. *paracasei* strain F-19 daily to age 13 months. All infants were immunized with DTaP (diphtheria and tetanus

toxoid and acellular pertussis), polio, and Hib-conjugate vaccines at 3, 5½, and 12 months of age. Parents recorded feedings, stooling, and any symptoms. Venous blood was collected at 5½, 6½, 12, and 13 months of age for analysis of IgG antibodies to tetanus toxoid and diphtheria toxin. Fecal samples were collected at enrolment and at 6½, 9, and 13 months of age for bacterial testing.

Five infants from the test group and 3 from the controls were withdrawn by their parents for reasons not associated with the intervention. *L. paracasei* ssp. *paracasei* strain F-19 was detected in feces of 90% of the infants receiving the probiotic. The number of days with fever, respiratory illness, or diarrhea did not differ between the groups, but days receiving antibiotics were significantly fewer among infants ingesting strain F-19. Exposure to *L. paracasei* ssp. *paracasei* strain F-19 enhanced anti-diphtheria concentrations. No adverse effects of the probiotic were reported, and the authors concluded that, “feeding LF-19 [West et al.’s term for *L. paracasei* ssp. *paracasei* strain F-19] did not prevent infections, but increased the capacity to raise immune responses to protein antigens.”

In additional analyses of the data from this study, West et al. (2009) found that the ratio of IFN-γ to IL4 mRNA was significantly greater in infants receiving *L. paracasei* ssp. *paracasei* strain F-19 and the incidence of eczema was significantly reduced, suggesting “enhancing effects of *Lactobacillus* F-19 on the T-cell mediated immune response.” In further analyses, West et al. (2012a), reported little difference between test and control groups in expression levels of IL-2, IFN-γ, IL-4, IL-17A, and IL-10 messenger RNAs, suggesting “modest effects by probiotics on T-cell maturation.”

West et al. (2013) conducted a follow-up with 121 of the infants from the trial described previously (West et al. 2008) at 8-9 years of age to assess the prevalence of allergic disease (eczema, allergic rhinitis, asthma, and food allergy) by clinical examination. The available participants included 59 from the probiotic group and 62 from the treatment group. Neither benefits nor adverse effects were reported. The authors concluded that, “There was no long-term effect of LF-19 on any diagnosed allergic disease, airway inflammation or IgE sensitization.”

In yet another report on data from the West et al. (2008) study, Chorell et al. (2013) reported that there were no effects on anthropometrics of serum lipids due to ingestion of F-19, but there was a significant increase in putrescine, which the authors suggested might contribute to increased gut integrity.

Hasslof et al. (2013) published still another report on the later experience of infants from the West et al. (2008) study—this time their dental health at 9 years of age. The authors explained that the rationale for the study was that “It has been suggested that the chronic use of probiotics . . . in young children may lead to undesirable side effects in the intestine and the oral cavity.” No differences were seen between those who had received *L. paracasei* ssp. *paracasei* strain F-19 and those who had not, either beneficial or adverse, and the authors concluded that, “early intervention with LF-19 [*L. paracasei* ssp. *paracasei* strain F-19] did not affect the frequency of dental caries, [mutans streptococci], or [lactobacilli]. LF-19 did not establish itself as a permanent facet of the oral microbiota in any of the subjects included in this study.”

Videhult et al. (2015a) reassessed the effects of *L. paracasei* ssp. *paracasei* strain F-19 on body composition, growth, and metabolic markers of 8-9-year-old children given the probiotic as infants, as reported in West et al. (2008). Of 179 children included in the original study, 120 were available for the follow-up at 8–9 years of age, 58 in the probiotic group and 62 in the placebo group. Anthropometrics, body composition, serum lipids, insulin, glucose, and transaminases were measured. There were no significant differences between groups on any measures, and the authors reported that, “Feeding LF-19 during infancy did not modulate body composition, growth or any of

the assessed metabolic markers at school age.” They concluded that “we observed no adverse effect of LF-19 on body composition or metabolism in this follow-up study.”

Videhult et al. (2015b) assessed the effects of *L. paracasei* ssp. *paracasei* strain F-19 given to infants, as reported in West et al. (2008), on the metabolic and inflammatory profile of 120 of the infants as 8-9-year-old children. Overweight or obese children had increased plasma C-peptide, plasminogen activator inhibitor-1, leptin, and serum high-sensitivity C-reactive protein compared with normal weight children, but there was no association with *L. paracasei* ssp. *paracasei* strain F-19. The authors concluded that “the probiotic LF-19 had no long-term effect on the inflammatory and metabolic profile.”

Zampieri et al. (2013) reported a prospective, randomized, unblinded trial enrolling 32 preterm infants (all delivered before gestational week 32; birth weight 600-1500 g) at Bell’s stage 2 necrotizing enterocolitis (NEC). Eighteen patients were randomly assigned to receive  $6 \times 10^9$  cfu/day of *L. paracasei* ssp. *paracasei* strain F-19 along with standard medical treatment, while 14 patients received only standard treatment. During the study, records were kept of ventilator support need and need for nasogastric feeding. Infants receiving *L. paracasei* ssp. *paracasei* strain F-19 had a significantly reduced probability of progression to Bell’s stage 3 NEC; the authors suggested that improved intestinal motility might have contributed to this result. The authors reported that, “No collateral effects were observed during the study period; none of our patients presented sepsis due to *L. paracasei* subsp. *paracasei* strain F-19. No patient required preventive exclusion from the trial. No patient [in the probiotic group] presented treatment-related intestinal complications (i.e., diarrhea).” Their conclusion was that “The use of *L. paracasei* subsp. *paracasei* strain F-19 is safe and effective.”

Growth of infants receiving formula supplemented with prebiotics or synbiotics was studied in a prospective, randomized, double-blind, two-arm, multicenter study (Szajewska et al. 2017). A total of 182 healthy term infants, 93 males and 89 females, aged 28 days were randomized to receive formula supplemented with fructooligosaccharides (FOS) and galactooligosaccharides (GOS) or with FOS, GOS, and  $10^9$  cfu of *L. paracasei* ssp. *paracasei* strain F-19 to 6 months of age. Study visits occurred every 28 days to age 4 months, then at 6, 9, and 12 months, for anthropometric measures, assessment of tolerance and health-related outcomes, and description of stool characteristics. The primary outcome measure was growth (body weight, length, and head circumference) during the first year of life. The secondary outcome measures were health-related parameters.

There was no difference in growth between the two groups, but the infants receiving synbiotics exhibited a weak but significant reduction of lower respiratory tract infections. With regard to safety, the authors reported that, “Probiotic supplementation of infant formula with *Lactobacillus* F-19 was well tolerated, and no significant differences between the experimental and control groups were observed in regard to adverse events,” and concluded:

“The lack of a significant difference between the formula-fed groups for growth and for the occurrence of serious adverse events supports the safety of using *Lactobacillus* F-19-supplemented synbiotic formula in healthy term infants.”

### 6.3.2. Studies in Children

Sixty-one apparently healthy and fully weaned children (sex not reported; mean age =  $13.1 \pm 5.3$  years) were enrolled in a prospective, randomized, double-blind, placebo-controlled study (Sullivan et al. 2002) of the effect of *L. paracasei* ssp. *paracasei* strain F-19 on their colonic microbiota. Both the test group (n = 30) and the placebo group (n = 31) consumed gelatin capsules containing 0 or  $10^{10}$  cfu of *L. paracasei* ssp. *paracasei* for 3 weeks. Children’s parents kept a daily record of

stooling characteristics and gastrointestinal symptoms. Fecal samples were collected at baseline and at weeks 2 and 5 for bacterial analysis.

Children receiving *L. paracasei* ssp. *paracasei* strain F-19 showed a significant increase in fecal levels of the strain, but not of other lactobacilli or bifidobacteria. The authors reported that, “The products were generally well tolerated . . . Fecal characteristics and occurrences of gastrointestinal symptoms were similar in the treatment and placebo groups.”

As part of a multicenter project named PROBDEMO (Crittenden et al. 2002) 61 apparently healthy male and female children aged 1-1.5 years enrolled in a prospective, randomized, double-blind, placebo-controlled study received gelatin capsules providing 0 or  $2 \times 10^{10}$  cfu of *L. paracasei* ssp. *paracasei* strain F-19 for 3 weeks. Fecal samples were taken for assessment of F-19 and other bacteria. Ingestion of the F-19 strain resulted in its presence in fecal samples, but did not significantly alter the balance of the major population groups within the intestinal microbiota. The study design included observations monitoring potential side-effects of probiotic consumption, including intestinal discomfort, increased flatulence, and changes in stool consistency and frequency. No such adverse effects were observed and the probiotic was well tolerated by all individuals. The authors concluded that, “The fact that no side-effects were detected in subjects . . . suggests that *Lactobacillus* F-19 is a safe microbial food supplement.”

### 6.3.3. Studies in Adults

Sullivan et al. (2001) reported a prospective, randomized, double-blind, placebo-controlled trial of the effects of treating elderly *Helicobacter pylori* patients with *L. paracasei* ssp. *paracasei* strain F-19 and inulin. Thirty-five patients (sex not reported; age range 58-89 years with a median age of 76 years) were randomized into treatment (n = 17) and placebo (n = 18) groups; the groups received cultured buttermilk with 0 or  $10^{11}$  cfu of F-19 and 0 or 6 g inulin for 12 weeks. Fecal samples were collected at baseline and at weeks 4, 12, and 20 for bacterial analysis. The numbers of *L. paracasei* ssp. *paracasei* strain F-19 and other lactobacilli increased significantly in the test group. The authors concluded that the probiotic “survives the passage through the gastrointestinal tract.” There was no discussion of any adverse events.

Thirty elderly *Helicobacter pylori* patients (sex not reported; mean age =  $76.8 \pm 6.5$  years) were enrolled in a prospective, randomized, double-blind, placebo-controlled study (Sullivan et al. 2002) of the effect of *L. paracasei* ssp. *paracasei* strain F-19 on their colonic microbiota. Both the test group (n = 13) and the placebo group (n = 17) consumed fermented milk twice daily for 12 weeks; the test group’s milk provided  $10^{11}$  cfu of *L. paracasei* ssp. *paracasei* daily. Patients kept a daily record of stooling characteristics and gastrointestinal symptoms, serum samples were examined for anti *H. pylori* IgG, and breath was analyzed for urea. Fecal samples were collected at baseline and at weeks 4, 12, and 20 for bacterial analysis.

Patients receiving *L. paracasei* ssp. *paracasei* strain F-19 showed a significant increase in fecal levels of the strain and other lactobacilli, but no effect was reported on *H. pylori* infection. The authors reported that, “The products were generally well tolerated . . . Fecal characteristics and occurrences of gastrointestinal symptoms were similar in the treatment and placebo groups.”

As part of a multicenter project named PROBDEMO (Crittenden et al. 2002), 5 apparently healthy adults (sex not reported) consumed fermented milk providing  $4 \times 10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus*, and *B. longum* per day for 12 days in an open-label study. Fecal samples were taken at baseline and at 12 days, and biopsy samples were taken from the colonic mucosa after 12 days, for enumeration of F-19 and other lactobacilli. In the fecal samples, lactobacilli were present at a level of  $2.0 \pm 1.3 \times 10^6$  cfu/g both before and after treatment, with the F-19 strain absent prior to treatment but constituting 65% of total lactobacilli after treatment. In the colonic mucosal biopsy, strain F-19 constituted 10% of the  $0.9 \pm 1.7 \times 10^5$  cfu/g lactobacilli present.

Similar levels of *L. paracasei* ssp. *paracasei* F-19 were found at all locations within the colon, ascending, transverse, and descending; “demonstrating,” according to the authors, “that it can colonize the length of the colon.”

In another study within the PROBDEMO project, 9 apparently healthy adults (sex not reported), 4 with milk hypersensitivity, consumed non-fermented milk providing 0 or  $4 \times 10^8$  cfu of *L. paracasei* ssp. *paracasei* strain F-19 for one week in a prospective, single-blind, placebo-controlled, crossover study (Crittenden et al. 2002). Fecal samples were taken for assessment of F-19 and other bacteria. As in the first study, the F-19 strain was found in fecal samples after a week of ingestion but did not significantly disturb the balance of the major population groups within the intestinal microbiota in individuals with or without milk hypersensitivity.

In another prospective, randomized, double-blind, placebo-controlled study within the PROBDEMO project, 30 elderly patients (>65 years) seropositive to *H. pylori* consumed fermented milk providing 0 or  $1.5 \times 10^{11}$  cfu of *L. paracasei* ssp. *paracasei* strain F-19 daily for 12 weeks. Fecal samples were taken for assessment of F-19 and other bacteria. Consumption of *L. paracasei* ssp. *paracasei* strain F-19 did not significantly disturb the balance of the major population groups within the intestinal microbiota in *H. pylori*-infected patients.

All PROBDEMO trials included observations monitoring potential side-effects of probiotic consumption (Crittenden et al. 2002). These included intestinal discomfort, increased flatulence, and changes in stool consistency and frequency. The trials included healthy subjects, adults with verified milk-hypersensitivity, and elderly people infected with *H. pylori*. The authors concluded that, “No adverse effects of probiotic administration were observed in any of the pilot studies. The probiotic was well tolerated by all individuals, including the elderly *H. pylori* patients who consumed *L. paracasei* ssp. *paracasei* strain F-19 daily for 3 months without adverse effects. The fact that no side-effects were detected in subjects ranging in age from 1 to 85 years, and in healthy individuals and subjects suffering mild illnesses (milk-hypersensitivity and *H. pylori* infection) suggests that *Lactobacillus* F-19 is a safe microbial food supplement.”

Rayes and her colleagues published a series of similar studies (2002a, 2002b, 2002c, 2005, 2007, 2012) investigating the ability of *L. paracasei* ssp. *paracasei* strain F-19 to reduce the incidence of post-operative bacterial infection in patients recovering from major abdominal surgery or organ transplantation.

In a prospective, randomized, double-blind, placebo-controlled trial of patients receiving liver transplants, Rayes et al. (2002a) divided 95 patients (49 males, 46 females with mean age =  $49.0 \pm 2.5$  years) into 3 groups—one group received standard enteral nutrition formula, a second group received the formula with added fiber and  $2 \times 10^9$  cfu of *L. paracasei* ssp. *paracasei* strain F-19, and a third group received the formula with added fiber plus heat-killed bacteria. All infections and isolated bacteria were recorded as well as days in the ICU and the hospital and any side effects. Non-infectious complications such as biliary fistulas, anastomotic leaks, and impaired kidney function were monitored, and blood was taken on post-operative days 1, 5, and 10 for hematology, clinical chemistries, and immunology. Body temperature was measured three times daily and surveillance cultures from urine, blood, bile, and intra-abdominal drainages were done twice a week.

The patients who received living lactobacilli plus fiber developed significantly fewer bacterial infections (13%) than the patients on standard formula (48%) or inactivated bacteria and fiber (34%). Both standard enteral formula and strain F-19 were well tolerated. Abdominal side effects (distension, cramps, or diarrhea) were seen in 8 of 32 patients in the standard-formula group, 6 of 31 patients in the live *Lactobacillus* group, and 11 of 32 patients in the inactivated-bacteria group.

Rayes et al. (2002b, abstract) enrolled 172 patients following major abdominal surgery or liver transplantation in a prospective, randomized, double-blind, placebo-controlled trial in which they received either conventional enteral nutrition, enteral nutrition with fiber and  $2 \times 10^9$  cfu of *L. paracasei* ssp. *paracasei* strain F-19, or enteral nutrition with fiber and heat inactivated lactobacilli. The groups receiving live *Lactobacillus* cultures experienced significantly reduced incidences of bacterial infection (among resection patients, 4% vs. 13% in the prebiotic group and 31% in the conventional-therapy group; among transplant recipients, 13% in the probiotic group vs. 34% in the prebiotic group and 48% in the conventional therapy group). The authors reported that “Fibre and lactobacilli were well tolerated in most cases.”

Ninety patients (48 males and 42 females, mean age =  $61.0 \pm 14.1$  years) recovering from major abdominal surgery were enrolled in a prospective, randomized, double-blind, placebo-controlled trial (Rayes et al. 2002c). Three randomly assigned groups of  $n = 30$  received conventional enteral nutrition, enteral nutrition with fiber and  $2 \times 10^9$  cfu of *L. paracasei* ssp. *paracasei* strain F-19, or enteral nutrition with fiber and heat inactivated bacteria. All infections and isolated bacteria were recorded as well as days in the ICU and the hospital and any side effects. Non-infectious complications such as biliary fistulas, anastomotic leaks, and impaired kidney function were monitored, and blood was taken on post-operative days 1, 5, and 10 for hematology, clinical chemistries, and immunology. Body temperature was measured three times daily and surveillance cultures from urine, blood, bile, and intra-abdominal drainages were done regularly.

All patients completed the study, receiving the full course of their assigned nutrition. The incidence of infections was significantly lower in groups receiving fiber and either live or killed lactobacilli (10% each) than in the group receiving standard nutritional formula (30%). Patients receiving live F-19 strain needed antibiotics for a significantly shorter time than did the patients in the other groups. The length of hospital stay and the incidence of non-infectious complications did not differ significantly. Fibers and lactobacilli were well tolerated; side effects did not differ between groups and there was no diarrhea. The authors concluded that “application of fiber and lactobacilli is feasible even after major abdominal surgery and caused few side effects.”

In a prospective, randomized, double-blind, placebo-controlled trial (Rayes et al. 2005), 66 liver transplant recipients (38 males, 28 females, mean age =  $51.5 \pm 2.5$  years), 33 per group, received enteral nutrition with prebiotics and with or without probiotics for 14 days beginning the day prior to surgery. The daily oral probiotic administration included  $2 \times 10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *Pediococcus pentosaceus* strain 5-33:3, *Leuconostoc mesenteroides* strain 77:1, and *L. plantarum* strain 2362. The primary endpoint was the occurrence of post-operative bacterial infection. All infections and isolated bacteria were recorded as well as days in the ICU and the hospital and any side effects. Non-infectious complications such as biliary fistulas, anastomotic leaks, and impaired kidney function were monitored, and blood was taken on post-operative days 1, 4, and 8 for hematology, clinical chemistries, and immunology. Body temperature was measured twice daily and surveillance cultures from urine, blood, bile, and intra-abdominal drainages were done three times a week.

Inclusion of probiotics in the enteral nutrition formula significantly reduced the incidence of post-operative bacterial infection and reduced the severity of infections that occurred. Hematology, immunology, and biochemistry parameters did not differ significantly between groups. The authors reported that “Enteral nutrition, containing the synbiotic combination, was well tolerated in all patients,” with fewer probiotic patients exhibiting diarrhea or abdominal cramps. “All side effects disappeared under temporary reduction in the amount of enteral nutrition.” The authors concluded that inclusion of probiotics “is an effective means to prevent post-operative bacterial infections in high-risk surgical patients, and since it causes no resistant strains and has no serious side effects, it could be widely used.”

Rayes et al. (2007) reported a fifth prospective, randomized, double-blind, placebo-controlled trial of the effect of synbiotics on the incidence of post-surgical bacterial infections. Eighty patients (45 males, 35 females, mean age =  $58.5 \pm 12.5$  years) recovering from pylorus-preserving pancreatoduodenectomy received enteral nutrition with prebiotics and with or without probiotics for 14 days beginning the day prior to surgery. The daily oral probiotic administration included  $2 \times 10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *Pediococcus pentosaceus* strain 5-33:3, *Leuconostoc mesenteroides* strain 77:1, and *L. plantarum* strain 2362. The primary endpoint was the occurrence of post-operative bacterial infection. All infections and isolated bacteria were recorded as well as days in the ICU and the hospital and any side effects. Non-infectious complications such as biliary fistulas, anastomotic leaks, and impaired kidney function were monitored, and blood was taken on post-operative days 1, 4, and 8 for hematology, clinical chemistries, and C-reactive protein. Body temperature was measured twice daily and surveillance cultures from urine, blood, bile, and intra-abdominal drainages were done in case of suspected infection.

Inclusion of probiotics in the enteral nutrition formula significantly reduced the incidence and duration of post-operative bacterial infection to 12.5% vs. 40% in the fiber-only formula group. The incidences of diarrhea and abdominal cramps were the same in both groups of patients. The authors reported that, “Enteral nutrition, containing the synbiotic combination, was well tolerated in all patients... All side effects disappeared under temporary reduction in the amount of enteral nutrition,” and concluded, “Early enteral nutrition with synbiotics was able to significantly reduce postoperative bacterial infections in patients following [pancreatoduodenectomy] with only single-shot antibiotic prophylaxis. In contrast to antibiotics, it is relatively cheap and does not cause resistant strains or serious side effects.”

The same combination of probiotics ( $2 \times 10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *Pediococcus pentosaceus* strain 5-33:3, *Leuconostoc mesenteroides* strain 77:1, and *L. plantarum* strain 2362) was tested for its effect on liver function after hepatic resection (Rayes et al. 2012). Nineteen right-hepatectomy patients (14 males and 5 females, mean age =  $60.1 \pm 13.9$  years) were enrolled in a prospective, randomized, double-blind, placebo-controlled pilot study in which they received enteral nutrition formula containing prebiotic or containing both pre- and probiotic for 10 days post-operatively. The primary study end point was restoration of liver function, tracked via laboratory parameters. The incidence of surgical complications, bacterial infections, and any side effects was monitored. Body temperature was measured twice daily and blood was drawn for tests of hematology, clinical chemistries, and C-reactive protein.

Results of liver function tests were ambiguous, and the authors suggested that patient numbers were too small and the clinical courses too heterogeneous to draw conclusions regarding the effectiveness of synbiotic intervention in restoration of liver function. The authors were able to conclude that, “The synbiotic combination was well-tolerated in all patients. Mild side-effects (abdominal distension and cramps) occurred in three patients of each group but disappeared under symptomatic therapy. No severe side effects were recorded.”

A prospective, randomized, double-blind, placebo-controlled study (Sullivan et al. 2003) was designed to test the ability of a probiotic mixture of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain NCFB 1748, and *B. animalis* ssp. *lactis* strain Bb12 to prevent antibiotic-associated ecological disturbances of the intestinal microbiota. Twenty-four apparently healthy adults (7 men and 17 women aged 21-48 years) ingested 4 clindamycin capsules daily for 7 days; 12 of them also consumed a yogurt product providing  $2 \times 10^8$  cfu of each strain while the other 12 consumed plain yogurt for 14 days. Stool samples were collected at baseline and on days 2, 5, 7, 10, 14, and 21 for microbiological analysis.

Consumption of the probiotic prevented ecological disturbances in the numbers of lactobacilli and *Bacteroides fragilis* group. One participant in the probiotic group developed diarrhea and one reported looser stools. The first person's stool was cytotoxin positive and was cured by metronidazole; the loose stools resolved spontaneously. No other adverse events were reported.

Riordan et al. (2003) reported an open label study in which 11 cirrhosis patients (age and sex not reported) and 5 healthy controls consumed sachets providing beta glucan, inulin, pectin, resistant starch, and  $8 \times 10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *L. plantarum* strain 2362, *Pediococcus pentosaceus* strain 5-33:3, and *Lactococcus raffinolactis* strain 32-77:1 for 7 days. Toll-like receptor (TLR) 2 expression, serum TNF- $\alpha$  levels, and production of TNF- $\alpha$  were measured at baseline and at 7 and 28 days after cessation of supplementation. Supplementation with the synbiotic regimen resulted in significant up-regulation of peripheral blood mononuclear cell expression of TLR2. Serum TNF- $\alpha$  levels were further increased while TNF- $\alpha$  production was reduced in most patients. The authors reported that, "Administration of the synbiotic supplement was well tolerated without any reported adverse events or change in general clinical state."

Liu et al. (2004) reported on a prospective, randomized, double-blind, placebo-controlled trial in which 55 patients (53 M, 2 F) aged  $56 \pm 11$  years with minimal hepatic encephalopathy were randomized to receive a synbiotic preparation containing fermentable fiber and  $10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *Pediococcus pentosaceus*, *Leuconostoc mesenteroides*, and *L. plantarum* (n = 20), fermentable fiber alone (n = 20), or a non-fermentable placebo (n = 15) for 30 days. Patients were re-assessed for minimal hepatic encephalopathy on day 30, while fecal samples were taken at baseline, on day 30, and 14 days later for bacterial analysis.

Either the synbiotic or fiber-alone produced significant improvement in minimal hepatic encephalopathy and reduction in potentially pathogenic *E. coli* and *Staphylococcal* species. The authors reported that, "The synbiotic, fermentable fiber and placebo preparations were well-tolerated by all patients, with no reports of adverse side effects. In particular, no patient reported diarrhea or abdominal pain or became noncompliant for other reasons."

Sullivan et al. (2004) studied the ability of *L. paracasei* ssp. *paracasei* strain F-19 to prevent the emergence of antibiotic-resistant microorganisms in patients receiving penicillin, ciprofloxacin, or norfloxacin. Twenty hospital patients (sex not reported) aged 18-89 years (mean = 59 years) receiving penicillin and 16 patients aged 28-86 years (mean = 61 years) receiving ciprofloxacin or norfloxacin were enrolled in a prospective, randomized, double-blind, placebo-controlled study in which half received  $2 \times 10^{10}$  cfu of *L. paracasei* ssp. *paracasei* strain F-19 daily for 14 days and half received placebo. Fecal samples were taken at baseline and on days 10 and 30 and isolates of enterococci, enterobacteria, and bacteroides were screened for antibiotic resistance.

The authors reported that, "No major differences were observed between the probiotic- and placebo-supplemented groups." They also reported that, "No adverse events were reported. Eight patients treated with penicillin reported looser stools, four in the active and four in the placebo group. In the quinolone-treated patients there were two individuals who experienced looser stools, one from each of the placebo and active groups."

Sixty otherwise apparently healthy women aged 18-40 years (mean =  $32.5 \pm 12/8$  years) with bacterial vaginosis were enrolled in a prospective, randomized, double-blind, placebo-controlled study (Delia et al. 2006) and treated with *Lactobacillus acidophilus* vaginal suppositories along with oral *L. paracasei* ssp. *paracasei* strain F-19 or placebo for 3 months. Both treatments were successful in reducing vaginal pH and subjective symptomatology scores. No adverse effects of the treatment were reported and the authors concluded, "The association of oral administration is useful to balance the vaginal environment with the intestinal microflora with improvement of long-term results."

Fourteen apparently healthy adults (5 males and 9 females aged 36-74 years) scheduled for routine colonoscopy were enrolled in an open-label study of the intestinal survival and persistence of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain NCFB 1748, and *B. animalis* ssp. *lactis* strain Bb-12 (Matto et al. 2006). The probiotics were provided in yogurt at daily levels of  $2 \times 10^{10}$  cfu for the 2 *Lactobacillus* strains and  $10^{11}$  cfu for the bifidobacteria for 10 days. Fecal samples were collected at the beginning and end of the ingestion period and at several time points up to 18-19 days after cessation of intake. Biopsy samples were taken at the end of ingestion, or 8-9 days later, or 18-19 days later. All samples were analyzed for bacterial content.

The F-19 and Bb-12 strains survived well and were detected in the feces of 100% and 79%, respectively, of the individuals consuming them while only 21% of them shed strain NCFB 1748. However, the probiotic bacteria washed out quickly and were rarely detected in fecal samples after a few weeks. Nor were the probiotics often isolated from colonic biopsies, suggesting that they do not exhibit strong adhesion properties. No adverse effects were reported by the authors.

To assess the effect of oral supplementation with synbiotics, glutamine, and peptide on intestinal permeability and the clinical outcome of critically ill trauma patients, Spindler-Vesel et al. (2007) enrolled 113 such patients (88 males and 25 females, mean age =  $41.0 \pm 18.9$  years) in a prospective, randomized, double-blind, placebo-controlled trial. Patients were randomized into 4 groups to receive glutamine (n = 32), soluble fiber (n = 29), peptide (n = 26), or fiber plus  $10^{10}$  cfu each of *L. paracasei* ssp. *paracasei* strain F-19, *L. plantarum* strain 2362, *Pediococcus pentosaceus* strain 5-33:3, and *Lactococcus raffinolactis* strain 32-77:1 (n = 26) for 7 days. Multiple organ failure was assessed daily and intestinal permeability was measured on days 2, 4, and 7.

No differences between groups were reported for multiple organ failure. However, out of a total of 51 bacterial infections, only 5 were observed in the synbiotic group, a significantly lower incidence than the other 3 groups. Only the synbiotic group showed evidence of decreased intestinal permeability. No adverse events associated with the synbiotic were reported. The authors concluded that, "Patients supplemented with synbiotics did better than the others, with lower intestinal permeability and fewer infections."

One hundred patients (38 male, 62 female, aged 18-68 years with mean =  $39 \pm 26$  years) suffering from irritable bowel syndrome (IBS) were enrolled in a prospective open-label study of the effectiveness of *L. paracasei* ssp. *paracasei* strain F-19 in treating the condition (Lombardo et al. 2009). Fifty two had diarrhea and 48 had constipation and all patients were given  $2.4 \times 10^{10}$  cfu per day of *L. paracasei* ssp. *paracasei* strain F-19 for 14 days. The probiotic treatment significantly reduced abdominal distension and pain and cured both diarrhea and constipation in more than 80% of those afflicted. The authors reported that tolerability of the probiotic was excellent, "No significant side effects that could definitely be attributed to the treatment were observed. Only one case of slight nausea was reported, but this did not require withdrawal from treatment."

In an open-label study, Sullivan et al. (2009) studied the effect of a probiotic mixture of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain NCFB 1748, and *B. animalis* ssp. *lactis* strain Bb12 on 15 chronic fatigue syndrome patients, 5 men and 10 women aged 30-56 years. Patients were observed for 2 weeks to establish a baseline of fecal bacteria and calprotectin levels, then they received  $4 \times 10^{10}$  cfu/day of the probiotic for 4 weeks, followed by a 4-week washout. Fatigue symptoms and fecal bacteria and calprotectin were measured at baseline, the end of the intervention, and the end of the washout period.

There were no significant changes in fatigue and physical activity scores and no major changes occurred in the gastrointestinal microflora; nevertheless, 6 of the 15 patients reported that they had improved. The authors did not report any adverse events. The authors concluded that the challenge is to "identify the responders to the therapy with probiotics as this pilot study demonstrates that

some individuals do respond with less fatigue, less bodily symptoms and better neurocognitive functions.”

Simren et al. (2010) reported a prospective, randomized, double-blind, placebo-controlled study of the effect of a probiotic mixture in patients with irritable bowel syndrome (IBS). A total of 74 IBS patients (22 males and 52 females with mean age =  $43.0 \pm 15.5$  years) were randomized to receive acidified milk with or without  $2 \times 10^{10}$  cfu of a mixture of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain La5, and *B. animalis* ssp. *lactis* strain Bb-12 for 8 weeks. The patients completed self-questionnaires weekly, blood was tested for clinical chemistries and hematology at baseline and following the intervention, and fecal samples were collected at the end of the intervention for enumeration of *L. paracasei* ssp. *paracasei* strain F-19 as a check on compliance.

IBS symptoms improved significantly in both groups over the study period, but with no significant difference between them. No clinically significant effects on biochemistry or hematology were noted and no adverse events were reported by the patients. The authors stated that, “The yoghurt was well tolerated.”

In a prospective, randomized, double-blind, placebo-controlled multicenter study of patients with irritable bowel syndrome (IBS), Sondergaard et al. (2011) enrolled 13 males and 39 females aged 29-67 years (mean age = 51.3 years) fulfilling Rome II criteria to consume acidified milk providing 0 or  $2.5 \times 10^{10}$  cfu of a combination of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain La5, and *B. animalis* ssp. *lactis* strain Bb-12 for 8 weeks. The patients completed weekly questionnaires and fecal samples were collected at the end of the intervention for enumeration of *L. paracasei* ssp. *paracasei* strain F-19 as a check on compliance.

While IBS symptoms significantly improved over the study period in both groups, there was no significant difference between treatment with acidified milk alone or acidified milk containing the probiotics. The authors reported that, “The investigational products were well tolerated and no serious adverse events occurred.”

Fifty patients (18 males and 32 females, mean age =  $65.2 \pm 8.1$  years) with symptomatic uncomplicated diverticular disease received a high-fiber diet and were randomized to the diet alone ( $n = 16$ ),  $2.4 \times 10^{10}$  cfu/day of *L. paracasei* ssp. *paracasei* strain F-19 ( $n = 18$ ), or  $4.8 \times 10^{10}$  cfu/day of the probiotic ( $N = 16$ ; Annibale et al. 2011) in a prospective, randomized, placebo-controlled, unblinded 6-month study. The probiotic was ingested for the first 14 days of each month. At enrolment and after 6 months, patients were examined for abdominal and dyspeptic symptoms and blood was taken for complete blood count, erythrocyte sedimentation rate, C-reactive protein, and protein electrophoresis.

Ingestion of the probiotic at either dose significantly reduced intestinal bloating and prolonged abdominal pain compared to the control group, with no difference between dosages. The authors concluded that, “Overall, the cyclic long-term (6 months) supplementation with *L. paracasei* sub. *paracasei* strain F-19 was found to be safe, since only in one patient was a side-effect (diarrhoea) found to lead to withdrawal from treatment.”

Pedersen et al. (2011) reported a prospective, randomized, double-blind, placebo-controlled study in which 61 patients (16 men and 45 women age 18-79 years; mean age = 42.5 years) with irritable bowel syndrome (IBS) consumed acidified milk with ( $n = 30$ ) or without ( $n = 31$ )  $2 \times 10^{10}$  cfu of each of the probiotics *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain LA-5, and *B. animalis* ssp. *lactis* strain Bb-12 for 8 weeks. Blood was drawn before and after the intervention and analyzed for serum glucose, lactate, glutamine, proline, creatinine/creatinine, and aspartic acid. While significant changes were reported in serum metabolites in both groups, these effects appeared to be due to the acidified milk since there were no significant differences between those receiving milk with or without the probiotics. The authors did not report any adverse events, and

concluded that “the delivering vector for probiotics, which was acidified milk in this study, can be very important for the study outcome.”

Begtrup et al. (2013) investigated the long-term effect of daily ingestion of  $5.2 \times 10^{10}$  cfu of a mixture of *L. paracasei* ssp. *paracasei* strain F-19, *L. acidophilus* strain La5, and *B. animalis* ssp. *lactis* strain Bb-12 in the management of irritable bowel syndrome (IBS). A total of 131 IBS patients (34 males, 97 females; mean age =  $30.52 \pm 9.42$  years) were enrolled in a prospective, randomized, double-blind, placebo-controlled study in which they received capsules containing either the probiotic mixture (n = 67) or placebo (n = 64) for 6 months with a 6-month follow-up. Patients visited a study nurse at 3, 6, and 12 months for questioning about compliance, symptoms, and adverse effects.

The authors concluded that, “During a 6-month treatment period, we were not able to detect a positive effect of probiotic when compared with placebo.” They also concluded that, “No serious adverse effects were reported.”

In a three-arm prospective, randomized, double-blind, placebo-controlled study of 58 obese postmenopausal women aged  $59.5 \pm 5.9$  years, Brahe et al. (2015) investigated the effect of daily ingestion of  $9.4 \times 10^{10}$  cfu of *L. paracasei* ssp. *paracasei* strain F-19 or of 10 g flaxseed mucilage on gut microbiota and metabolic risk markers. Nineteen women each were randomized to receive either probiotic or mucilage for 6 weeks while 20 women received maltodextrin placebo. Bacterial viability was confirmed at the end of the intervention. The participants were interviewed about all types of potential adverse effects at each visit by the use of broad, open-ended questions, and asked about changes in stool characteristics. Additionally, body composition was assessed; blood samples were taken for analysis of glucose, insulin, total cholesterol, LDL- and HDL-cholesterol, triacylglycerol, leucocytes, C-reactive protein, lipopolysaccharide-binding protein, TNF- $\alpha$ , IL-6, and angiopoietin-like protein 4; and DNA in fecal samples was analyzed for identification of shed bacteria.

Intake of *L. paracasei* ssp. *paracasei* strain F-19 did not modulate metabolic markers compared with placebo. The authors reported on probiotic-related adverse events as follows:

“More adverse events were reported in the flaxseed group compared with the placebo group, while there was no difference in the occurrence of adverse events between the placebo and the *L. paracasei* F-19 group. . . There was no difference between the adverse events reported following the probiotic and placebo intervention that included both more frequent and less-frequent defecation. No serious adverse events were registered during the study” (Brahe et al. 2015).

Compare et al. (2015) reported a prospective, randomized, double-blind, placebo-controlled, multicenter study of the effect of *L. paracasei* ssp. *paracasei* strain F-19 on bowel symptom onset in gastroesophageal reflux disease (GERD) patients on long-term proton-pump inhibitors. The study had 4 arms:  $2.4 \times 10^{10}$  cfu of F-19 for 3 days per week for 6 months,  $2.4 \times 10^{10}$  cfu of F-19 for 3 days/week for 3 months then placebo for 3 months, placebo for 3 months then  $2.4 \times 10^{10}$  cfu of F-19 for 3 days/week for 3 months, and placebo for 6 months. The 100 enrolled patients (56 male, 44 female; mean age =  $39 \pm 10.4$  years) were assigned 25 per arm. Patients completed questionnaires each month to assess bowel habits and symptoms.

Treatment with *L. paracasei* ssp. *paracasei* strain F-19 significantly reduced flatulence and bloating but not abdominal pain. There was no discussion of any adverse effects of the probiotic intervention.

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference                                                                                                                                                                     | Objective                                                                      | Study Design                                                                              | Subjects                                          | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|---------------------------------------------------|------------------------------------------------------------|----------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Studies in Infants</b>                                                                                                                                                     |                                                                                |                                                                                           |                                                   |                                                            |          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Szajewska et al. 2017                                                                                                                                                         | Growth of infants receiving formula supplemented with prebiotics or synbiotics | Prospective, randomized, double-blind, parallel-group, multicenter                        | 182 healthy term infants, 93M 89F, aged 28 days   | $10^9$ cfu/day                                             | 5 months | There was no difference in growth between the two groups. The authors reported that, "Probiotic supplementation of infant formula with <i>Lactobacillus</i> F-19 was well tolerated, and no significant differences between the experimental and control groups were observed in regard to adverse events," and concluded: "The lack of a significant difference between the formula-fed groups for growth and for the occurrence of serious adverse events supports the safety of using <i>Lactobacillus</i> F-19-supplemented synbiotic formula in healthy term infants."                                                                                                              |
| West et al. 2008<br>West et al. 2009<br>West et al. 2012a<br>West et al. 2013<br>Chorell et al. 2013<br>Hasslof et al. 2013<br>Videhult et al. 2015a<br>Videhult et al. 2015b | Effects of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 in infants    | Prospective, randomized, double-blind, placebo-controlled with follow-up at age 8-9 years | 179 healthy term infants, 75M 104F, aged 4 months | $10^8$ cfu/day                                             | 9 months | 5 infants from the test group and 3 from the controls were withdrawn by their parents for reasons not associated with the intervention. Days receiving antibiotics were fewer among infants ingesting F-19. No adverse effects of the probiotic were reported.<br><br>In additional analyses of the data from this study, neither benefits nor adverse effects were reported.<br><br>In follow-ups with infants when they reached the age of 8-9 years, the authors concluded that "we observed no adverse effect of LF-19 on body composition or metabolism in this follow-up study," and that "the probiotic LF-19 had no long-term effect on the inflammatory and metabolic profile." |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference                  | Objective                                                                                 | Study Design                                              | Subjects                                                                                                     | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|----------------------------|-------------------------------------------------------------------------------------------|-----------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Zampieri et al. 2013       | Prevention of progression of NEC from Bell's stage 2                                      | Prospective, randomized, unblinded                        | 32 preterm infants (all delivered before gestational week 32; birth weight 600-1500 g) at Bell's stage 2 NEC | $6 \times 10^9$ cfu/day                                    | 21 days  | The authors reported that, "No collateral effects were observed during the study period; none of our patients presented sepsis due to <i>L. paracasei</i> subsp. <i>paracasei</i> strain F-19. No patient required preventive exclusion from the trial. No patient [in the probiotic group] presented treatment-related intestinal complications (i.e., diarrhea)." Their conclusion was that "The use of <i>L. paracasei</i> subsp. <i>paracasei</i> strain F-19 is safe and effective." |
| <b>Studies in Children</b> |                                                                                           |                                                           |                                                                                                              |                                                            |          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Crittenden et al. 2002     | Assess the suitability of strain F-19 as a probiotic                                      | Prospective, randomized, double-blind, placebo-controlled | 61 apparently healthy young children aged 1-1.5 years                                                        | $2 \times 10^{10}$ cfu/day                                 | 3 weeks  | The study design included observations monitoring potential side-effects of probiotic consumption, including intestinal discomfort, increased flatulence, and changes in stool consistency and frequency. No such adverse effects were observed and the probiotic was well tolerated by all individuals. The authors concluded that, "The fact that no side-effects were detected in subjects . . . suggests that <i>Lactobacillus</i> F-19 is a safe microbial food supplement."         |
| Sullivan et al. 2002       | Effect of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 on the colonic microbiota | Prospective, randomized, double-blind, placebo-controlled | 61 apparently healthy and fully weaned children (mean age = $13.1 \pm 5.3$ years)                            | $10^{10}$ cfu/day                                          | 3 weeks  | Children receiving <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 showed a significant increase in fecal levels of the strain, but not of other lactobacilli or bifidobacteria. The authors reported that, "The products were generally well tolerated . . . Fecal characteristics and occurrences of gastrointestinal symptoms were similar in the treatment and placebo groups."                                                                                                 |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference                | Objective                                                              | Study Design                                                            | Subjects                                                                                               | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                            | Duration                   | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------|------------------------------------------------------------------------|-------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Studies in Adults</b> |                                                                        |                                                                         |                                                                                                        |                                                                                                                                                                       |                            |                                                                                                                                                                                                                                                                                                                                                                                                              |
| Annibale et al. 2011     | Treatment of symptomatic uncomplicated diverticular disease            | Prospective, randomized, placebo-controlled, unblinded                  | 50 patients (18M, 32F) with symptomatic uncomplicated diverticular disease, mean age = 65.2±8.1 years) | 4.8x10 <sup>10</sup> cfu/day                                                                                                                                          | 14 days/month for 6 months | Ingestion of the probiotic significantly reduced intestinal bloating and prolonged abdominal pain compared to the control group. The authors concluded that, "Overall, the cyclic long-term (6 months) supplementation with <i>L. paracasei</i> sub. <i>paracasei</i> strain F-19 was found to be safe, since only in one patient was a side-effect (diarrhoea) found to lead to withdrawal from treatment." |
| Begtrup et al. 2013      | Long-term effect of daily ingestion of probiotics in management of IBS | Prospective, randomized, double-blind, placebo-controlled               | 131 IBS patients (34M, 97F); mean age = 30.52±9.42 years                                               | 5.2x10 <sup>10</sup> cfu/day of a mixture of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. acidophilus</i> , and <i>B. animalis</i> ssp. <i>lactis</i> |                            | The authors concluded that, "During a 6-month treatment period, we were not able to detect a positive effect of probiotic when compared with placebo." They also concluded that, "No serious adverse effects were reported."                                                                                                                                                                                 |
| Brahe et al. 2015        | Effect on gut microbiota and metabolic risk markers in obese women     | Three-arm prospective, randomized, double-blind, placebo-controlled     | 58 obese postmenopausal women aged 59.5±5.9 years                                                      | 9.4x10 <sup>10</sup> cfu/day                                                                                                                                          | 6 weeks                    | The authors stated that "there was no difference in the occurrence of adverse events between the placebo and the <i>L. paracasei</i> F-19 group. . . There was no difference between the adverse events reported following the probiotic and placebo intervention that included both more frequent and less-frequent defecation. No serious adverse events were registered during the study."                |
| Compare et al. 2015      | Effect on symptom onset in gastro-esophageal reflux disease            | Prospective, randomized, double-blind, placebo-controlled, multi-center | 100 GERD patients (56M, 44 F); mean age = 39±10.4 years                                                | 2.4x10 <sup>10</sup> cfu/day                                                                                                                                          | 3 days/week for 6 months   | Treatment with <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 significantly reduced flatulence and bloating but not abdominal pain. There was no discussion of any adverse effects of the probiotic intervention.                                                                                                                                                                                     |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference              | Objective                                            | Study Design                                             | Subjects                                                  | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|------------------------|------------------------------------------------------|----------------------------------------------------------|-----------------------------------------------------------|------------------------------------------------------------|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Crittenden et al. 2002 | Assess the suitability of strain F-19 as a probiotic | Open-label                                               | 5 apparently healthy adults                               | $2 \times 10^{10}$ cfu/day                                 | 12 days  | Side effects monitored included intestinal discomfort, increased flatulence, and changes in stool consistency and frequency. The authors concluded that, "No adverse effects of probiotic administration were observed in any of the pilot studies. The probiotic was well tolerated by all individuals, including the elderly <i>H. pylori</i> patients who consumed <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 daily for 3 months without adverse effects. The fact that no side-effects were detected in subjects ranging in age from 1 to 85 years, and in healthy individuals and subjects suffering mild illnesses (milk-hypersensitivity and <i>H. pylori</i> infection) suggests that <i>Lactobacillus</i> F-19 is a safe microbial food supplement." |
| Crittenden et al. 2002 | Assess the suitability of strain F-19 as a probiotic | Prospective, single-blind, placebo-controlled, crossover | 9 apparently healthy adults, 4 with milk hypersensitivity | $2 \times 10^{10}$ cfu/day                                 | 1 week   | Side effects monitored included intestinal discomfort, increased flatulence, and changes in stool consistency and frequency. The authors concluded that, "No adverse effects of probiotic administration were observed. The probiotic was well tolerated by all individuals. The fact that no side-effects were detected in subjects ranging in age from 1 to 85 years, and in healthy individuals and subjects suffering mild illnesses (milk-hypersensitivity and <i>H. pylori</i> infection) suggests that <i>Lactobacillus</i> F-19 is a safe microbial food supplement."                                                                                                                                                                                            |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference              | Objective                                                 | Study Design                                              | Subjects                                                                                                       | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                                   | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|------------------------|-----------------------------------------------------------|-----------------------------------------------------------|----------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Crittenden et al. 2002 | Assess the suitability of strain F-19 as a probiotic      | Prospective, randomized, double-blind, placebo-controlled | 30 elderly (>65 years) patients seropositive to <i>H. pylori</i>                                               | $7.5 \times 10^{10}$ cfu/day                                                                                                                                                 | 12 weeks | Side effects monitored included intestinal discomfort, increased flatulence, and changes in stool consistency and frequency. The authors concluded that, "No adverse effects of probiotic administration were observed. The probiotic was well tolerated by all individuals, including the elderly <i>H. pylori</i> patients who consumed <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19 daily for 3 months without adverse effects. The fact that no side-effects were detected in subjects ranging in age from 1 to 85 years, and in healthy individuals and subjects suffering mild illnesses (milk-hypersensitivity and <i>H. pylori</i> infection) suggests that <i>Lactobacillus</i> F-19 is a safe microbial food supplement." |
| Delia et al. 2006      | Treatment of bacterial vaginosis                          | Prospective, randomized, double-blind, placebo-controlled | 60 otherwise apparently healthy women aged 18-40 years (mean = $32.5 \pm 12/8$ years) with bacterial vaginosis | Dose not reported                                                                                                                                                            | 3 months | No adverse effects of the treatment were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Liu et al. 2004        | Treatment of patients with minimal hepatic encephalopathy | Prospective, randomized, double-blind, placebo-controlled | 55 patients aged $56 \pm 11$ years with minimal hepatic encephalopathy                                         | $10^{10}$ cfu/day each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>Pediococcus pentosaceus</i> , <i>Leuconostoc mesenteroides</i> , and <i>L. plantarum</i> | 30 days  | The authors reported that, "The synbiotic, fermentable fiber and placebo preparations were well-tolerated by all patients, with no reports of adverse side effects. In particular, no patient reported diarrhea or abdominal pain or became noncompliant for other reasons."                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference            | Objective                                                                                     | Study Design                                              | Subjects                                                                                 | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                  | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                |
|----------------------|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------|------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Lombardo et al. 2009 | Treatment of IBS                                                                              | Open-label                                                | 100 patients (38M, 62F) aged 18-68 years with mean = 39±26 years) with IBS               | 2.4x10 <sup>10</sup> cfu/day                                                                                                                                | 14 days  | The authors reported that tolerability of the probiotic was excellent; "No significant side effects that could definitely be attributed to the treatment were observed. Only one case of slight nausea was reported, but this did not require withdrawal from treatment."                                                                                                             |
| Matto et al. 2006    | Study the intestinal survival and persistence of probiotics                                   | Open-label                                                | 14 apparently healthy adults (5M, 9F) aged 36-74 years scheduled for routine colonoscopy | 2x10 <sup>10</sup> cfu/day                                                                                                                                  | 10 days  | The F-19 survived well and were detected in the feces of 100% of the individuals consuming them. However, they washed out quickly and were rarely detected in fecal samples after a few days or weeks. Nor were the probiotics often isolated from colonic biopsies, suggesting that they do not exhibit strong adhesion properties. No adverse effects were reported by the authors. |
| Pedersen et al. 2011 | Treatment of IBS                                                                              | Prospective, randomized, double-blind, placebo-controlled | 61 IBS patients (16M, 45W) aged 18-79 years; mean age = 42.5 years)                      | 2x10 <sup>10</sup> cfu/day each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. acidophilus</i> , and <i>B. animalis</i> ssp. <i>lactis</i> | 8 weeks  | There were no significant differences between those receiving milk with or without the probiotics. The authors did not report any adverse events.                                                                                                                                                                                                                                     |
| Rayes et al. 2002a   | Reduce post-operative bacterial infection in patients recovering from major abdominal surgery | Prospective, randomized, double-blind, placebo-controlled | 95 patients (49M, 46F) with mean age = 49.0±2.5 years recovering from abdominal surgery  | 2x10 <sup>9</sup> cfu/day                                                                                                                                   | 10 days  | Both standard enteral formula and strain F-19 were well tolerated. Abdominal side effects (distension, cramps, or diarrhea) were seen in 8 of 32 patients in the standard-formula group, 6 of 31 patients in the live <i>Lactobacillus</i> group, and 11 of 32 patients in the inactivated-bacteria group.                                                                            |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference          | Objective                                                                                     | Study Design                                              | Subjects                                                                                          | <i>L. paracasei</i><br>ssp. <i>paracasei</i><br>Strain F-19<br>Dose | Duration     | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|--------------------|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------|---------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|--------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Rayes et al. 2002b | Reduce post-operative bacterial infection in patients recovering from major abdominal surgery | Prospective, randomized, double-blind, placebo-controlled | 172 patients following major abdominal surgery or liver transplantation                           | $2 \times 10^9$ cfu/day                                             | Not reported | The groups receiving live <i>Lactobacillus</i> cultures experienced significantly reduced incidences of bacterial infection (among resection patients, 4% vs. 13% in the prebiotic group and 31% in the conventional-therapy group; among transplant recipients, 13% in the probiotic group vs. 34% in the prebiotic group and 48% in the conventional therapy group). The authors reported that "Fibre and lactobacilli were well tolerated in most cases."                                                                                                                                                                                                                                                                                                                                              |
| Rayes et al. 2002c | Reduce post-operative bacterial infection in patients recovering from major abdominal surgery | Prospective, randomized, double-blind, placebo-controlled | 90 patients (48M, 42F), mean age = $61.0 \pm 14.1$ years, recovering from major abdominal surgery | $2 \times 10^9$ cfu/day                                             | 10 days      | All patients completed the study, receiving the full course of their assigned nutrition. The incidence of infections was significantly lower in groups receiving fiber and either live or killed lactobacilli (10% each) than in the group receiving standard nutritional formula (30%). Patients receiving live F-19 strain needed antibiotics for a significantly shorter time than did the patients in the other groups. The length of hospital stay and the incidence of non-infectious complications did not differ significantly. Fibers and lactobacilli were well tolerated; side effects did not differ between groups and there was no diarrhea. The authors concluded that "application of fiber and lactobacilli is feasible even after major abdominal surgery and caused few side effects." |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference         | Objective                                                                                     | Study Design                                              | Subjects                                                                                                     | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                                            | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-------------------|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Rayes et al. 2005 | Reduce post-operative bacterial infection in patients recovering from liver transplantation   | Prospective, randomized, double-blind, placebo-controlled | 66 liver transplant recipients (38M, 28F), mean age = 51.5±2.5 years)                                        | 2x10 <sup>10</sup> cfu/day each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>Pediococcus pentosaceus</i> , <i>Leuconostoc mesenteroides</i> , and <i>L. plantarum</i> | 14 days  | Inclusion of probiotics in the enteral nutrition formula significantly reduced the incidence of post-operative bacterial infection and reduced the severity of infections that did occur. Hematology, immunology, and biochemistry parameters did not differ significantly between groups. The authors reported that "Enteral nutrition, containing the synbiotic combination, was well tolerated in all patients," with fewer probiotic patients exhibiting diarrhea or abdominal cramps. "All side effects disappeared under temporary reduction in the amount of enteral nutrition." The authors concluded that inclusion of probiotics "is an effective means to prevent post-operative bacterial infections in high-risk surgical patients, and since it causes no resistant strains and has no serious side effects, it could be widely used."        |
| Rayes et al. 2007 | Reduce post-operative bacterial infection in patients recovering from major abdominal surgery | Prospective, randomized, double-blind, placebo-controlled | 80 patients (45M, 35F, mean age = 58.5±12.5 years) recovering from pylorus-preserving pancreato-duodenectomy | 2x10 <sup>10</sup> cfu/day each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>Pediococcus pentosaceus</i> , <i>Leuconostoc mesenteroides</i> , and <i>L. plantarum</i> | 8 days   | Inclusion of probiotics in the enteral nutrition formula significantly reduced the incidence and duration of post-operative bacterial infection to 12.5% vs 40% in the fiber-only formula group. The incidences of diarrhea and abdominal cramps were the same in both groups of patients. The authors reported that, "Enteral nutrition, containing the synbiotic combination, was well tolerated in all patients... All side effects disappeared under temporary reduction in the amount of enteral nutrition," and concluded, "Early enteral nutrition with synbiotics was able to significantly reduce postoperative bacterial infections in patients following [pancreatoduodenectomy] with only single-shot antibiotic prophylaxis. In contrast to antibiotics, it is relatively cheap and does not cause resistant strains or serious side effects." |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference           | Objective                                                                                     | Study Design                                              | Subjects                                                            | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                                            | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                |
|---------------------|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------|---------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Rayes et al. 2012   | Reduce post-operative bacterial infection in patients recovering from major abdominal surgery | Prospective, randomized, double-blind, placebo-controlled | 19 right-hepatectomy patients (14M, 5F), mean age = 60.1±13.9 years | 2x10 <sup>10</sup> cfu/day each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>Pediococcus pentosaceus</i> , <i>Leuconostoc mesenteroides</i> , and <i>L. plantarum</i> | 10 days  | The authors concluded that, "The synbiotic combination was well-tolerated in all patients. Mild side-effects (abdominal distension and cramps) occurred in three patients of each group but disappeared under symptomatic therapy. No severe side effects were recorded."                                             |
| Riordan et al. 2003 | Effect of F-19 on TNF-α level in cirrhosis patients                                           | Open-label                                                | 11 cirrhosis patients and 5 healthy controls                        | 8x10 <sup>10</sup> cfu/day each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. plantarum</i> , <i>Pediococcus pentosaceus</i> , and <i>Lactococcus raffinolactis</i> | 7 days   | The authors reported that, "Administration of the synbiotic supplement was well tolerated without any reported adverse events or change in general clinical state."                                                                                                                                                   |
| Simren et al. 2010  | Treatment of IBS                                                                              | Prospective, randomized, double-blind, placebo-controlled | 74 IBS patients (22M, 52F) with mean age = 43.0±15.5 years)         | 2x10 <sup>10</sup> cfu/day of a mixture of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. acidophilus</i> , and <i>B. animalis</i> ssp. <i>lactis</i>                   | 8 weeks  | IBS symptoms improved significantly in both groups over the study period, but with no significant difference between them. No clinically significant effects on biochemistry or hematology were noted and no adverse events were reported by the patients. The authors stated that, "The yoghurt was well tolerated." |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference                  | Objective                                                                                    | Study Design                                                           | Subjects                                                                                             | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                               | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                       |
|----------------------------|----------------------------------------------------------------------------------------------|------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sondergaard et al. 2011    | Treatment of IBS                                                                             | Prospective, randomized, double-blind, placebo-controlled multi-center | 52 patients (13M, 39F) aged 29-67 years (mean age = 51.3 years) fulfilling Rome II criteria          | $2.5 \times 10^{10}$ cfu/day of a mixture of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. acidophilus</i> , and <i>B. animalis</i> ssp. <i>lactis</i>    | 8 weeks  | There was no significant difference between treatment with acidified milk alone or acidified milk containing the probiotics. The authors reported that, "The investigational products were well tolerated and no serious adverse events occurred."                                                                                                                                                                                           |
| Spindler-Vesel et al. 2007 | Effect on intestinal permeability and the clinical outcome of critically ill trauma patients | Prospective, randomized, double-blind, placebo-controlled              | 113 trauma patients (88M, 25F), mean age = $41.0 \pm 18.9$ years)                                    | $10^{10}$ cfu each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. plantarum</i> , <i>Pediococcus pentosaceus</i> , and <i>Lactococcus raffinolactis</i> | 7 days   | Out of 51 bacterial infections, only 5 were observed in the synbiotic group, a significantly lower incidence than the other 3 groups. Only the synbiotic group showed evidence of decreased intestinal permeability. No adverse events associated with the synbiotic were reported. The authors concluded that, "Patients supplemented with synbiotics did better than the others, with lower intestinal permeability and fewer infections." |
| Sullivan et al. 2001       | Treatment of elderly <i>Helicobacter pylori</i> patients                                     | Prospective, randomized, double-blind, placebo-controlled              | 35 <i>H. pylori</i> patients (sex not reported; age range 58-89 years with a median age of 76 years) | $10^{11}$ cfu/day                                                                                                                                                        | 12 weeks | . The authors concluded that the probiotic "survives the passage through the gastrointestinal tract." There was no discussion of any adverse events.                                                                                                                                                                                                                                                                                         |
| Sullivan et al. 2002       | Treatment of elderly <i>Helicobacter pylori</i> patients                                     | Prospective, randomized, double-blind, placebo-controlled              | 30 elderly <i>H. pylori</i> patients (sex not reported; mean age = $76.8 \pm 6.5$ years)             | $10^{11}$ cfu/day                                                                                                                                                        | 12 weeks | The authors reported that, "The products were generally well tolerated . . . Fecal characteristics and occurrences of gastrointestinal symptoms were similar in the treatment and placebo groups."                                                                                                                                                                                                                                           |

**Table 5. Studies of *L. paracasei* ssp. *paracasei* Strain F-19.**

| Reference            | Objective                                                                                                                           | Study Design                                              | Subjects                                                                                                                                                               | <i>L. paracasei</i> ssp. <i>paracasei</i> Strain F-19 Dose                                                                                                          | Duration | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                  |
|----------------------|-------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sullivan et al. 2003 | Prevention of antibiotic-associated ecological disturbances of the intestinal microbiota                                            | Prospective, randomized, double-blind, placebo-controlled | 24 apparently healthy adults (7M, 17W) aged 21-48 years receiving antibiotic therapy                                                                                   | $2 \times 10^8$ cfu of each of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. acidophilus</i> , and <i>B. animalis</i> ssp. <i>lactis</i>             | 14 days  | Consumption of the probiotic prevented ecological disturbances in the numbers of lactobacilli and <i>Bacteroides fragilis</i> group. One participant in the probiotic group developed diarrhea and one reported looser stools. The first person's stool was cytotoxin positive, cured by metronidazole; the loose stools resolved spontaneously. No other adverse events were reported. |
| Sullivan et al. 2004 | Prevention of the emergence of antibiotic-resistant micro-organisms in patients receiving penicillin, ciprofloxacin, or norfloxacin | Prospective, randomized, double-blind, placebo-controlled | 20 hospital patients aged 18-89 years (mean = 59 years) receiving penicillin and 16 patients aged 28-86 years (mean = 61 years) receiving ciprofloxacin or norfloxacin | $2 \times 10^{10}$ cfu/day                                                                                                                                          | 14 days  | The authors reported that, "No adverse events were reported. Eight patients treated with penicillin reported looser stools, four in the active and four in the placebo group. In the quinolone-treated patients there were two individuals who experienced looser stools, one from each of the placebo and active groups."                                                              |
| Sullivan et al. 2009 | Treatment of chronic fatigue syndrome                                                                                               | Open-label                                                | 15 chronic fatigue syndrome patients, (5M, 10W) aged 30-56 years                                                                                                       | $4 \times 10^{10}$ cfu/day of a mixture of <i>L. paracasei</i> ssp. <i>paracasei</i> strain F-19, <i>L. acidophilus</i> , and <i>B. animalis</i> ssp. <i>lactis</i> | 4 weeks  | There were no significant changes in fatigue and physical activity scores and no major changes occurred in the gastrointestinal microflora; nevertheless, 6 of the 15 patients reported that they had improved. The authors did not report any adverse events.                                                                                                                          |

### **6.5. Studies of Other Strains of *L. paracasei* ssp. *paracasei***

Human studies of other strains of the species are summarized in Table 6. These fifty studies include 28 different strains of *L. paracasei* ssp. *paracasei*: CUL08, LPC-S01, LPC-37, W8, ST11, CBA L74, LP-33, DG, CNCM I-2116, B21060, NCC 2461, BRAP01, 317, CNCM I-1518, HF A000232, 431, IMC502, HN019, N24, N1115, CNCM I-4034, 8700, IMPC 2.1, SD1, K71, IMC 502, LPC09, and LMGP22043. No adverse events clearly attributable to the probiotic were reported in any study.

The studies summarized in Table 6 do not include the extensive literature on *L. casei* strain Shirota, the subject of GRN 000429, although the GRAS submission observes:

“The taxonomy of the genus *Lactobacillus* is currently in a state of flux, and nowhere is taxonomic placement less certain than in the complex of *L. casei* strains. According to the most recent opinion (as of this writing) of the Judicial Commission of the International Committee on Systematics of Bacteria in 2008 (JCICSB 2008), *L. casei* strain Shirota would now be regarded as a strain of *L. paracasei* rather than *L. casei*” (p8).

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference              | Objective                                                                              | Study Design                                                                | Subjects                                            | Strain, Dose, and Duration                                                                                                                                                                                        | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------|----------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Allen et al. 2014      | Evaluate efficacy in prevention of eczema.                                             | Prospective, randomized, double-blind, placebo-controlled, parallel-group   | 454 pregnant women and their infants                | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CUL08, $1.25 \times 10^9$ cfu (& <i>L. salivarius</i> , <i>B. animalis</i> ssp. <i>lactis</i> , and <i>B. bifidum</i> ), mother ~4 weeks, infant to age 6 months | Authors' conclusion: "Probiotic administration was not associated with adverse effects in either mothers or their infants."                                                                                                                                                                                                                                                                                                                                   |
| Balzaretti et al. 2015 | Assess use of <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC-S01 as a probiotic. | Prospective, randomized, double-blind, placebo-controlled, cross-over       | 11 apparently healthy adults                        | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC-S01, $2.4 \times 10^{10}$ cfu, 7 days                                                                                                                        | "Capsules, which contained about 24 billion CFU of <i>L. paracasei</i> , were well tolerated by all participants and no adverse events were reported."<br>Authors' conclusion: " <i>L. paracasei</i> LPC-S01 is a safe bacterial strain for human consumption, which does not contain any acquired antibiotic resistance, does not produce biogenic amines and can be administered in high number (24 billion CFU) to healthy people without adverse events." |
| Barker et al. 2017     | Evaluate use against <i>C. difficile</i> infection.                                    | Prospective, randomized, double-blind, placebo-controlled, pilot trial      | 33 patients with mild <i>C. difficile</i> infection | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC-37 (& <i>B. lactis</i> and <i>L. acidophilus</i> ), 28 days                                                                                                  | "There was no significant difference in the rate of CDI recurrence or functional improvement over time between treatment groups."                                                                                                                                                                                                                                                                                                                             |
| Bjerg et al 2015a,b    | Effect of probiotic on blood TAG levels.                                               | Prospective, randomized, double-blind, placebo-controlled, two-arm parallel | 64 apparently healthy young adults                  | <i>L. paracasei</i> ssp. <i>paracasei</i> strain W8, $10^{10}$ cfu, 4 weeks                                                                                                                                       | "Four weeks supplementation with <i>L. casei</i> W8 did not affect the overall composition of the gut microbiota."                                                                                                                                                                                                                                                                                                                                            |
| Chouraqui et al. 2008  | Effect on risk of diarrhea in infants.                                                 | Prospective, randomized, double-blind, placebo-controlled, parallel group   | 284 healthy term infants                            | <i>L. paracasei</i> ssp. <i>paracasei</i> strain ST11 (& <i>B. longum</i> ), 14 weeks                                                                                                                             | Infants in all groups had similar weight gain, length, head circumference, digestive tolerance, and adverse events at 2, 4, 8, 12, 16, and 52 weeks of age. The authors concluded that: "Infants fed formulas containing probiotics or synbiotics show a similar rate in weight gain compared with those fed a control formula and tolerate these formulas well."                                                                                             |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference            | Objective                                                                   | Study Design                                                                    | Subjects                                                                    | Strain, Dose, and Duration                                                                   | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|----------------------|-----------------------------------------------------------------------------|---------------------------------------------------------------------------------|-----------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Corsello et al. 2017 | Effect on risk of common infectious diseases in children.                   | Prospective, randomized, double-blind, placebo-controlled, multi-center         | 146 health children (84M, 62F) aged 12-48 months (mean = 33±9 months)       | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CBA L74, 5.9x10 <sup>11</sup> cfu, 3 months | "No changes were observed for body weight and height in the two study groups, indicating that the consumption of the study products was safe, at least in the short term. No adverse events related to the consumption of the active or placebo products were recorded." The authors concluded that: "The dietary supplementation was well accepted by the children and safe, as demonstrated by the low dropout rate together with the high level of adherence, and the absence of ad-verse events observed during the study period." |
| Costa et al. 2014    | Study effect of probiotics in patients with allergic rhinitis.              | Prospective, randomized, double-blind, placebo-controlled                       | 425 patients with allergic rhinitis to grass pollen treated with loratadine | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LP-33, 5 weeks                              | Improved quality of life; no adverse events reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Farrario et al. 2014 | Ability of probiotic to modulate fecal Clostridiales bacteria and butyrate. | Prospective, randomized, double-blind, placebo-controlled, crossover design     | Apparently healthy adults age 23-55 years                                   | <i>L. paracasei</i> ssp. <i>paracasei</i> strain DG 2.4x10 <sup>10</sup> cfu, 4 weeks        | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Gore et al. 2012     | Treatment and secondary prevention of early infant eczema.                  | Prospective, randomized, double-blind, placebo-controlled with 3-year follow-up | 208 infants aged 3-6 months with eczema                                     | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CNCM I-2116, 3 months                       | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Grossi et al. 2010   | Treatment of acute diarrhea.                                                | Prospective, randomized, double-blind, placebo-controlled                       | 174 adult patients with acute diarrhea                                      | <i>L. paracasei</i> ssp. <i>paracasei</i> strain B 21060, 10 days                            | "The 2 treatments showed a very good tolerability profile, with negligible and similar adverse event rates and similar concomitant medication usage rates."                                                                                                                                                                                                                                                                                                                                                                            |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference             | Objective                                                        | Study Design                                                                            | Subjects                                                                         | Strain, Dose, and Duration                                                                         | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|-----------------------|------------------------------------------------------------------|-----------------------------------------------------------------------------------------|----------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Gueniche et al. 2014  | Effect of probiotic on skin reactivity.                          | Prospective, randomized, double-blind, placebo-controlled                               | 64 women with reactive/ sensitive skin                                           | <i>L. paracasei</i> ssp. <i>paracasei</i> strain NCC 2461, 60 days                                 | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Hemalatha et al. 2017 | Effect of probiotic on SCFA in children.                         | Prospective, randomized, double-blind, placebo-controlled                               | 379 apparently healthy children aged 2-5 years                                   | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC-37, $2 \times 10^9$ cfu, 9 months             | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Ho et al. 2014        | Effect of probiotic on cytotoxicity of human NK cells.           | Prospective multi-arm open-label study                                                  | 100 apparently healthy adults with average age = 55 years                        | <i>L. paracasei</i> ssp. <i>paracasei</i> strain BRAP01, $2 \times 10^{10}$ CFU, 1 day             | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Hutt et al. 2011      | Evaluate the safety of probiotics.                               | Prospective open-label study                                                            | 9 apparently healthy adults                                                      | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 317 (vs. other probiotics), $10^{10}$ cfu, 5 days | Data on gut health, blood parameters, and liver and kidney function were collected. The authors reported that "The administration of high doses of different <i>Lactobacillus</i> strains did not result in any severe adverse effects in GIT and/or abnormal values of blood indices."                                                                                                                                                                                                                                                                                                                           |
| Jespersen et al. 2015 | Effect of probiotic on immune response to influenza vaccination. | Prospective, randomized, double-blind, placebo-controlled, parallel-group, multi-center | 1104 apparently healthy adults (453M, 651F) aged 18-60 years (mean = 31.4 years) | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 431, $10^9$ cfu, 42 days                          | The authors reported that: "A total number of 2212 AEs in 914 subjects were reported during the study. Of these, 41 events in 34 subjects (21 in the probiotic group and 20 in the placebo group) were assessed as study product related. The most prevalent of the product-related AEs were gastrointestinal disorders (48% of events) and nasopharyngitis (29%). In total, 373 events in 344 subjects (186 in the probiotic group and 187 in the placebo group) were assessed as vaccine related. Five AEs were defined as serious; none of these were assessed to be related to the study product or vaccine." |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference                   | Objective                                               | Study Design                                                                       | Subjects                                                                                      | Strain, Dose, and Duration                                                                                                         | Safety-Related Results                                                                                                                                                                                                                                                                     |
|-----------------------------|---------------------------------------------------------|------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Lahner et al 2012           | Use of probiotic in treatment of diverticular disease.  | Prospective, randomized, double-blind, placebo-controlled, multi-center            | 52 patients (17M, 35F) aged 40-80 years with diverticular disease (mean age = 66.3±9.6 years) | <i>L. paracasei</i> ssp. <i>paracasei</i> strain B21060, 5x10 <sup>9</sup> cfu, 6 months                                           | The authors reported that: "None of the patients developed altered biochemical inflammatory parameters, acute diverticulitis or other diverticular disease complications throughout the 6-mo study period. In both groups no adverse event was registered over the 6-mo treatment period." |
| Lee et al. 2017             | Effect of probiotics on immune function.                | Prospective, randomized, placebo-controlled, open-label                            | 152 healthy adults (45M, 107F) aged >60 years (mean age = 65.7±0.4 years)                     | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 431, 1.2x10 <sup>9</sup> cfu, 12 weeks (& <i>B. animalis</i> ssp. <i>lactis</i> ) | The authors reported that: "No adverse events were reported from the participants."                                                                                                                                                                                                        |
| Lenoir-Wijnkoop et al. 2014 | Prevention of antibiotic-associated diarrhea.           | Prospective, randomized, double-blind, placebo-controlled                          | 135 elderly hospitalized patients (mean age = 74 years)                                       | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CNCM I-1518, approx.. 2 weeks                                                     | The authors reported that: "There were no adverse events in the probiotic group."                                                                                                                                                                                                          |
| Lin et al. 2014             | Treatment of children with perennial allergic rhinitis. | Prospective, randomized, double-blind, placebo-controlled                          | 60 children (47M, 13F) aged 6-13 years with allergic rhinitis                                 | <i>L. paracasei</i> ssp. <i>paracasei</i> strain HF.A00232, 5x10 <sup>9</sup> cfu, 12 weeks                                        | The authors reported that: "No serious adverse events were recorded in either group. The vital signs and physical examination of all systems revealed no differences between these two groups."                                                                                            |
| Lundelin et al. 2017        | Long-term safety of perinatal probiotic intervention.   | Prospective, randomized, double-blind, placebo-controlled with long-term follow-up | 303 mother-infant pairs                                                                       | <i>L. paracasei</i> ssp. <i>paracasei</i> strain ST11                                                                              | The authors reported that: "We found no differences in growth or non-communicable disease prevalence between children receiving perinatally probiotics or placebo" and concluded: "Perinatal probiotic administration is safe in long-term follow-up."                                     |
| Maretti and Cavallini 2017  | Treatment of idiopathic oligoastheno-teratospermia.     | Prospective, randomized, double-blind, placebo-controlled                          | 41 patients with idiopathic oligoastheno-teratospermia                                        | <i>L. paracasei</i> ssp. <i>paracasei</i> strain B21060, 5x10 <sup>9</sup> cfu, 6 months                                           | The authors concluded that: "There were no side effects in either group. These data showed that [the probiotic] constitutes a safe therapy."                                                                                                                                               |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference               | Objective                                                                         | Study Design                                              | Subjects                                                                                    | Strain, Dose, and Duration                                                                        | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                  |
|-------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------|---------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Martarelli et al. 2011  | Effect of probiotics on oxidant and antioxidant parameters in plasma of athletes. | Prospective, randomized, open-label                       | 24 healthy athletes                                                                         | <i>L. paracasei</i> ssp. <i>paracasei</i> strain IMC 502, $10^9$ cfu, 4 weeks                     | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                        |
| Nembrini et al. 2015    | Modulation of allergic rhinitis.                                                  | Prospective, randomized, double-blind, placebo-controlled | 131 allergic rhinitis patients (64M, 67F) aged 18-65 years                                  | <i>L. paracasei</i> ssp. <i>paracasei</i> strain NCC 2461, $5 \times 10^9$ cfu, 8 weeks           | The authors reported that: "Subjects receiving NCC 2461 reported fewer episodes of nasopharyngitis, however the frequency of minor adverse events was not a trial outcome and was therefore not considered in the statistical analysis." They concluded that: "Probiotic administration was considered safe as no serious adverse event was recorded during the trial." |
| Ouwehand et al. 2014    | Reduce incidence & severity of diarrhea in hospital.                              | Prospective, randomized, double-blind, multiple-dose      | 503 patients in hospital                                                                    | <i>L. paracasei</i> ssp. <i>paracasei</i> strain HN019, $1.7 \times 10^{10}$ cfu, approx. 10 days | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                        |
| Passariello et al. 2012 | Treatment of acute diarrhea in children.                                          | Prospective, randomized, double-blind, placebo-controlled | 107 children (58M,49F) aged 17-24 months with acute diarrhea (mean age = 20.5 months)       | <i>L. paracasei</i> ssp. <i>paracasei</i> strain B21060, $5 \times 10^9$ cfu, 5 days              | The authors reported that: "The rate of patients requiring hospitalisation because of worsening of symptoms was slightly but not significantly higher in placebo group. No adverse event was observed in the two groups."                                                                                                                                               |
| Perrin et al. 2014      | Modulation of allergic rhinitis.                                                  | Prospective, randomized, double-blind, cross-over         | 31 allergic rhinitis patients (21M, 10F) aged 20-35 years (mean age = $26.8 \pm 3.7$ years) | <i>L. paracasei</i> ssp. <i>paracasei</i> strain NCC2461, $10^{10}$ cfu, 4 weeks                  | The authors reported that: "There was no noticeable clinical issue and no formulation-related adverse event during the study." They concluded that: "This suggests that both probiotic treatments are safe and do not affect the immune system steady state."                                                                                                           |
| Pino et al. 2017        | Test viability after passing GI tract.                                            | Prospective open-label study                              | Apparently healthy adult volunteers                                                         | <i>L. paracasei</i> ssp. <i>paracasei</i> strain N24, 15 days                                     | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                        |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference              | Objective                                                       | Study Design                                                            | Subjects                                                            | Strain, Dose, and Duration                                                                 | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------|-----------------------------------------------------------------|-------------------------------------------------------------------------|---------------------------------------------------------------------|--------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Plaza-Diaz et al. 2013 | Safety and immune-modulatory effects of probiotics.             | Prospective, randomized, double-blind, placebo-controlled, multi-center | 100 healthy adults (46M, 54F) with mean age = 28.6 years            | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CNCM I-4034, $9 \times 10^9$ cfu, 30 days | The authors reported: "All symptom scores were less than 2, and there was no significant difference between the control group and the probiotic-treated group. The median score of the daily recorded GI symptoms of acid regurgitation, nausea, vomiting, abdominal distension, and eructation did not change during the probiotic supplementation (intervention) and subsequent follow-up period. Additionally, the stool consistency and defecation frequency did not change during the supplementation period and the subsequent follow-up period in the probiotic and placebo groups. Therefore, no serious adverse events occurred during the supplementation period in any of the groups, which shows that the differences between the probiotic and placebo groups were not significant for any of the reported symptoms. Likewise, no difference between placebo and probiotic groups occurred in any of the haematological parameters." They concluded: "These results demonstrate that the consumption of these three bacterial strains was safe." |
| Pu et al. 2017         | Probiotics to protect older people from respiratory infections. | Prospective, randomized, placebo-controlled, open-label                 | 205 healthy adults (59M, 146F) with mean age = $58.5 \pm 8.3$ years | <i>L. paracasei</i> ssp. <i>paracasei</i> strain N1115, $10^{10}$ cfu, 12 weeks            | The authors reported that: "No significant difference in the distribution of subjects leaving the study was observed. Furthermore, no adverse events associated with the consumption of the tested yogurt were reported."                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Rask et al. 2013       | Effect of probiotics on cell-mediated immunity.                 | Prospective, randomized, open-label                                     | 57 apparently healthy adults (20M, 37F) aged 18-55 (20M, 37F)       | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 8700:2, $10^{10}$ cfu, 5 weeks            | The authors reported that: "Only mild adverse gastrointestinal side effects were reported following intake of study products and there was no apparent relation between symptoms and treatment group."                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Rautava et al. 2012    | Maternal probiotics to reduce risk of eczema in the infant.     | Prospective, randomized, double-blind, placebo-controlled               | 241 mother-infant pairs; mother with allergic disease               | <i>L. paracasei</i> ssp. <i>paracasei</i> strain ST11, 4 months                            | The authors reported that: "No adverse effects were related to the use of probiotics," and concluded that: "Prevention regimen with specific probiotics administered to the pregnant and breast-feeding mother, that is, prenatally and postnatally, is safe and effective in reducing the risk of eczema in infants with allergic mothers positive for skin prick test."                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference              | Objective                                                 | Study Design                                                          | Subjects                                                                                 | Strain, Dose, and Duration                                                                 | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|------------------------|-----------------------------------------------------------|-----------------------------------------------------------------------|------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reygagne et al. 2017   | Effect of probiotics on severe dandruff.                  | Prospective, randomized, double-blind, placebo-controlled             | 60 men aged 18-60 years with severe dandruff                                             | <i>L. paracasei</i> ssp. <i>paracasei</i> strain ST11, $10^9$ cfu, 56 days                 | The authors stated that "No adverse events were reported" and concluded that "Regular intake of ST11 over 56 days is safe."                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Riezzo et al. 2012     | Effect of probiotics on functional constipation.          | Prospective, randomized, double-blind, placebo-controlled, cross-over | 20 functional constipation patients (3M, 17F), mean age = $38.8 \pm 14.4$ years          | <i>L. paracasei</i> ssp. <i>paracasei</i> strain IMPC 2.1, $2 \times 10^{10}$ cfu, 15 days | The authors reported that: "No adverse events were noted during the study period" and concluded that use of probiotics is "a useful and safe tool for managing constipation."                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Ritthagol et al. 2014  | Probiotics on mutans streptococci in cleft lip patients.  | Prospective, randomized, double-blind, placebo-controlled             | 30 patients aged $19.2 \pm 3.7$ years with cleft lip and palate                          | <i>L. paracasei</i> ssp. <i>paracasei</i> strain SD1, 4 weeks                              | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Rizzardini et al. 2012 | Immune benefits of probiotics.                            | Prospective, randomized, double-blind, placebo-controlled             | 211 healthy adults (93M, 118 F) aged $33.2 \pm 13.1$ years                               | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 431, $10^9$ cfu, 6 weeks                  | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Roessler et al. 2012   | Effects of probiotics in patients with atopic dermatitis. | Prospective, randomized, double-blind, placebo-controlled, crossover  | 15 apparently healthy adults & 15 atopic dermatitis patients                             | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC-37, 8 weeks                           | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Saito et al. 2017      | Safety of extreme intake of strain K71.                   | Open-label safety study                                               | 10 apparently healthy adults (5/sex) aged 20-64 years (mean age = $41.6 \pm 14.0$ years) | <i>L. paracasei</i> ssp. <i>paracasei</i> strain K71, $10^{12}$ cfu, 4 weeks               | The authors reported: "During this study, the following adverse events were observed: fullness feeling (n=1), loose stool (n=1), diarrhea (n=1), and abdominal pain loose stool (n=1). Each of these adverse events was mild and judged by the principal investigator as being unrelated to test food consumption. There were several slight but significant changes (from pretrial values) in parameters in tests for hematology and blood biochemistry, whereas the principal investigator judged that none of these deviations was of any clinical significance." They concluded that "doses of $10^{12}$ bacteria daily for 4 weeks were shown to be safe." |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference                 | Objective                                                                    | Study Design                                                         | Subjects                                                                            | Strain, Dose, and Duration                                                                                      | Safety-Related Results                                                                                                                                                                                                                                                                         |
|---------------------------|------------------------------------------------------------------------------|----------------------------------------------------------------------|-------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Simakachom et al. 2011    | Tolerance and safety of synbiotic formula in critically ill children.        | Prospective, randomized, double-blind, placebo-controlled            | 94 patients under mechanical ventilation aged 1-3 years                             | <i>L. paracasei</i> ssp. <i>paracasei</i> strain NCC 2461, 14 days                                              | The authors reported that “Abdominal distention, vomiting, and stool frequency were not affected by the supplementation with pre- and probiotics” and concluded that : “The enteral formula supplemented with synbiotics was well tolerated by children in intensive care units; it was safe.” |
| Teanpaisan and Piwat 2014 | Effect of probiotics on mutans streptococci.                                 | Prospective, randomized, double-blind, placebo-controlled            | 40 apparently healthy young adults                                                  | <i>L. paracasei</i> ssp. <i>paracasei</i> strain SD1, 4 weeks                                                   | No adverse events were reported.                                                                                                                                                                                                                                                               |
| Trautvetter et al. 2012   | Effect of probiotics on colonic lactobacilli & cholesterol metabolism.       | Prospective, randomized, double-blind, placebo-controlled, crossover | 32 apparently healthy adults                                                        | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC-37, $10^{10}$ cfu, 4 weeks                                 | No adverse events were reported.                                                                                                                                                                                                                                                               |
| Valerio et al. 2011       | Effect of probiotics on human biochemistry.                                  | Prospective, randomized, double-blind, placebo-controlled, crossover | 20 healthy adults (3M, 17F) aged $37.8 \pm 13.9$ years                              | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LMGP22043, $2 \times 10^{10}$ cfu, 15 days                     | No adverse events were reported.                                                                                                                                                                                                                                                               |
| Valsecchi et al. 2014     | Effect of probiotics on secretory IgA during antibiotic therapy in children. | Prospective, randomized, double-blind, placebo-controlled            | 77 pediatric patients with recurrent airway infections receiving antibiotic therapy | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CFL-431 (& <i>B. lactis</i> & <i>S. thermophilus</i> ), 7 days | No adverse events were reported.                                                                                                                                                                                                                                                               |
| Verdenelli et al. 2011    | Influence of probiotics on adult bowel habits.                               | Prospective, randomized, double-blind, placebo-controlled            | 50 apparently healthy adults (23M, 27F) aged 23-65 years                            | <i>L. paracasei</i> ssp. <i>paracasei</i> strain IMC 502, $2 \times 10^9$ cfu, 12 weeks                         | The authors reported that “No significant differences were detected with respect to constipation and flatulence by both groups. The probiotic-enriched foods consumption was very well tolerated, and no side effects were experienced.”                                                       |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference              | Objective                                         | Study Design                                                          | Subjects                                                                | Strain, Dose, and Duration                                                                                                         | Safety-Related Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------------|---------------------------------------------------|-----------------------------------------------------------------------|-------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Vicariotto 2014        | Effect of probiotics on urinary tract infections. | Open-label                                                            | 33 premenopausal women with acute cystitis                              | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LPC09, 60 days                                                                    | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Vlieger et al. 2009    | Tolerance and safety of probiotics.               | Prospective, randomized, double-blind, placebo-controlled             | 126 healthy term neonates <7 days of age                                | <i>L. paracasei</i> ssp. <i>paracasei</i> strain CRL-431, 10 <sup>7</sup> cfu, 3 months (& <i>B. animalis</i> ssp. <i>lactis</i> ) | <p>The authors stated that: “No serious adverse events were reported that could be related to the study formula. Parents were asked if they had noticed any symptoms that could have been caused by the study feed. Fewer infants in the probiotics group had developed a rash in the first 3 months. No differences were seen in other adverse effects between the two groups in both the first and second trimester.”</p> <p>They concluded: “Normal growth occurred in all infants and no statistically significant differences were detected between the probiotics group and the control group for gain in weight, length and head circumference. Infants in the probiotics group produced softer and more frequent stools during the first 3 months of life. No differences were found in crying and sleeping hours, number of parent-diagnosed infections, antibiotic use, visits to the general practitioner and number of adverse events. The use of a prebiotic-containing starter formula supplemented with <i>L. paracasei</i> ssp. <i>paracasei</i> and <i>B. animalis</i> ssp. <i>lactis</i> in early infancy is safe, well tolerated and has no adverse effects on growth and infant behaviour.”</p> |
| Wang et al. 2004       | Effect of probiotics on allergic rhinitis.        | Prospective, randomized, double-blind, placebo-controlled             | 80 allergic rhinitis patients (41M, 39F) with mean age = 15.4±1.8 years | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 33, 2x10 <sup>9</sup> cfu, 30 days                                                | The authors reported that no patients left the study due to adverse events, and “Subjects reported no severe adverse effects such as fever, abdominal pain, or diarrhea.” They concluded that: “The results suggest that ingestion of LP-33-fortified fermented milk for 30 days can effectively and safely improve the quality of life of patients with allergic rhinitis.”                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Wassenberg et al. 2011 | Effect of probiotics on allergic rhinitis.        | Prospective, randomized, double-blind, placebo-controlled, cross-over | 31 patients with allergic rhinitis                                      | <i>L. paracasei</i> ssp. <i>paracasei</i> strain ST11, 4 weeks                                                                     | No adverse events were reported.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |

**Table 6. Human Studies in *L. paracasei* ssp. *paracasei* Strains Other Than F-19.**

| Reference              | Objective                                                   | Study Design                                                              | Subjects                                                   | Strain, Dose, and Duration                                                                                                                               | Safety-Related Results                                                                                                                                                                                        |
|------------------------|-------------------------------------------------------------|---------------------------------------------------------------------------|------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Wattanarat et al. 2015 | Effect of probiotics on salivary HNP1-3 levels in children. | Prospective, randomized, double-blind, placebo-controlled, parallel-group | 60 apparently healthy children (26M, 34F) aged 13-15 years | <i>L. paracasei</i> ssp. <i>paracasei</i> strain SD1, $3.75 \times 10^9$ cfu, 6 months                                                                   | The authors reported that: "No adverse side effects from probiotics or milk powder intake in this cohort were reported."                                                                                      |
| West et al. 2012       | Effect of probiotics on immune system.                      | Prospective, randomized, double-blind, placebo-controlled, parallel-group | 22 healthy males with mean age = $32.9 \pm 4.5$ years      | <i>L. paracasei</i> ssp. <i>paracasei</i> strain 431, $4.6 \times 10^8$ cfu, 21 days (& <i>B. lactis</i> , <i>L. acidophilus</i> & <i>L. rhamnosus</i> ) | The authors reported that: "There were five episodes of mild GI symptoms that included flatulence and stomach rumbles in both groups during supplementation. Both supplements were otherwise well tolerated." |
| Zhang et al. 2013b     | Effect of probiotics on colonic microbiota                  | Prospective, randomized, double-blind, placebo-controlled, cross-over     | 52 apparently healthy adults                               | <i>L. paracasei</i> ssp. <i>paracasei</i> strain LC01, $10^{10}$ cfu, 4 weeks                                                                            | No adverse events were reported.                                                                                                                                                                              |

## 6.6. Toxicity Studies of *L. paracasei* ssp. *paracasei*

Oral toxicity studies were conducted on 4 strains of *L. paracasei* ssp. *paracasei*: strain Lpc-37, LC-01, NTU 101, and GM080 (Morovic et al. 2017; Zhang et al. 2013a; Tseng et al. 2015, and Jia et al. 2011). The first was an acute study, the next two were subacute, and the fourth was subchronic.

Morovic et al. (2017) investigated the safety of HOWARU® Restore, a blend of *Lactobacillus acidophilus* NCFM, *Lactobacillus paracasei* Lpc-37, *Bifidobacterium animalis* subsp. *lactis* Bl-04 and *B. lactis* Bi-07, as well as the individual strains. Acute oral toxicity testing was done in 10-week-old female Crl:CD(SD) rats that received single gavage doses of 5000 mg/kg bw, equivalent to  $2.64 \times 10^{12}$  cfu/kg bw for HOWARU® Restore and  $3.35 \times 10^{12}$  CFU/kg bw for *L. paracasei* Lpc-37 tested individually. Rats were observed for 14 days and weighed on days -1, 1, 8, and 15 prior to euthanasia. Necropsy was performed to detect grossly observable evidence of organ or tissue damage.

There were no incidents of mortality, clinical abnormalities, or overall (test day 1–15) body weight losses reported in any animal administered either HOWARU® Restore or *L. paracasei* Lpc-37 and no gross findings detected at necropsy suggestive of acute toxicity.

In an assessment of the safety of *L. paracasei* ssp. *paracasei* strain LC-01 (Zhang et al. 2013a), 40 7-9-week-old female Balb/C mice were assigned to 4 groups (10 mice/group) to receive daily gavage of 0,  $10^8$ ,  $10^9$ , or  $10^{10}$  cfu of strain LC-01 for 28 days. Mice were observed daily; feed intake and body weight were recorded weekly. On day 29, blood was collected, organs and tissues were observed for gross abnormalities, and liver, spleen, and kidney were excised. Blood was analyzed for red blood cell and platelet counts, hemoglobin, hematocrit, mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, glutamic-oxalacetic transaminase, glutamic-pyruvic transaminase activity, total protein, albumin, glucose, cholesterol, and bacterial translocation.

No noticeable activity or behavioral changes were reported in the mice during the in-life portion of the study, and no treatment-related illness or death were reported. There were no significant differences in feed intake, bodyweight gain, or organ weights between the control and treatment groups, and the hematological and biochemical parameters did not show statistically significant differences between control and strain-treated groups. The authors stated that, “Our findings suggest that LC-01 do not have adverse effect on murine haematology or blood biochemistry.”

Additionally, no bacteremia was reported in any of the groups, and there were no significant differences in the incidence of translocation to the liver, spleen, or kidney between the control and treated groups at any of the tested doses. The authors concluded that, “Therefore, it can be concluded that LC-01 is likely to be non-pathogenic and safe for human consumption.”

After completing *in vitro* and *in vivo* genotoxicity studies (reverse bacterial mutation, micronucleus assay, and mammalian chromosomal aberration test) of *L. paracasei* ssp. *paracasei* strain NTU 101, finding no evidence of mutagenicity, Tseng et al. (2015) performed a subacute oral toxicity study in Wistar rats. Eighty rats, 40 of each sex, weighing 170-200 g were assigned to 4 groups (10 rats/sex/group) to receive 0, 300, 1500, or 5000 mg/kg bw/day of the bacterial powder for 28 days; these doses provided 0,  $1.8 \times 10^9$ ,  $4.5 \times 10^9$ , and  $1.5 \times 10^{10}$  cfu of strain NTU 101/kg bw/day. Animals were observed and weighed daily and feed intake was measured semiweekly. On the day of euthanasia, blood was taken from the abdominal aorta and analyzed for hematology (hemoglobin, hematocrit, lymphocytes, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, mean corpuscular volume, mean platelet volume, platelet distribution width, platelet large-cell ratio, platelet count, red blood cell count, red blood cell distribution width, and white blood cell count) and clinical chemistry parameters (alanine aminotransferase, aspartate aminotransferase, blood urea nitrogen, calcium, cholesterol, chloride, creatinine, glucose,

potassium, magnesium, sodium, inorganic phosphorus, and triacylglycerol). All organs and tissues were examined macroscopically (and microscopically for rats in the high-dose and control groups) and weights were obtained for the liver, kidney, heart, spleen, lung, adrenal gland, epididymis, testis, uterus, and ovary.

During the study period, no clinical signs or deaths were reported in either the control or the treatment groups. The animals showed no significant differences in feed consumption, body weight, hematological parameters, absolute or relative organ weights, or histology between the control and treatment groups. Although there were some significant differences between treated and control rats in potassium, aspartate aminotransferase, and triacylglycerol levels, all values were within the normal range. The authors concluded that:

“This study demonstrates that Vigiis 101 has no mutagenic/genotoxic effects based on the results of the Ames test, the *in vitro* chromosomal aberration test, or the *in vivo* micronucleus assay; there was no evidence of toxicity in the 28-day oral toxicity assay at 5000 mg/kg/day in rats. Taken together, these results support the safety of Vigiis 101 made from *L. paracasei* subsp. *paracasei* NTU 101.”

The safety of a fourth strain, *L. paracasei* ssp. *paracasei* strain GM080, was assessed in a sub-chronic study (Jia et al. 2011) in Sprague-Dawley rats weighing  $66.6 \pm 6.5$  g (males) and  $64.2 \pm 7.1$  g (females). Eighty rats were divided into 4 groups of 10 rats/sex/group to receive by gavage 0, 1.25, 2.5, or 5.0 g strain GM080/kg bw/day for 90 days; the doses were equivalent to 0,  $2.5 \times 10^9$ ,  $5 \times 10^9$ , and  $10^{10}$  cfu/kg bw/day. Rats were individually caged and observed twice daily; feed intake and weight were measured weekly. Blood was tested at day 46 and day 91 for hematology (red blood cell count, hemoglobin, platelet count, white blood cell count, and differential blood cell count) and clinical chemistry (alanine aminotransferase, aspartate aminotransferase, total protein, albumin, glucose, blood urea nitrogen, creatinine, cholesterol, triglyceride and alkaline phosphatase). After euthanasia, selected organs (heart, kidneys, liver, spleen, testes and thymus) were excised and weighed and macro- and microscopic histological examinations were performed on organs and tissues from rats in the control and high-dose groups (cecum, colon, duodenum, esophagus, femur with bone marrow, ileum, jejunum, lacrimal gland, lung, lymph node, mammary gland, nasal turbinates, pancreas, pituitary gland, prostate, rectum, salivary gland, sciatic nerve, seminal vesicles, skeletal muscle (thigh), skin, spinal cord, sternum with bone marrow, trachea, urinary bladder and vagina).

No mortality, adverse clinical reactions, or differences in feed intake or bodyweight gain were reported. There were some sporadic, statistically significant changes in some hematology and clinical chemistry parameters, but none were consistent and there was no dose-response relationship. There were no statistically significant differences in absolute or relative organ between treatment groups and control groups and no macroscopic pathology findings. Slightly sporadic focal necrosis in liver was found in one animal in the control group; focal necrosis was reported in the heart in some rats (1/20 animal in the control group and 3/20 animals in the high-dose group).

The authors concluded that, “the results of the present study demonstrate that *L. paracasei* GM080 are non-toxic up to a level of 5.0 g/kg body weight, when given orally. The no-observed-adverse-effect-level (NOAEL) for *L. paracasei* GM080 was 5.0 g/kg body weight (approximately equivalent to  $1 \times 10^{10}$  cfu/kg bw) in male and female rats, the highest dose tested.”

In summary, all toxicity testing of 4 different strains of *L. paracasei* ssp. *paracasei* has found no evidence that would suggest any adverse reactions in mice or rats.

## 6.6. Decision-Tree Analysis

The decision tree of Pariza et al. (2015) was utilized to assess the evidence regarding the safety of the intended use of *L. paracasei* ssp. *paracasei* strain F-19. Significant questions follow:

1. Has the strain been characterized for the purpose of assigning an unambiguous genus and species name using currently accepted methodology? **YES**
2. Has the strain genome been sequenced? **YES**
3. Is the strain genome free of genetic elements encoding virulence factors and/or toxins associated with pathogenicity? **YES**
4. Is the strain genome free of functional and transferable antibiotic resistance gene DNA? **YES**
5. Does the strain produce antimicrobial substances? **NO**
6. Has the strain been genetically modified using rDNA techniques? **NO**
7. Was the strain isolated from a food that has a history of safe consumption for which the species, to which the strain belongs, is a substantial and characterizing component (not simply an 'incidental isolate')? **NO**—Isolated from a healthy human colon
8. Does the strain induce undesirable physiological effects in appropriately designed safety evaluation studies? **NO**

The outcome of this decision-tree analysis is that “the strain is deemed to be safe for use in the manufacture of food, probiotics, and dietary supplements for human consumption” (Pariza et al., 2015).

## 6.7. Evaluations by Authoritative Bodies

Noting that a wide variety of microbial species is used in food, some with a long history of apparent safe use, and facing the need to set priorities for risk assessment, the European Food Safety Authority (EFSA) proposed a system referred to as “Qualified Presumption of Safety” (QPS; EFSA 2007a, 2007b). This system proposed basing the safety assessment of a defined taxonomic group (e.g., a genus or a species) on 4 pillars: established identity, body of knowledge, possible pathogenicity, and end use. If the taxonomic group did not raise safety concerns or, if safety concerns existed but could be defined and excluded, the grouping could be granted QPS status. Thereafter, “any strain of microorganism the identity of which could be unambiguously established and assigned to a QPS group would be freed from the need for further safety assessment other than satisfying any qualifications specified” (EFSA 2007a, p1).

EFSA’s Scientific Committee was asked to recommend organisms regarded as suitable for QPS status. The list of such organisms proposed by the Committee included *L. paracasei*. In listing *L. paracasei* and other species of *Lactobacillus* as suitable for QPS status, the Committee stated, “Where QPS status is proposed, the Scientific Committee is satisfied that the body of knowledge available is sufficient to provide adequate assurance that any potential to produce adverse effects in humans, livestock or the wider environment is understood and capable of exclusion” (EFSA 2007a, p8) and that the recommendations are “based on a thorough review of the available scientific literature and the knowledge and experience of the scientists involved” (EFSA 2007a, p8).

EFSA has issued annual updates of the QPS status of bacterial strains from 2008 through 2016; no need for either review or change of the QPS status of *L. paracasei* was reported (EFSA 2008b, 2009, 2010, 2011, 2012b, 2013, 2014, 2015a, 2015b, 2016, 2017, 2018).

The Danish Veterinary and Food Administration, having previously approved *L. paracasei* ssp. *paracasei* strain F-19 for use in dietary supplements and several types of foods, was petitioned

to extend this approval to use in infant formula, transition foods, and other food types. This approval was granted in a letter dated 1 December 2009 (DVFA 2009).

## **6.8. Safety Assessment and GRAS Determination**

This section presents an assessment that demonstrates that the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is safe and is GRAS based on scientific procedures.

This safety assessment and GRAS determination entail two steps. In the first step, the safety of the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is demonstrated. Safety is established by demonstrating a reasonable certainty that the exposure of consumers to *L. paracasei* ssp. *paracasei* strain F-19 under its intended conditions of use is not harmful. In the second step, the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is determined to be GRAS by demonstrating that the safety of this product under its intended conditions of use is generally recognized among qualified scientific experts and is based on publicly available and accepted information.

The regulatory framework for establishing whether the intended use of a substance (or organism) is GRAS, in accordance with Section 201(s) of the Federal Food Drug and Cosmetic Act, is set forth under 21 CFR §170.30. This regulation states that general recognition of safety may be based on the view of experts qualified by scientific training and experience to evaluate the safety of substances directly or indirectly added to food. A GRAS determination may be made either: 1) through scientific procedures under §170.30(b); or 2) through experience based on common use in food, in the case of a substance used in food prior to January 1, 1958, under §170.30(c). This GRAS determination employs scientific procedures established under §170.30(b).

A scientific procedures GRAS determination requires the same quantity and quality of scientific evidence as is needed to obtain approval of the substance as a food additive. In addition to requiring scientific evidence of safety, a GRAS determination also requires that this scientific evidence of safety be generally known and accepted among qualified scientific experts. This “common knowledge” element of a GRAS determination consists of two components:

1. Data and information relied upon to establish the scientific element of safety must be generally available; and
2. There must be a basis to conclude that there is a consensus among qualified experts about the safety of the substance for its intended use.

The criteria outlined above for a scientific-procedures GRAS determination are applied below in an analysis of whether the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is safe and is GRAS.

### **6.8.1. Evidence of Safety**

The body of evidence supporting the safety of oral administration of *Lactobacillus* strains in general, and *L. paracasei* ssp. *paracasei* strain F-19 in particular, is large and convincing. Numerous commentators and authoritative bodies such as EFSA have noted the safe history of human ingestion of *L. paracasei* ssp. *paracasei* strains over many years. *L. paracasei* ssp. *paracasei* strain F-19 produces no deleterious metabolites and is not destructive of mucin. Any effects that this probiotic microorganism has on intestinal permeability appear to be beneficial in strengthening barrier function. While it is theoretically possible for biogenic amines to be produced as a result of fermentation of dairy products by lactobacilli, this phenomenon has not been reported. When *Lactobacillus* strains are ingested as probiotics, they produce lactic acid, lowering the intestinal pH and reducing the opportunity for production of harmful biogenic amines by putrefactive bacteria.

Lactobacilli are not regarded as pathogens, although some strains are capable of opportunistic infection in extremely favorable circumstances invariably involving severe underlying disease states and most often also involving a facilitated pathway such as surgical intervention or the presence of central lines. Documented cases of *Lactobacillus* bacteremia are so rare, in comparison to the widespread use of *Lactobacillus* strains in the environment, in food production, and in probiotic applications, that the participants in the 2007 EU-PROSAFE project (Vankerckhoven et al. 2008) suggested that “they are more medical exceptions, or even curiosities, than a genuine public health issue.”

Consumption of live lactic acid bacteria included in lactic-acid-fermented foods has been a regular part of the food intake of humans for hundreds of years and individuals consuming lactic-acid-fermented products of dairy origin also consume large amounts of *L. paracasei* ssp. *paracasei*.

The safe history of human exposure to *L. paracasei* ssp. *paracasei* strains is strongly supported by a large body of published research. In addition to *in vitro* work, the published literature includes 4 oral toxicity studies in mice and rats and more than 50 studies in human infants, children, and adults enrolling more than 7,000 individuals. Participants in these studies received the probiotic at daily levels up to  $10^{12}$  cfu/day for as long as 9 months with no reported adverse events associated with the probiotic intervention.

The strain *L. paracasei* ssp. *paracasei* strain F-19 originates from human intestinal mucosa (Ljungh et al. 2002) and it is sold in food products and as a dietary supplement in Europe. Since 2001, products containing *L. paracasei* ssp. *paracasei* strain F-19 have been widely consumed without any reported adverse events.

The genome of *L. paracasei* ssp. *paracasei* strain F-19 was sequenced and genes were annotated and compared with databases of antibiotic resistance genes and virulence factors; additionally, terms were searched for possible associations with antibiotic resistance or virulence. No findings were suggestive of potential risk to consumers. The absence of genes encoding antibiotic resistance was confirmed by phenotypic testing for such resistance in which no resistance was found other than intrinsic resistance to vancomycin found in many *Lactobacillus* strains. Additional phenotypic testing demonstrated that the strain does not produce biogenic amines and only trace levels of D-lactate.

*L. paracasei* ssp. *paracasei* strain F-19 has been studied in 3 trials with infants; the highest test dose was  $6 \times 10^9$  cfu/day for 21 days in a study with low-birthweight preterm infants. Two studies in children and 27 in adults included doses as high as  $10^{11}$  cfu/day for up to 9 months. No adverse events attributable to the probiotic were reported in any of these studies.

The available evidence demonstrates that there is no reason to suspect harm to individuals from the intended use of *L. paracasei* ssp. *paracasei* strain F-19.

#### **6.8.2. Conclusion of the Expert Panel**

The intended use of *L. paracasei* ssp. *paracasei* strain F-19 has been determined to be safe through scientific procedures set forth under 21 CFR §170.30(b). This safety was shown by establishing the identity and probiotic characteristics of the strain, demonstrating its freedom from pathogenic or other risk factors using the decision-tree analysis, and concluding that the expected exposure to *L. paracasei* ssp. *paracasei* strain F-19 is without significant risk of harm. Finally, because this safety assessment satisfies the common knowledge requirement of a GRAS determination, this intended use can be considered GRAS.

Determination of the safety and GRAS status of the intended use of *L. paracasei* ssp. *paracasei* strain F-19 has been made through the deliberations of an Expert Panel consisting of Joseph F. Borzelleca, Ph.D., Berthold V. Koletzko, M.D., Ph.D., and Michael W. Pariza, Ph.D., who

reviewed a monograph prepared by James T. Heimbach, Ph.D., and other information available to them. These individuals are qualified by scientific training and experience to evaluate the safety of food and food ingredients, including probiotic bacteria. They independently critically reviewed and evaluated the publicly available information and the potential human exposure to *L. paracasei* ssp. *paracasei* strain F-19 anticipated to result from its intended use, and individually and collectively determined that no evidence exists in the available information on *L. paracasei* ssp. *paracasei* strain F-19 that demonstrates, or suggests reasonable grounds to suspect, a hazard to consumers under the intended conditions of use of *L. paracasei* ssp. *paracasei* strain F-19.

It is the Expert Panel's opinion that other qualified scientists reviewing the same publicly available data and information would reach the same conclusion regarding the safety of the strain under its intended conditions of use. Therefore, the intended use of *L. paracasei* ssp. *paracasei* strain F-19 is GRAS by scientific procedures.

#### **6.9. Statement Regarding Information Inconsistent with GRAS**

I have reviewed the available data and information and am not aware of any data or information that are, or may appear to be, inconsistent with our conclusion of GRAS status of the intended use of *L. paracasei* ssp. *paracasei* strain F-19.

A rectangular area of the document has been redacted with a solid grey box. Below the box, there are faint, illegible handwritten marks in blue ink.

## Part 7: List of Supporting Data and Information

- Alander M, De Smet I, Nollet L, Verstraete W, von Wright A, Mattila-Sandholm T. 1999. The effect of probiotic strains on the microbiota of the simulator of the human intestinal microbial ecosystem (SHIME). *Int J Food Microbiol* 46:71-79.
- Allen SJ, Jordan S, Storey M, Thornton CA, Gravenor MB, Garaiova I, Plummer SF, Wang D, Morgan G. 2014. Probiotics in the prevention of eczema: a randomized controlled trial. *Arch Dis Child* 99:1014-1019.
- Andersen AD, Cilieborg MS, Lauridsen C, Morkbak AL, Sangild PT. 2017. Supplementation with *Lactobacillus paracasei* or *Pediococcus pentosaceus* does not prevent diarrhoea in neonatal pigs infected with *Escherichia coli* F18. *Br J Nutr* 118:109-120.
- Annibale B, Maconi G, Lahner E, De Giorgi F, Cuomo R. 2011. Efficacy of *Lactobacillus paracasei* subsp. *paracasei* F-19 on abdominal symptoms in patients with symptomatic uncomplicated diverticular disease: a pilot study. *Minerva Gastroenterol Dietol* 57:13-22.
- Aronsson L, Huang Y, Parini P, Korach-Andre M, Hakansson J, Gustafsson J. A, Petersson S, Arulampalam V, Rafter J. 2010. Decreased fat storage by *Lactobacillus paracasei* is associated with increased levels of angiopoietin-like 4 protein (ANGPTL4). *PLoS One*. 5: e13087.
- Balzaretti S, Taverniti V, Rondini G, Marcollegio G, Minuzzo M, Remagni MC, Fiore W, Arioli S, Guglielmetti S. 2015. The vaginal isolate *Lactobacillus paracasei* LPC- S01 (DSM26760) is suitable for oral administration. *Front Microbiol* 6:952-964.
- Barker AK, Duster M, Valentine S, Hess T, Archbald-Pannone L, Guerrant R, Safdar N. 2017. A randomized controlled trial of probiotics for *Clostridium difficile* infection in adults (PICO). *J Antimicrob Chemother* 72:3177-3180.
- Begtrup LM, Schaffalitzky de Muckadell OB, Kjeldsen J, Christensen RdeP, Jarbol DE. 2013. Long-term treatment with probiotics in primary care patients with irritable bowel syndrome – a randomised, double-blind, placebo controlled trial. *Scand J Gastroenterol* 48:1127-1135.
- Bennedsen M, Stuer-Lauridsen B, Danielsen M, Johansen E. 2011. Screening for antimicrobial resistance genes and virulence factors via genome sequencing. *Appl Environ Microbiol* 77:2785–2787.
- Bjerg AT, Sorensen MB, Krych L, Hansen LH, Astrup A, Kristensen M, Nielsen DS. 2015a. The effect of *Lactobacillus paracasei* subsp. *paracasei* L. casei W8® on blood levels of triacylglycerol is independent of colonisation. *Benef Microbes* 6:263-269.
- Bjerg AT, Kristensen M, Ritz C, Stark KD, Holst JJ, Leser TD, Wellejus A, Astrup A. 2015b. Four weeks supplementation with *Lactobacillus paracasei* subsp. *paracasei* L. casei W8® shows modest effect on triacylglycerol in young healthy adults. *Benef Microbes* 6:29-39.
- Bjorneholm S, Eklow A, Saarela M, Matto J. 2002. Enumeration and identification of *Lactobacillus paracasei* subsp. *paracasei* F-19. *Microb Ecol Health Dis* 14:7-13
- Brahe LK, Le Chatelier E, Prifti E, Pons N, Kennedy S, Blaedel T, Hakansson J, Dalsgaard TK, Hansen T, Pedersen O, Astrup A, Ehrlich SD, Larsen LH. 2015. Dietary modulation of the gut microbiota – a randomised controlled trial in obese postmenopausal women. *Br J Nutr* 114:406-417.
- Charteris W, Kelly P, Morelli L, Collins K. 1998a. Antibiotic susceptibility of potentially probiotic *Lactobacillus* species. *J Food Prot* 61:1636-1643.

- Charteris W, Kelly P, Morelli L, Collins K. 1998b. Development and application of an *in vitro* methodology to determine the transit tolerance of potentially probiotic *Lactobacillus* and *Bifidobacterium* species in the upper gastrointestinal tract. *J Appl Microbiol* 84:759-768.
- Charteris W, Kelly P, Morelli L, Collins K. 2001. Quality control *Lactobacillus* strains for use with the API 50CH and API ZYM systems at 37°C. *J Basic Microbiol* 41:241-251.
- Chen L, Zheng D, Liu B, Yang J, Jin Q. 2016. VFDB 2016: hierarchical and refined dataset for big data analysis - 10 years on. *Nucleic Acids Res* 44:D694–D697.
- Chorell E, Videhult FK, Hernell O, Antti H, West CE. 2013. Impact of probiotic feeding during weaning on the serum lipid profile and plasma metabolome in infants. *Br J Nutr* 110:116-126.
- Chouraqui JP, Grathwohl D, Labaune JM, Hascoet JM, de Montgolfier I, Leclaire M, Giarre M, Steenhout P. 2008. Assessment of the safety, tolerance, and protective effect against diarrhea of infant formulas containing mixtures of probiotics or probiotics and prebiotics in a randomized controlled trial. *Am J Clin Nutr* 87:1365-1373.
- Compare D, Rocco A, Sgamato C, Coccoli P, Campo SMA, Nazionale I, Larussa T, Luzzza F, Chiodini P, Nardone G. 2015. *Lactobacillus paracasei* F-19 versus placebo for the prevention of proton pump inhibitor-induced bowel symptoms: a randomized clinical trial. *Digest Liver Dis* 47:273-279.
- Corsello G, Carta M, Marinello R, Picca M, De Marco G, Micillo M, Ferrara D, Vigneri P, Cecere G, Ferri P, Roggero P, Bedogni G, Mosca F, Paparo L, Nocerino R, Canani RB. 2017. Preventive effect of cow's milk fermented with *Lactobacillus paracasei* CBA L74 on common infectious diseases in children: a multicenter randomized controlled trial. *Nutrients* 9:669-683.
- Costa DJ, Marteau P, Amouyal M, Poulsen LK, Hamelmann E, Cazaubiel M, Housez B, Leuillet S, Stavnsbjerg M, Molimard P, Courau S, Bousquet J. 2014. Efficacy and safety of the probiotic *Lactobacillus paracasei* LP-33 in allergic rhinitis: a double-blind, randomized, placebo controlled trial (GA2LEN Study). *Eur J Clin Nutr* 68:602-607.
- Crittenden R, Saarela M, Matto J, Ouwehand AC, Salminen S, Peltto L, Vaughan EE, de Vos WM, von Wright A, Fonden R, Mattila-Sandholm T. 2002. *Lactobacillus paracasei* subsp. *paracasei* F-19: survival, ecology and safety in the human intestinal tract - a survey of feeding studies within the PROBDEMO project. *Microb Ecol Health Dis* 14:22-26.
- Danish Veterinary and Food Administration (DVFA). 2009. *Approval of extended application range of L. paracasei subsp. paracasei*. Letter to Chr. Hansen Holding A/S, 1 December.
- Delia A, Morgante G, Rago G, Musacchio MC, Petraglia F, De Leo V. 2006. Effectiveness of oral administration of *Lactobacillus paracasei* subsp. *paracasei* F-19 in association with vaginal suppositories of *Lactobacillus acidophilus* in the treatment of vaginosis and in the prevention of recurrent vaginitis. *Minerva Ginecol* 58:227-231.
- EFSA Panel on Additives and Products or Substances used in Animal Feed (FEEDAP). 2012. Guidance on the assessment of bacterial susceptibility to antimicrobials of human and veterinary importance. *EFSA J* 10:2740.
- European Food Safety Authority (EFSA). 2007a. Introduction of a qualified presumption of safety (QPS) approach for assessment of selected microorganisms referred to EFSA: opinion of the scientific committee. *EFSA J* 5:8-16.

- European Food Safety Authority (EFSA). 2007b. Introduction of a qualified presumption of safety (QPS) approach for assessment of selected microorganisms referred to EFSA: opinion of the scientific committee. Appendix A: Assessment of gram-positive non-sporulating bacteria. *EFSA J* 587.
- European Food Safety Authority (EFSA). 2008a. Update of the criteria used in the assessment of bacterial resistance to antibiotics of human or veterinary importance. Scientific opinion of the Panel on Additives and Products or Substances Used in Animal Feed. *EFSA J* 732:1-15.
- European Food Safety Authority (EFSA). 2008b. The maintenance of the list of QPS microorganisms intentionally added to food or feed. Scientific opinion of the Panel on Biological Hazards. *EFSA J* 923:1-48.
- European Food Safety Authority (EFSA). 2009. Scientific opinion on the maintenance of the list of QPS microorganisms intentionally added to food or feed (2009 update). *EFSA J* 7(12):1431.
- European Food Safety Authority (EFSA). 2010. Scientific opinion on the maintenance of the list of QPS microorganisms intentionally added to food or feed (2010 update). *EFSA J* 8(12):1944.
- European Food Safety Authority (EFSA). 2011. Scientific Opinion on the maintenance of the list of QPS biological agents intentionally added to food and feed (2011 update). *EFSA J* 9(12):2497-2578.
- European Food Safety Authority (EFSA). 2012a. Guidance on the assessment of bacterial susceptibility to antimicrobials of human and veterinary importance. *EFSA J* 10(6) 2740-2749.
- European Food Safety Authority (EFSA). 2012b. Scientific Opinion on the maintenance of the list of QPS biological agents intentionally added to food and feed (2012 update). *EFSA J* 10(12):3020-3103.
- European Food Safety Authority (EFSA). 2013. Scientific Opinion on the maintenance of the list of QPS biological agents intentionally added to food and feed (2013 update). *EFSA J* 11(11):3449-3554.
- European Food Safety Authority (EFSA). 2014. Statement on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 1: Suitability of taxonomic units notified to EFSA until October 2014. *EFSA J* 12(12):3938-3978.
- European Food Safety Authority (EFSA). 2015. Statement on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA. 2: Suitability of taxonomic units notified to EFSA until March 2015. *EFSA J* 13(6):4138-4166.
- European Food Safety Authority (EFSA). 2015. Statement on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 3: Suitability of taxonomic units notified to EFSA until September 2015. *EFSA J* 13(12):4331-4355.
- European Food Safety Authority (EFSA). 2016. Update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 4: suitability of taxonomic units notified to EFSA until March 2016. *EFSA J* 14(7):4522-4558.
- European Food Safety Authority (EFSA). 2017. Update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 6: suitability of taxonomic units notified to EFSA until March 2017. *EFSA J* 15(7):4884-4932.

- European Food Safety Authority (EFSA). 2018. Update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 7: suitability of taxonomic units notified to EFSA until September 2017. *EFSA J* 16(1):5131-5173.
- Ferrario C, Taverniti V, Milani C, Fiore W, Laureati M, De Noni I, Stuknyte M, Chouaia B, Riso P, Guglielmetti S. 2014. Modulation of fecal *Clostridiales* bacteria and butyrate by probiotic intervention with *Lactobacillus paracasei* DG varies among healthy adults. *J Nutr* 144:1787-1796.
- Foerst P, Reitmaier J, Kulozik U. 2010. <sup>1</sup>H NMR investigation on the role of sorbitol for the survival of *Lactobacillus paracasei* ssp. *paracasei* in vacuum-dried preparations. *J Appl Microbiol* 108:841-850.
- Fomon SJ. 1993. Energy intake by normal infants. In *Nutrition of Normal Infants*, p. 104–111. Baltimore, MD: Mosby.
- Gore C, Custovic A, Tannock GW, Munro K, Kerry G, Johnson K, Peterson C, Morris J, Chaloner C, Murray CS, Woodcock A. 2012. Treatment and secondary prevention effects of the probiotics *Lactobacillus paracasei* or *Bifidobacterium lactis* on early infant eczema: randomized controlled trial with follow-up until age 3 years. *Clin Exp Allergy* 42:112-122.
- Grossi E, Buresta R, Abbiati R, Cerutti R, Pro-DIA study group. 2010. Clinical trial on the efficacy of a new symbiotic formulation, Flortec, in patients with acute diarrhea: a multicenter, randomized study in primary care. *J Clin Gastroenterol* 44 Suppl 1:S35-41.
- Gueniche A, Philippe D, Bastien P, Reuteler G, Blum S, Castiel-Higounenc I, Breton L, Benyacoub J. 2014. Randomised double-blind placebo-controlled study of the effect of *Lactobacillus paracasei* NCC 2461 on skin reactivity. *Benef Microbes* 5:137-145.
- Hasslof P, Hedberg M, Twetman S, Stecksén-Blicks C. 2010. Growth inhibition of oral mutans streptococci and candida by commercial probiotic lactobacilli--an *in vitro* study. *BMC Oral Health* 2;10:18.
- Hasslof P, West CE, Videhult FK, Brandelius C, Stecksén-Blicks C. 2013. Early intervention with probiotic *Lactobacillus paracasei* F-19 has no long-term effect on caries experience. *Caries Res* 47:559-565.
- Hedberg M, Hasslof P, Sjöström I, Twetman S, Stecksén-Blicks S. 2008. Sugar fermentation in probiotic bacteria – an *in vitro* study. *Oral Microbiol Immunol* 23:482-485.
- Hemalatha R, Ouwehand AC, Saarinen MT, Prasad UV, Swetha K, Bhaskar V. 2017. Effect of probiotic supplementation on total lactobacilli, bifidobacteria and short chain fatty acids in 2–5-year-old children. *Microb Ecol Health Dis* 28:1298340.
- Hill C. 2012. Virulence or niche factors: what's in a name? *J Bacteriol* 194:5725–5727.
- Ho Y-H, Lu Y-C, Chang H-C, Lee S-Y, Tsai M-F, Huang Y-T, Hsu T-Y. 2014. Daily Intake of Probiotics with High IFN- $\gamma$ /IL-10 Ratio Increases the Cytotoxicity of Human Natural Killer Cells: A Personalized Probiotic Approach. *J Immunol Res* 721505.
- Hutt P, Koll P, Stsepetova J, Alvarez B, Mandar R, Krogh-Andersen K, Marcotte H, Hammarström L, Mikelsaar M. 2011. Safety and persistence of orally administered human *Lactobacillus* sp. strains in healthy adults. *Benef Microbes* 2:79-90.
- Jagadeesan B, Koo OK, Kim KP, Burkholder KM, Mishra KK, Aroonnu A, Bhunia AK. 2010. LAP, an alcohol acetaldehyde dehydrogenase enzyme in *Listeria*, promotes bacterial adhesion to enterocyte-like Caco-2 cells only in pathogenic species. *Microbiology* 156:2782–2795.

- Jernberg C, Sullivan A, Edlund C, Jansson JK. 2005. Monitoring of antibiotic-induced alterations in the human intestinal microflora and detection of probiotic strains by use of terminal restriction fragment length polymorphism. *Appl Environ Microbiol* 71:501-506.
- Jespersen L, Tarnow I, Eskesen D, Morberg CM, Michelsen B, Bügel S, Dragsted LO, Rijkers GT, Calder PC. 2015. Effect of *Lactobacillus paracasei* subsp. *paracasei*, *L. casei* 431 on immune response to influenza vaccination and upper respiratory tract infections in healthy adult volunteers: a randomized, double-blind, placebo-controlled, parallel-group study. *Am J Clin Nutr* 101:188-196.
- Jia X, Wang W, Song Y, Li N. 2011. A 90-day oral toxicity study on a new strain of *Lactobacillus paracasei* in rats. *Food Chem Toxicol* 49:1148-1151.
- Juntunen M, Kirjavainen PV, Ouwehand AC, Salminen SJ, Isolauri E. 2001. Adherence of probiotic bacteria to human intestinal mucus in healthy infants and during rotavirus infection. *Clin Diagn Lab Immunol* 8:293-296.
- Kirjavainen P, Ouwehand AC, Isolauri E, Salminen SJ. 1998. The ability of probiotic bacteria to bind to human intestinal mucus. *FEMS Microbiol Lett* 167:185-189.
- Kruszewska D, Lan J, Lorca G, Yanagisawa N, Marklinder I, Ljungh A. 2002. Selection of lactic acid bacteria as probiotic strains by *in vitro* tests. *Microecol Ther* 29:37-49.
- Lahner E, Esposito G, Zullo A, Hassan C, Cannaviello C, Di Paolo MC, Pallotta L, Garbagna N, Grossi E, Annibale B. 2012. High-fibre diet and *Lactobacillus paracasei* B21060 in symptomatic uncomplicated diverticular disease. *World J Gastroenterol* 18:5918-5924.
- Larsen N, Michaelsen KF, Paerregaard A, Vogensen FK, Jakobsen M. 2009. A comparative study on adhesion and recovery of potential probiotic strains of *Lactobacillus* spp. by *in vitro* assay and analysis of human colon biopsies. *Microb Ecol Health Dis* 21:95-99.
- Lee A, L YJ, Yoo HJ, Kim M, Chang Y, Lee DS, Lee JH. 2017. Consumption of dairy yogurt containing *Lactobacillus paracasei* ssp. *paracasei*, *Bifidobacterium animalis* ssp. *lactis* and heat-treated *Lactobacillus plantarum* improves immune function including natural killer cell activity. *Nutrients* 9:558-566.
- Lenoir-Wijnkoop I, Nuijten MJC, Craig J, Butler CC. 2014. Nutrition economic evaluation of a probiotic in the prevention of antibiotic-associated diarrhea. *Front Pharmacol* 5:13.
- Lin W-Y, Fu L-S, Lin H-K, Shen C-Y, Chen Y-J. 2014. Evaluation of the effect of *Lactobacillus paracasei* (HF.A00232) in children (6-13 years old) with perennial allergic rhinitis: a 12-week, double-blind, randomized, placebo-controlled study. *Pediatr Neonatol* 55:181-188.
- Liu Q, Duan ZP, Ha DK, Bengmark S, Kurtovic J, Riordan SM. 2004. Synbiotic modulation of the gut flora: effect on minimal hepatic encephalopathy in patients with cirrhosis. *Hepatology* 39:1441-1449.
- Ljungh A, Lan J, Yanagisawa N. 2002. Isolation, selection and characteristics of *Lactobacillus paracasei* subsp. *paracasei* F-19. *Microb Ecol Health Dis* 14:4-6
- Lombardo L, Vernetto A, Blanco I. 2009. Clinical evaluation of *Lactobacillus paracasei* subsp. *paracasei* F-19 with gluco-oligosaccharides in the short-term treatment of irritable bowel syndrome. *Microb Ecol Health Dis* 21:28-32.
- Lundelin K, Poussa T, Salminen S, Isolauri E. 2017. Long-term safety and efficacy of perinatal probiotic intervention: evidence from a follow-up study of four randomized, double-blind, placebo-controlled trials. *Pediatr Allergy Immunol* 28:170-175.

- Maretti C and Cavallini G. 2017. The association of a probiotic with a prebiotic (Flortec, Bracco) to improve the quality/quantity of spermatozoa in infertile patients with idiopathic oligoasthenoteratospermia: a pilot study. *Andrology* 5:439-444.
- Martarelli D, Verdenelli MC, Scuri S, Cocchioni M, Silvi S, Cecchini C, Pompei P. 2011. Effect of a probiotic intake on oxidant and antioxidant parameters in plasma of athletes during intense exercise training. *Curr Microbiol* 62:1689-1696.
- Matto J, Fonden R, Tolvanen T, von Wright A, Vilpponen-Salmela T, Satokari R, Saarela M. 2006. Intestinal survival and persistence of probiotic *Lactobacillus* and *Bifidobacterium* strains administered in triple-strain yoghurt. *Int Dairy J* 16:1174-1180.
- Miettinen M, Vuopio-Varkila J, Varkila K. 1996. Production of human tumor necrosis factor alpha, interleukin-6 and interleukin-10 is induced by lactic acid bacteria. *Infect Immun* 64:5403-5404.
- Morelli L and Campominski E. 2002. Genetic stability of *Lactobacillus paracasei* subsp. *paracasei* F-19. *Microb Ecol Health Dis* 14:14-16.
- Morovic W, Roper JM, Smith AB, Mukerji P, Stahl B, Rae JC, Ouwehand AC. 2017. Safety evaluation of HOWARU® Restore (*Lactobacillus acidophilus* NCFM, *Lactobacillus paracasei* Lpc-37, *Bifidobacterium animalis* subsp. *lactis* Bl-04 and *B. lactis* Bi-07) for antibiotic resistance, genomic risk factors, and acute toxicity. *Food Chem Toxicol* 110:316-324.
- Nardone G, Compare D, Liguori E, Di Mauro V, Rocco A, Barone M, Napol, A, Lapi D, Iovene M R, Colantuoni A. 2010. Protective effects of *Lactobacillus paracasei* F-19 in a rat model of oxidative and metabolic hepatic injury. *Am J Physiol Gastrointest Liver Physiol* 299:G669-676.
- Nembrini C, Singh A, De Castro CA, Mercenier A, Nutten S. 2015. Oral administration of *Lactobacillus paracasei* NCC 2461 for the modulation of grass pollen allergic rhinitis: a randomized, placebocontrolled study during the pollen season. *Clin Transl Allergy* 5:41-46.
- Nerstedt A, Nilsson EC, Ohlson K, Hakansso, J, Svensson T, Lowenadler B, Svensson UK, Mahlapuu M. 2007. Administration of *Lactobacillus* evokes coordinated changes in the intestinal expression profile of genes regulating energy homeostasis and immune phenotype in mice. *Br J Nutr* 16:1-11.
- Ohlson K, Bjorneholm S, Fonden R, Svensson U. 2002. *Lactobacillus* F-19 - a probiotic strain suitable for consumer products. *Microb Ecol Health Dis* 14:27-32.
- Ouwehand AC, DongLian C, Weijian X, Stewart M, Ni J, Stewart T, Miller LE. 2014. Probiotics reduce symptoms of antibiotic use in a hospital setting: a randomized dose response study. *Vaccine* 32:458-463.
- Passariello A, Terrin G, Cecere G, Micillo M, De Marco G, Di Costanzo M, Cosenza L, Leone L, Nocerino R, Canani RB. 2012. Randomised clinical trial: efficacy of a new synbiotic formulation containing *Lactobacillus paracasei* B21060 plus arabinogalactan and xilooligosaccharides in children with acute diarrhea. *Aliment Pharmacol Ther* 35:782-788.
- Pedersen SMM, Nebel C, Nielsen NC, Andersen HJ, Olsson J, Simren M, Ohman L, Svensson U, Bertram HC, Malmendal A. 2011. A GC-MS-based metabonomic investigation of blood serum from irritable bowel syndrome patients undergoing intervention with acidified milk products. *Eur Food Res Technol* 233:1013-1021.

- Peluso I, Fina D, Caruso R, Stolfi C, Caprioli F, Fantini MC, Caspani G, Grossi E, Di Iorio L, Paone FM, Pallone F, Monteleone G. 2007. *Lactobacillus paracasei* subsp. *Paracasei* B21060 suppresses human T-cell proliferation. *Infect Immun* 75:1730-1737.
- Perrin Y, Nutton S, Audran R, Berger B, Bibiloni R, Wassenberg J, Barbier N, Aubert V, Moulin J, Singh A, Magliola C, Mercenier A, Spertini F. 2014. Comparison of two oral probiotic preparations in a randomized crossover trial highlights a potentially beneficial effect of *Lactobacillus paracasei* NCC2461 in patients with allergic rhinitis. *Clin Transl Allergy* 4:1-12.
- Pino A, Van Hoorde K, Pitino I, Russo N, Carpino S, Caggia C, Randazzo CL. 2017. Survival of potential probiotic lactobacilli used as adjunct cultures on Pecorino Siciliano cheese ripening and passage through the gastrointestinal tract of healthy volunteers. *Int J Food Microbiol* 252:42-52.
- Plaza-Diaz J, Gomez-Liorente C, Campana-Martin L, Matencio E, Ortuno I, Martinez-Silla R, Gomez-Gallego C, Periago MJ, Ros, G, Chenoli E, Genoves S, Casinos B, Silva A, Corella D, Portoles O, Romero F, Ramon D, Perezde la Cruz A, Gil A, Fontana L. 2013. Safety and immunomodulatory effects of three probiotic strains isolated from the feces of breast-fed infants in healthy adults: SETOPROB study. *PLoS ONE* 8:e78111.
- Pu F, Guo Y, Li M, Zhu H, Wang S, Shen X, He M, Huang C, He Fang. 2017. Yogurt supplemented with probiotics can protect the healthy elderly from respiratory infections: A randomized controlled open-label trial. *Clin Interv Aging* 12:1223-1231.
- Rask C, Adlerberth I, Berggren A, Ahren IL, Wold AE. 2013. Differential effect on cell-mediated immunity in human volunteers after intake of different lactobacilli. *Clin Exp Immunol* 172:321-332.
- Rautava S, Kainonen E, Salminen S, Isolauri E. 2012. Maternal probiotic supplementation during pregnancy and breast-feeding reduces the risk of eczema in the infant. *J Allergy Clin Immunol* 130:1355-1360.
- Rayes N, Seehofer D, Hansen S, Boucsein K, Muller AR, Serke S, Bengmark S, Neuhaus P. 2002a. Early enteral supply of *Lactobacillus* and fiber versus selective bowel decontamination: a controlled trial in liver transplant recipients. *Transplantation* 74:123-128.
- Rayes N, Seehofer D, Müller AR, Hansen S, Bengmark S, Neuhaus P. 2002b. Influence of probiotics and fibre on the incidence of bacterial infections following major abdominal surgery - results of a prospective trial. *Z Gastroenterol* 40:869-876. [Article in German with English abstract]
- Rayes N, Hansen S, Seehofer D, Muller AR, Serke S, Bengmark S, Neuhaus P. 2002c. Early enteral supply of fiber and lactobacilli versus conventional nutrition: a controlled trial in patients with major abdominal surgery. *Nutrition* 18:609-615.
- Rayes N, Seehofer D, Theruvath T, Schiller RA, Langrehr JM, Jonas S, Bengmark S, Neuhaus P. 2005. Supply of pre- and probiotics reduces bacterial infection rates after liver transplantation – a randomized, double-blind trial. *Am J Transplant* 5:125-130.
- Rayes N, Seehofer D, Theruvath T, Mogl M, Langrehr JM, Nussler NC, Bengmark S, Neuhaus P. 2007. Effect of enteral nutrition and synbiotics on bacterial infection rates after pylorus-preserving pancreatoduodenectomy: a randomized, double-blind trial. *Ann Surg* 246:36-41.
- Rayes N, Pilarski T, Stockmann M, Bengmark S, Newhaus P, Seehofer D. 2012. Effect of pre- and probiotics on liver regeneration after resection: a randomised, double-blind pilot study. *Benef Microb* 2:237-244.

- Reygagne P, Bastien P, Couavoux MP, Philippe D, Renouf M, Castiel-Higounenc I, Gueniche A. 2017. The positive benefit of *Lactobacillus paracasei* NCC2461 ST11 in healthy volunteers with moderate to severe dandruff. *Benef Microbes* 8:671-680.
- Riezzo G, Orlando A, D'Attoma B, Guerra V, Valerio F, Lavermicocca P, De Candia S, Russo F. 2012. Randomised clinical trial: efficacy of *Lactobacillus paracasei* enriched artichokes in the treatment of patients with functional constipation – a double-blind, controlled, crossover study. *Aliment Pharmacol Ther* 35:441-450.
- Riordan SM, Skinner N, Nagree A, McCallum H, McIver CJ, Kurtovic J, Hamilton JA, Bengmark S, Williams R, Visvanathan K. 2003. Peripheral blood mononuclear cell expression of toll-like receptors and relation to cytokine levels in cirrhosis. *Hepatology* 37:1154-1164.
- Ritthagol W, Saetang C, Teanpaisan R. 2014. Effect of probiotics containing *Lactobacillus paracasei* SD1 on salivary mutans streptococci and lactobacilli in orthodontic cleft patients: a double-blinded, randomized, placebo- controlled study. *Cleft Palate Craniofac J* 51:257-263.
- Roessler A, Forssten SD, Gleis M, Ouwehand AC, Jahreis G. 2012. The effect of probiotics on faecal microbiota and genotoxic activity of faecal water in patients with atopic dermatitis: a randomized, placebo-controlled study. *Clin Nutr* 31:22-29.
- Saito Y, Fujii M, Watanabe T, Maruyama K, Kowatari Y, Ogata H, Kumagai T. 2017. Randomized, double-blind, placebo-controlled, parallel-group study of the effect of *Lactobacillus paracasei* K71 intake on salivary release of secretory immunoglobulin A. *Food Health* 36:55-63.
- Saxelin M, Grenov B, Svensson U, Fonden R, Reniero R, Mattila-Sandholm T. 1999. The technology of probiotics. *Trends Food Sci Technol* 10:387-392.
- Schott AS, Behr J, Quinn J, Vogel RF. 2016. MALDI-TOF mass spectrometry enables a comprehensive and fast analysis of dynamics and qualities of stress responses of *Lactobacillus paracasei* subsp. *paracasei* F-19. *PLoS One* 11.
- Simakachorn N, Bibiloni R, Yimyaem P, Tongpenyai Y, Varavithaya W, Grathwohl D, Reuteler G, Maire JC, Blum S, Steenhout P, Benyacoub J, Schiffrin EJ. 2011. Tolerance, safety, and effect on the faecal microbiota of an enteral formula supplemented with pre- and probiotics in critically ill children. *J Pediatr Gastroenterol Nutr* 53:174-181.
- Simren M, Ohman L, Olsson J, Svensson U, Ohlson K, Posserud I, Strid, H. 2010. Clinical trial: the effects of a fermented milk containing three probiotic bacteria in patients with irritable bowel syndrome - a randomized, double-blind, controlled study. *Aliment Pharmacol Ther* 31:218-227.
- Sondergaard B, Olsson J, Ohlson K, Svensson U, Bytzer P, Ekesbo R. 2011. Effects of probiotic fermented milk on symptoms and intestinal flora in patients with irritable bowel syndrome: a randomized, placebo-controlled trial. *Scand J Gastroenterol* 46:663-672.
- Spindler-Vesel A, Bengmark S, Vovk I, and Kompan L. 2007. Synbiotics, prebiotics, glutamine, or peptide in early enteral nutrition: a randomized study in trauma patients. *J Parent Ent Nutr* 31:1-8.
- Sullivan A, Palmgren AC, Nordh CE. 2001. Effect of *Lactobacillus paracasei* on intestinal colonisation of *Lactobacilli*, *Bifidobacteria* and *Clostridium difficile* in elderly persons. *Anaerobe* 7:67-70.

- Sullivan A, Bennet R, Viitanen M, Palmgren AC, Nord CE. 2002. Influence of *Lactobacillus* F-19 on intestinal microflora in children and elderly persons and impact on *Helicobacter pylori* infections. *Microb Ecol Health Dis* 14:17-21.
- Sullivan A, Barkholt L, Nord CE. 2003. *Lactobacillus acidophilus*, *Bifidobacterium lactis* and *Lactobacillus* F-19 prevent antibiotic-associated ecological disturbances of *Bacteroides fragilis* in the intestine. *J Antimicrob Chemother* 52:308-311.
- Sullivan A, Johansson A, Svennungsson B, Nord CE. 2004. Effect of *Lactobacillus* F-19 on the emergence of antibiotic-resistant microorganisms in the intestinal microflora. *J Antimicrob Chemother* 54:791-797.
- Sullivan A and Nord CE. 2006. Probiotic lactobacilli and bacteraemia in Stockholm. *Scand J Inf Dis* 38:327-331.
- Sullivan A, Nord CE, Evengard B. 2009. Effect of supplement with lactic-acid producing bacteria on fatigue and physical activity in patients with chronic fatigue syndrome. *Nutr J* 8:1-6.
- Szajewska H, Ruszczynski M, Szymanski H, Sadowska-Krawczenko I, Piwowarczyk A, Rasmussen PB, Kristensen MB, West CE, Hernell O. 2017. Effects of infant formula supplemented with prebiotics compared with synbiotics on growth up to the age of 12 mo: a randomized controlled trial. *Pediatr Res* 81:752-758.
- Teanpaisan R and Piwat S. 2014. *Lactobacillus paracasei* SD1, a novel probiotic, reduces mutans streptococci in human volunteers: a randomized placebocontrolled trial. *Clin Oral Investig* 18:857-862.
- Trautvetter U, Ditscheid B, Kiehntopf M, Jahreis G. 2012. A combination of calcium phosphate and probiotics beneficially influences intestinal lactobacilli and cholesterol metabolism in humans. *Clin Nutr* 31:230-237.
- Tseng W-T, Shih T-W, Liu S-H, Pan T-M. 2015. Safety and mutagenicity evaluation of Vigii 101 powder made from *Lactobacillus paracasei* subsp. *paracasei* NTU 101. *Regul Toxicol Pharmacol* 71:148-157.
- Twetman L, Larsen U, Fiehn NE, Stecksen-Blicks C, Twetman S. 2009. Coaggregation between probiotic bacteria and caries-associated strains: An *in vitro* study. *Acta Odontol Scand* 27: 1-5.
- Valerio F, de Candia S, Lonigro SL, Russo F, Riezzo G, Orlando A, De Bellis P, Sisto A, Lavermicocca P. 2011. Role of the probiotic strain *Lactobacillus paracasei* LMGP22043 carried by artichokes in influencing faecal bacteria and biochemical parameters in human subjects. *J Appl Microbiol* 111:155-164.
- Valsecchi C, Marseglia A, Montagna L, Tagliacarne SC, Elli M, Licari A, Marseglia GL, Castellazzi AM. 2014. Evaluation of the effects of a probiotic supplementation with respect to placebo on intestinal microflora and secretory IgA production, during antibiotic therapy, in children affected by recurrent airway infections and skin symptoms. *J Biol Regul Homeost Agents* 28:117-124.
- Vankerckhoven V, Huys G, Vancanneyt M, Vael C, Klare I, Romond M-B, Entenze JM, Moreillon P, Wind RD, Knol J, Wiertz E, Pot B, Vaughan EE, Kahlmeter G, Goossens H. 2008. Biosafety assessment of probiotics used for human consumption: recommendations from the EU-PROSAFE project. *Trends Food Sci Technol* 19:102-114.

- Verdenelli MC, Silvi S, Cecchini C, Orpianesi C, Cresci A. 2011. Influence of a combination of two potential probiotic strains, *Lactobacillus rhamnosus* IMC 501<sup>®</sup> and *Lactobacillus paracasei* IMC 502<sup>®</sup> on bowel habits of healthy adults. *Lett Appl Microbiol* 52:596-602.
- Vicariotto F. 2014. Effectiveness of an association of a cranberry dry extract, D-mannose, and the two microorganisms *Lactobacillus plantarum* LP01 and *Lactobacillus paracasei* LPC09 in women affected by cystitis: a pilot study. *J Clin Gastroenterol* 48 Suppl 1:S96-101.
- Videhult FK, Ohlund I, Stenlund H, Hernell O, West CE. 2015a. Probiotics during weaning: a follow-up study on effects on body composition and metabolic markers at school age. *Eur J Nutr* 54:355-363.
- Videhult FK, Andersson Y, Ohlund I, Stenlund H, Hernell O, West CE. 2015b. Impact of probiotics during weaning on the metabolic and inflammatory profile: follow-up at school age. *Int J Food Sci Nutr* 66:686-691.
- Vlieger AM, Robroch A, van Buuren S, Kiers J, Rijkers G, Benninga MA, te Biesebeke R. 2009. Tolerance and safety of *Lactobacillus paracasei* ssp. *paracasei* in combination with *Bifidobacterium animalis* ssp. *lactis* in a prebiotic-containing infant formula: a randomised controlled trial. *Br J Nutr* 102:869-875.
- Wang MF, Lin HC, Wang YY, Hsu CH. 2004. Treatment of perennial allergic rhinitis with lactic acid bacteria. *Pediatr Allergy Immunol* 15:152-158.
- Wassenaar TM, Gaastra W. 2001. Bacterial virulence: can we draw the line? *FEMS Microbiol Lett* 201:1-7.
- Wassenberg J, Nutten S, Audran R, Barbier N, Aubert V, Moulin J, Mercenier A, Spertini F. 2011. Effect of *Lactobacillus paracasei* ST11 on a nasal provocation test with grass pollen in allergic rhinitis. *Clin Exp Allergy* 41:565-573.
- Wattanarat O, Makeudom A, Sastraruji T, Piwat S, Tianviwat S, Teanpaisan R, Krisanaprakornkit S. 2015. Enhancement of salivary human neutrophil peptide 1-3 levels by probiotic supplementation. *BMC Oral Health* 15:19-29.
- West C, Gothefors L, Granström M, Kayhty H, Hammarstrom M-L, Hernell O. 2008. Effects of feeding probiotics during weaning on infections and antibody responses to diphtheria, tetanus and Hib vaccines. *Pediatr Allergy Immunol* 19:53-60.
- West C, Hammarstrom M-L, Hernell O. 2009. Probiotics during weaning reduce the incidence of eczema. *Pediatr Allergy Immunol* 20:430-437.
- West CE, Hernell O, Andersson Y, Sjostedt M, Hammarstrom ML. 2012a. Probiotic effects on T-cell maturation in infants during weaning. *Clin Exp Allergy* 42:540-549.
- West NP, Pyne DB, Cripps AW, Christophersen CT, Conlon MA, Fricker PA. 2012b. Gut Balance, a synbiotic supplement, increases fecal *Lactobacillus paracasei* but has little effect on immunity in healthy physically active individuals. *Gut Microbes* 3:221-227.
- West CE, Hammarstrom ML, Hernell O. 2013. Probiotics in primary prevention of allergic disease—follow-up at 8-9 years of age. *Allergy* 68:1015-1020.
- Zampieri N, Pietrobelli A, Biban P, Soffiati M, Dall'Agnola A, Camoglio FS. 2013. *Lactobacillus paracasei* subsp. *paracasei* F-19 in Bell's Stage 2 of necrotizing enterocolitis. *Minerva Pediatr* 65:353-360.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640-2644.

- Zhang H, Wang Y, Sun J, Guo Z, Guo H, Ren F. 2013a. Safety evaluation of *Lactobacillus paracasei* subsp. *paracasei* LC-01, a probiotic bacterium. *J Microbiol* 51:633-638.
- Zhang H, Sun J, Liu X, Hong C, Zhu Y, Liu A, Li S, Guo H, Ren F. 2013b. *Lactobacillus paracasei* subsp. *paracasei* LC01 positively modulates intestinal microflora in healthy young adults. *J Microbiol* 51:777-782.

**From:** [Jim](#)  
**To:** [Harry, Molly](#)  
**Cc:** [Jim Heimbach](#)  
**Subject:** Re: GRAS Notice No. 000810  
**Date:** Thursday, February 28, 2019 3:53:49 PM

---

Molly—

Thanks for your inquiry. It means that no organisms were detected in any of ten samples of 10 g each.

Regards,  
Jim

Sent from my iPhone

On Feb 28, 2019, at 2:42 PM, Harry, Molly <[Molly.Harry@fda.hhs.gov](mailto:Molly.Harry@fda.hhs.gov)> wrote:

Hi Dr. Heimbach,

Our scientific review of GRN 000810 is almost complete. However, we'll like you to provide some minor clarification regarding information provided in the notice. On Table 1 (page 8 of the GRN), the specifications provided for *Enterobacteriaceae*, *Cronobacter sakazakii*, and *Salmonella* spp. is "Absent/10x10g." Please clarify what 10x10g means.

Thanks,

**Molly A. Harry**  
*Consumer Safety Officer*

**Center for Food Safety and Applied Nutrition**  
**Office of Food Additive Safety**  
**U.S. Food and Drug Administration**  
Tel: 240-402-1075  
[Molly.Harry@fda.hhs.gov](mailto:Molly.Harry@fda.hhs.gov)

[<image001.png>](#)

**From:** Jim <jheimbach@va.metrocast.net>  
**Sent:** Friday, March 29, 2019 2:55 PM  
**To:** Harry, Molly <Molly.Harry@fda.hhs.gov>  
**Subject:** Re: GRAS Notice No. 000810

Dear Molly—

No, they are not.

Regards,  
Jim

Sent from my iPhone

On Mar 29, 2019, at 1:12 PM, Harry, Molly <[Molly.Harry@fda.hhs.gov](mailto:Molly.Harry@fda.hhs.gov)> wrote:

Hi Dr. Heimbach,

I have a quick question for you. The method of production for *L. paracasei* F-19 includes addition of cryoprotectants and excipients steps. Please clarify whether (or not) the cryoprotectants and excipients are from allergenic sources.

Thanks, Molly

**CONCLUSION OF THE EXPERT PANEL:  
DETERMINATION OF THE GRAS  
STATUS OF THE USE OF *LACTOBACILLUS*  
*PARACASEI* SSP. *PARACASEI* STRAIN F-19**

**Prepared for:  
Arla Foods Ingredients**

**July 2018**

## CONCLUSION OF THE EXPERT PANEL:

We, the members of the Expert Panel, have individually and collectively critically evaluated the publicly available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 summarized in a monograph, *Generally Recognized As Safe (GRAS) Determination for the Intended Use of Lactobacillus paracasei* ssp. *paracasei* strain F-19 (July 2018), prepared by JHeimbach LLC, and other material deemed appropriate or necessary. Our evaluation included critical evaluation of the identity, genotypic and phenotypic characteristics of the strain, production methods, potential exposure resulting from the intended use of the strain, and published research bearing on the safety of *Lactobacillus paracasei* ssp. *paracasei* strain F-19. Our summary and conclusion resulting from this critical evaluation are presented below.

### Summary

- *Lactobacillus paracasei* ssp. *paracasei* strain F-19 is intended to be added to powdered milk-based non-exempt infant formula intended for consumption by healthy term infants at a level consistent with current Good Manufacturing Practice to result in a daily intake of  $10^9$  cfu of the strain. The target population is healthy infants and toddlers aged 0-3 years of age. It is not expected that these infants and toddlers will have other dietary sources of *L. paracasei* ssp. *paracasei* strain F-19.
- *Lactobacillus paracasei* ssp. *paracasei* strain F-19 was isolated from the colon of a healthy adult and deposited in the Belgian Coordinated Collections of Microorganisms, Microbiology Laboratory (BCCM/LMG) under deposit number LMG-P-17806. The taxonomic placement of *L. paracasei* ssp. *paracasei* strain F-19 was determined by ribotyping and confirmed by pulsed-field gel electrophoresis and amplified fragment length polymorphism.
- A complete genome sequence of the strain revealed a circular chromosome of 3.1 Mb, a linear plasmid of 93.7 Kb, three circular plasmids of 8.7 Kb, 6.3 Kb, and 2.2 Kb, and two linear fragments of 2.3 Kb and 3.4 Kb. The genome was analyzed for the presence of antibiotic resistance and virulence genes; none was detected. One gene labeled “ornithine decarboxylase” was found, but it is not accompanied by a transporter required for generation of putrescine.
- *Lactobacillus paracasei* ssp. *paracasei* strain F-19 is produced under certification that systems are in full compliance with ISO 9001:2008, ISO 22000:2005, ISO/TS 2202-1:2009, and Food Safety System Certification 22000. All processing aids meet food-grade specifications, are used consistent with their regulatory status, and are suitable for human consumption.
- *Lactobacillus paracasei* received Qualified Presumption of Safety (QPS) assignment in the initial review in 2007 and this status was renewed each year from 2008 through 2018. The safe history of human exposure to *L. paracasei* ssp. *paracasei* strains is strongly supported by a large body of published research, including 4 oral toxicity studies in mice and rats and more than 50 studies in human infants, children, and adults enrolling more than 7,000 individuals. Participants in these studies received the probiotic at daily levels

up to  $10^{12}$  cfu/day for as long as 9 months with no reported adverse events associated with the probiotic intervention.

- A decision-tree analysis (Pariza et al. 2015) determined that *Lactobacillus paracasei* ssp. *paracasei* strain F-19 is deemed to be safe for use in the manufacture of food, probiotics, and dietary supplements for human consumption based on the following responses:
  - Has the strain been characterized for the purpose of assigning an unambiguous genus and species name using currently accepted methodology? **YES**
  - Has the strain genome been sequenced? **YES**
  - Is the strain genome free of genetic elements encoding virulence factors and/or toxins associated with pathogenicity? **YES**
  - Is the strain genome free of functional and transferable antibiotic resistance gene DNA? **YES**
  - Does the strain produce antimicrobial substances? **NO**
  - Has the strain been genetically modified using rDNA techniques? **NO**
  - Was the strain isolated from a food that has a history of safe consumption for which the species, to which the strain belongs, is a substantial and characterizing component (not simply an 'incidental isolate')? **NO** (THE STRAIN WAS ISOLATED FROM THE COLON OF A HEALTHY HUMAN)
  - Does the strain induce undesirable physiological effects in appropriately designed safety evaluation studies? **NO**

## Conclusion

We, the undersigned members of the Expert Panel, are qualified by scientific education and experience to evaluate the safety of microorganisms intended for addition to foods, including infant formulas, as probiotics. We have individually and collectively critically evaluated the publicly available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 summarized in a monograph, *Generally Recognized As Safe (GRAS) Determination for the Intended Use of Lactobacillus paracasei* ssp. *paracasei* strain F-19 (July 2018), prepared by JHeimbach LLC, and other material deemed appropriate or necessary.

We have individually and collectively determined that no evidence exists in the available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 that demonstrates, or suggests reasonable grounds to suspect, a hazard to infants under the intended conditions of use of *Lactobacillus paracasei* ssp. *paracasei* strain F-19.

We unanimously conclude that the intended addition to powdered milk-based non-exempt infant formula intended for consumption by healthy term infants of *Lactobacillus paracasei* ssp. *paracasei* strain F-19, produced consistent with current good manufacturing practice (cGMP) and meeting the food-grade specifications presented in the monograph, at a level consistent with cGMP to provide a daily intake of  $10^9$  cfu of the strain, is safe and is GRAS by scientific procedures.

It is our opinion that other qualified and competent scientists reviewing the same publicly available information would reach the same conclusions.

Joseph F. Borzelleca, Ph.D.  
Professor Emeritus  
Virginia Commonwealth University School of Medicine  
Richmond, Virginia

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

3/July 2018

Berthold V. Koletzko, Dr med, Dr med habil (M.D., Ph.D.)  
Professor of Pediatrics  
University of Munich  
Munich, Germany

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Michael W. Pariza, Ph.D.  
Professor Emeritus  
University of Wisconsin—Madison  
Madison, Wisconsin

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

## Conclusion

We, the undersigned members of the Expert Panel, are qualified by scientific education and experience to evaluate the safety of microorganisms intended for addition to foods, including infant formulas, as probiotics. We have individually and collectively critically evaluated the publicly available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 summarized in a monograph, *Generally Recognized As Safe (GRAS) Determination for the Intended Use of Lactobacillus paracasei* ssp. *paracasei* strain F-19 (July 2018), prepared by JHeimbach LLC, and other material deemed appropriate or necessary.

We have individually and collectively determined that no evidence exists in the available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 that demonstrates, or suggests reasonable grounds to suspect, a hazard to infants under the intended conditions of use of *Lactobacillus paracasei* ssp. *paracasei* strain F-19.

We unanimously conclude that the intended addition to powdered milk-based non-exempt infant formula intended for consumption by healthy term infants of *Lactobacillus paracasei* ssp. *paracasei* strain F-19, produced consistent with current good manufacturing practice (cGMP) and meeting the food-grade specifications presented in the monograph, at a level consistent with cGMP to provide a daily intake of  $10^9$  cfu of the strain, is safe and is GRAS by scientific procedures.

It is our opinion that other qualified and competent scientists reviewing the same publicly available information would reach the same conclusions.

Joseph F. Borzelleca, Ph.D.  
Professor Emeritus  
Virginia Commonwealth University School of Medicine  
Richmond, Virginia

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Berthold V. Koletzko, Dr med, Dr med habil (M.D., Ph.D.)  
Professor of Pediatrics  
University of Munich  
Munich, Germany

Signature: \_\_\_\_\_ Date: 4 Aug 2017

Michael W. Pariza, Ph.D.  
Professor Emeritus  
University of Wisconsin—Madison  
Madison, Wisconsin

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

## Conclusion

We, the undersigned members of the Expert Panel, are qualified by scientific education and experience to evaluate the safety of microorganisms intended for addition to foods, including infant formulas, as probiotics. We have individually and collectively critically evaluated the publicly available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 summarized in a monograph, *Generally Recognized As Safe (GRAS) Determination for the Intended Use of Lactobacillus paracasei* ssp. *paracasei* strain F-19 (July 2018), prepared by JHeimbach LLC, and other material deemed appropriate or necessary.

We have individually and collectively determined that no evidence exists in the available information on *Lactobacillus paracasei* ssp. *paracasei* strain F-19 that demonstrates, or suggests reasonable grounds to suspect, a hazard to infants under the intended conditions of use of *Lactobacillus paracasei* ssp. *paracasei* strain F-19.

We unanimously conclude that the intended addition to powdered milk-based non-exempt infant formula intended for consumption by healthy term infants of *Lactobacillus paracasei* ssp. *paracasei* strain F-19, produced consistent with current good manufacturing practice (cGMP) and meeting the food-grade specifications presented in the monograph, at a level consistent with cGMP to provide a daily intake of  $10^9$  cfu of the strain, is safe and is GRAS by scientific procedures.

It is our opinion that other qualified and competent scientists reviewing the same publicly available information would reach the same conclusions.

Joseph F. Borzelleca, Ph.D.  
Professor Emeritus  
Virginia Commonwealth University School of Medicine  
Richmond, Virginia

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Berthold V. Koletzko, Dr med, Dr med habil (M.D., Ph.D.)  
Professor of Pediatrics  
University of Munich  
Munich, Germany

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Michael W. Pariza, Ph.D.  
Professor Emeritus  
University of Wisconsin—Madison  
Madison, Wisconsin

Signature: Michael W. Pariza, Ph.D. Digitally signed by Michael W. Pariza, Ph.D.  
DN: cn=Michael W. Pariza, Ph.D., o=Michael W. Pariza  
Consulting LLC, ou=Member, email=rmwpariza@gmail.com,  
c=US  
Date: 2018.08.01 10:12:59 -05'00' Date: August 1, 2018