



SedgeStop™

NUTSEGE WEED KILLER

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Do not allow people or pets to enter the treated area until dusts have settled.

1. Product Description

Sedge Stop™ Nutsedge Weed Killer is a selective herbicide for post-emergent control of listed broadleaf weeds, sedges, kyllinga, and rush species commonly found in turfgrasses. It can be used on established warm-season turfgrasses (including improved varieties of St. Augustinegrass) and cool-season turfgrasses. Fall applications provide control of winter weeds such as henbit, false dandelion and chickweed.

2. Weeds Controlled


Sedge Stop Nutsedge Weed Killer will control the following weeds:

Sedges, Kyllinga, and Rushes: Yellow Nutsedge, Purple Nutsedge, Green Kyllinga, False Green Kyllinga, Cockscomb Kyllinga, and Path Rush.

Broadleaf Weeds: Common Chickweed, Hairy Bittercress, Henbit, Buckhorn Plantain, Dollarweed (Pennywort), False Dandelion, Florida Betony, Ground Ivy, and Virginia Buttonweed.

3. How To Identify Nutsedge

Yellow Nutsedge and Purple Nutsedge are common weeds in residential lawns. They are sometimes referred to as Nutgrass or Watergrass. They have distinctive characteristics that make them easy to identify. Both weeds become visible in a lawn during the summer. The leaves are glossy, light green, hairless, and usually grow faster than the surrounding turf. Stems are distinctly triangular. Leaves are arranged in groups of three and are V-shaped with a prominent midrib.



Getting Started

How To Use:

Shaker can allows for targeted applications. Apply to weed leaves when wet or dry.

How Much To Use:

Apply 2 lb. of product per 500 sq. ft.

ACTIVE INGREDIENT:

Pyrimisulfan..... 0.025%

OTHER INGREDIENTS: 99.975%

TOTAL 100.000%

THIS PRODUCT CONTAINS:

0.5 lb. Pyrimisulfan per ton or 0.025% (0.00025 lb. pyrimisulfan per pound).

**KEEP OUT OF REACH OF CHILDREN
CAUTION**



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Physical and Chemical Hazards

Do not mix or allow to come into contact with an oxidizing agent. Hazardous chemical reaction may occur.

4. Where To Use

Ornamental turfgrass established around residential dwellings and buildings and other non-cropland sites.

The following established turfgrass species may be treated:

- **Cool-Season Species:** Kentucky bluegrass, fine fescues, tall fescue, perennial ryegrass, bentgrass.
- **Warm-Season Species:** Bermudagrass (common or hybrid), bahia-grass, buffalograss, centipedegrass, kikuyugrass, seashore paspalum, St. Augustinegrass (including improved varieties such as Bitterblue and Floratam), zoysiagrass.

Use Restrictions:

- Do not apply more than 8.26 lb. of product per 1000 sq.ft. per year.
- Do not retreat within 30 days of application.
- Do not use on carpetgrass, dichondra, legumes, or turfgrass where desirable clovers are present.
- Do not apply to bentgrass mowed under 1/2 inch.
- Do not collect grass clippings from treated turfgrass for use as mulch around plants.
- Do not apply directly to flowers, vegetables, groundcovers, gardens or landscape plantings including ornamental shrubs or trees.
- Do not apply this product to grass seedlings until after the second mowing.
- Do not apply this product until 3 weeks after sodding, sprigging, or plugging operations.

Use Precautions:

- Delay sodding, sprigging, seeding or plugging operations until 3 weeks after application of this product.

5. Application Directions

How Much to Apply:

Apply 2 lb. of product per 500 sq.ft. Retreatment after 30 days is allowed but do not apply more than a total of 8.26 lb. per 1000 sq.ft. per year.

When to Apply:

Apply when sedge or kyllinga plants are actively growing, typically from May through September. Retreatment may be necessary when weed pressures are high, when soil moisture is low, or when weeds are not actively growing.

Application Tips:

- Moisture on leaf surface is not required. Apply when weed leaves are wet or dry.
- Temperatures below 50°F will reduce weed control activity.
- For best results, water-in granules within 48 hours after treatment.
- For best results, postpone mowing until granules are watered-in.
- For best winter annual control (henbit, false dandelion, and chickweed) apply in early fall.

This 2 lb. canister will cover 500 sq.ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Vexis® is a registered trademark, and Sedge Stop™ is a trademark of PBI-Gordon Corporation.

842/4-2019 AP120618
EPA REG. NO. 2217-1021



MANUFACTURED FOR
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.