



U.S. Department of Health and Human Services
Food and Drug Administration
Center for Drug Evaluation and Research
Office of Pharmacoeconomics and Statistical Science
Office of Biostatistics

STATISTICAL REVIEW AND EVALUATION

CLINICAL STUDIES

NDA/Serial Number: N20-905/SE012

Drug Name: AVARATTM (leflunomide 20 mg)

Indication(s): Rheumatoid Arthritis

Applicant: Aventis Pharmaceuticals, Inc.

Date(s): Received: October 1, 2003

Review Priority: Priority Review

Biometrics Division: Division of Biometrics III (HFD-725)

Statistical Reviewer: Suktae Choi, Ph.D.

Concurring Reviewers: Stan Lin, Ph.D.

Medical Division: Division of Anti-Inflammatory, Analgesic, and Ophthalmic Drug Products (HFD-550)

Clinical Team: Carolyn Yancey, M.D.

Project Manager: Jane Dean

Keywords: Active Control/Superiority, Clinical Studies, Confidence Interval

Table of Contents

LIST OF TABLES.....	3
LIST OF FIGURES.....	3
1. EXECUTIVE SUMMARY	4
1.1 CONCLUSIONS AND RECOMMENDATIONS	4
1.2 BRIEF OVERVIEW OF CLINICAL STUDIES	4
1.3 STATISTICAL ISSUES AND FINDINGS	4
2. INTRODUCTION	6
2.1 OVERVIEW.....	6
2.2 DATA SOURCES	6
3. STATISTICAL EVALUATION	7
3.1 EVALUATION OF EFFICACY	7
3.1.1 <i>Study Design and Endpoints</i>	7
3.1.2 <i>Patient Disposition, Demographic and Baseline Characteristics</i>	8
3.1.3 <i>Statistical Methodologies</i>	8
3.1.4 <i>Results and Conclusions</i>	9
3.2 EVALUATION OF SAFETY	10
4. FINDINGS IN SPECIAL/SUBGROUP POPULATIONS	11
4.1 GENDER, RACE AND AGE	11
4.2 OTHER SPECIAL/SUBGROUP POPULATIONS	11
5. SUMMARY AND CONCLUSIONS	12
5.1 STATISTICAL ISSUES AND COLLECTIVE EVIDENCE	12
5.2 CONCLUSIONS AND RECOMMENDATIONS	12
APPENDIX	13
<u>TABLES</u>	13
<u>FIGURES</u>	19

LIST OF TABLES

Table 1. Dosage Calculations for leflunomide and leflunomide placebo	13
Table 2. Analysis population and subjects disposition	13
Table 3. Demographics of patients	13
Table 4. JRA DOI 30% Responder Rate; Primary efficacy variable, ITT	14
Table 5. Percent Improvement Index; Primary efficacy variable, ITT	14
Table 6. JRA DOI 50% responder rate; ITT	14
Table 7. JRA DOI 70% responder rate; ITT	14
Table 8. JRA DOI responder-at-endpoint rate; ITT	15
Table 9. AUC for JRA DOI 30%, 50%, 70%; ITT	15
Table 10. Global assessment by physician; Change from baseline, ITT	15
Table 11. Global assessment by patient/parent; Change from baseline, ITT	15
Table 12. Number of active joints; Change from baseline, ITT	16
Table 13. Number of limited ROM joints; Change from baseline, ITT	16
Table 14. CHAQ Disability Index; Change from baseline, ITT	16
Table 15. ESR (mm/hr); Change from baseline, ITT	16
Table 16. Pain score; Change from baseline, ITT	17
Table 17. CRP (mg/L); Change from baseline, ITT	17
Table 18. JRA DOI 30%; logistic regression results by subgroup, ITT	17
Table 19. Percent Improvement Index subgroup analyses; ITT	18

LIST OF FIGURES

Figure 1. JRA DOI 30% Responder Rate over time	19
Figure 2. % Improvement Index (LS Mean).....	19

1. EXECUTIVE SUMMARY

1.1 Conclusions and Recommendations

This NDA supplement failed to establish the efficacy of leflunomide comparing to methotrexate in treatment of JRA. The efficacy results demonstrated statistical superiority of methotrexate for DOI 30% responder rate which is one of two co-primary efficacy variables. The other co-primary efficacy variable, % Improvement Index, showed in favor of methotrexate but the difference was not statistically significant.

1.2 Brief Overview of Clinical Studies

This submission is being made to supplement the current approved NDA with pediatric data pertaining to the clinical utility of leflunomide in juvenile rheumatoid arthritis. The sponsor submitted three studies (1037, 3503, and 3504) under the pediatric program. Study 1037 was an open-label, non-controlled, five-centers, Phase IB study over a 6 month treatment period with up to a 24-month extension phase. Study 3503 was a randomized, double blind, parallel group 16-week treatment trial comparing leflunomide to methotrexate, in pediatric subjects with polyarticular course JRA who were DMARD-therapy naïve. This study was originally planned 240 subjects (120 per treatment group) for a non-inferiority design, but amended to 94 (47 per group) because of the difficulty of recruitment. Study 3504 was an eight month extension of study 3503 and still ongoing. This review will focus only on study 3503.

In study 3503, following efficacy variables were observed at screening, baseline, week 4, 8, 12, and 16: Percent Improvement Index and JRA DOI $\geq$ 30% responder status using the 6 core set measures of the JRA Definition of Improvement. Additional response assessments were time to response, DOI $\geq$ 30%, $\geq$ 50%, and $\geq$ 70% responder-at-endpoint rates, AUC, physician's global assessment, subject/parent global assessment, number of active joints, number of joints with limitation of motion plus pain and/or tenderness, functional assessment (CHAQ), ESR, CRP, and pain score.

1.3 Statistical Issues and Findings

- Methotrexate performed statistically better than leflunomide as measured by the JRA DOI $\geq$ 30% responder rate. The rate in the methotrexate group was 89.4% vs. 68.1 % in the leflunomide group. P-value is 0.0091 and 95% Confidence Interval of the difference is (-37.3%, -5.3%).

- The percent Improvement Index demonstrated no significant difference between treatment groups at week 16, LS Mean improvement was –44.41% (SE4.51) in the leflunomide group and –52.87% (SE4.39) in the methotrexate group.
- JRA DOI $\geq$ 30% responder rate was requested to add as a primary efficacy variable by agency, because this variable is one of the most commonly used efficacy variable. In fact, percent Improvement Index is rarely used as a primary efficacy variable.
- Secondary analyses showed similar results with primary analyses. All the secondary efficacy variables at week 16 showed in favor of methotrexate compared with leflunomide except CHAQ, and some of them showed significant differences. At week 4, 8, 12, and 16, majority of them showed in favor of methotrexate.
- Since both primary efficacy variables showed in favor of methotrexate, and one of them showed significant difference, we cannot conclude that the efficacy of leflunomide is as good as the efficacy of methotrexate in this study.

2. INTRODUCTION

2.1 Overview

This submission is being made to supplement the current approved NDA with pediatric data pertaining to the clinical utility of leflunomide in juvenile rheumatoid arthritis. The sponsor submitted three studies (1037, 3503, and 3504) under the pediatric program. Study 1037 was an open-label, non-controlled, five-centers, Phase IB study over a 6 month treatment period with up to a 24-month extension phase. Study 3503 was a randomized, double blind, parallel group 16-week treatment trial comparing leflunomide to methotrexate, in pediatric subjects with polyarticular course JRA who were DMARD-therapy naïve. This study was originally planned 240 subjects (120 per treatment group) for a non-inferiority design, but amended to 94 (47 per group) because of the difficulty of recruitment. Study 3504 was an eight month extension of study 3503 and still ongoing. This review will focus only on study 3503.

2.2 Data Sources

Hard Copies: Volume 1 through 7 submitted 10/1/2003

Electric files: \\CDSESUB1\N20905\S_012\2003-09-30

3. STATISTICAL EVALUATION

3.1 Evaluation of Efficacy

3.1.1 Study Design and Endpoints

The study was a multinational, multicenter, two arms, double-blind, double-dummy, randomized, parallel, and active controlled study. Duration was 16 weeks. Among 103 patients screened, 94 were randomized (47 per each group). Patients were between the ages of 3-17 years. Visits were at week 4, 8, 12 and 16.

Dosage schedule

Randomized to leflunomide: each subject received a leflunomide loading dose ranging from one 100 mg tablet /day for 1 day to one 100 mg tablet /day for 3 consecutive days, depending on body weight. Thereafter, subjects received a maintenance dose of 10 mg every other day, 10 mg daily, or two-10 mg tablets daily (20 mg daily), depending on weight. Detail of dosage schedule is summarized in Table 1 of appendix. Subjects also received methotrexate placebo tablets weekly based on body weight.

Randomized to methotrexate: each subject received methotrexate 2.5 mg tablets weekly, based on body weight, for a dose of 0.5 mg/kg/wk to a maximum of 25 mg/wk. If the calculated methotrexate dose was not a multiple of 2.5 mg, the subject was dosed at the closest whole number of methotrexate tablets. Subjects also received leflunomide placebo.

For those children who were unable to swallow a tablet, tablets were permitted to be crushed and mixed in applesauce or jam.

Efficacy data

Primary efficacy variables:

1. JRA DOI $\geq$ 30% responder rate at week 16
2. Percent Improvement Index at week 16

Percent Improvement Index is defined as follow:

- This variable is based on the JRA DOI's 6 core set measures.
- For each subject, the % Improvement Index was the mean of the 6 core set percent changes from baseline.

The percent change from baseline to end of treatment was calculated as follows:

$(\text{value at end of treatment} - \text{value at baseline}) / \text{value at baseline} \times 100$

- In the event that the mean percent change was positive (worsened), then the % Improvement Index for that subject was set to zero.

Secondary efficacy variables:

1. JRA DOI $\geq 50\%$ and $\geq 70\%$ responder rates
2. JRA DOI $\geq 30\%$, $\geq 50\%$, and $\geq 70\%$ responder-at-endpoint rates (this variable considers non-completers as not responders)
3. AUC of DOI $\geq 30\%$, $\geq 50\%$, and $\geq 70\%$ responses: Months of response
4. Time to JRA DOI 30% response
5. Physician global assessment of disease activity
6. Patient/parent global assessment of disease activity
7. Number of active joints
8. Joints with limited range of motion
9. CHAQ disability index
10. Erythrocyte sedimentation rate (ESR)
11. Pain score
12. C-reactive protein (CRP)

3.1.2 Patient Disposition, Demographic and Baseline Characteristics

As shown in Table 2 and Table 3 of appendix, two treatment groups are similar in disposition and in key demographic characteristics. Primary disease were compared between two treatment groups at baseline by sponsor, and the variables were JRA type at onset, JRA duration, Active joint count, Limited ROM joint count, MD global assessment score, Patient global assessment, Disability index, ESR, CRP, and Pain score, but none of them showed significant difference.

3.1.3 Statistical Methodologies

The following inferential null hypothesis was tested:

- H_0 : no treatment difference between leflunomide and methotrexate for JRA DOI 30% Responder rate at endpoint (or mean % Improvement Index).
 H_1 : treatment difference between leflunomide and methotrexate for JRA DOI 30% Responder rate at endpoint (or mean % Improvement Index).

The null hypothesis H_0 will be tested against the alternative H_1 two-sided with $\alpha=0.05$. Since both comparisons have to show significant difference, multiple comparison adjustment is not necessary.

For the analysis for JRA DOI 30% Responder at Endpoint, the difference of responder rates of treatment groups was supposed to be compared using the normal approximation in

statistical analysis plan (Appendix B of sponsor's NDA submission). However, in the sponsor's NDA final report, CMH method was used to calculate p-values, which was not specified in the statistical analysis plan. The p-values using the protocol specified method were calculated by this reviewer and replaced with CMH p-values in this review, because the primary analysis must be the one specified in the protocol.

For the analysis of % Improvement Index, ANOVA was used on the mean % Improvement Index with treatment and country as fixed effects. This was specified in the statistical analysis plan.

For secondary analyses, 95% CI of responder rate difference between treatment groups using normal approximation was used for binary variables (p-values are correspondent to this CI), and ANCOVA with factors of treatment and baseline was used for changes from baseline continuous variables.

ITT was used in efficacy analyses for primary population, and LOCF was used as an imputation method for early dropout for all the efficacy analysis as specified in the protocol.

3.1.4 Results and Conclusions

- Methotrexate performed statistically better than leflunomide as measured by the JRA DOI $\geq 30\%$ responder rate. The rate in the methotrexate group was 89.4% vs. 68.1 % in the leflunomide group. P-value is 0.0091 and 95% Confidence Interval of the difference is (-37.3%, -5.3%). The comparison results during the study period are summarized in Table 4 and Figure 1.
- The percent Improvement Index demonstrated no significant difference between treatment groups at week 16, LS Mean improvement was -44.41% (SE4.51) in the leflunomide group and -52.87% (SE4.39) in the methotrexate group. The comparison results during the study period are summarized in Table 5 and Figure 2.
- JRA DOI $\geq 30\%$ responder rate was requested to add as a primary efficacy variable by agency, because this variable is one of the most commonly used efficacy variable. In fact, percent Improvement Index is rarely used as a primary efficacy variable.
- Secondary analyses showed similar with primary analysis results. All the secondary efficacy variables at week 16 showed in favor of methotrexate compared with leflunomide except CHAQ, and some of them showed significant differences. At week 4, 8, 12, and 16, majority of them showed in favor of methotrexate. Details of secondary analysis results are summarized in Table 6 to Table 17 of appendix.

- Since both primary efficacy variables showed in favor of methotrexate, and one of them showed significant difference, we cannot conclude that the efficacy of leflunomide is as good as the efficacy of methotrexate in this study.

3.2 Evaluation of Safety

Safety data were not reviewed.

4. FINDINGS IN SPECIAL/SUBGROUP POPULATIONS

4.1 Gender, Race and Age

Subgroup analysis results for gender and age are summarized in Table 18 and Table 19 for JRA DOI 30% and percent Improvement Index, respectively. Race was not included because most of RA patients are white. Since these subgroup analyses were not planned in the protocol, CMH method is acceptable for analysis of DOI 30%. As shown, there is no significant interaction between subgroup and treatment group.

4.2 Other Special/Subgroup Populations

Subgroup analysis result for Body weight is summarized in . As shown, there is no significant interaction between subgroup and treatment group.

5. SUMMARY AND CONCLUSIONS

5.1 Statistical Issues and Collective Evidence

- Methotrexate performed statistically better than leflunomide as measured by the JRA DOI $\geq 30\%$ responder rate. The rate in the methotrexate group was 89.4% vs. 68.1 % in the leflunomide group. P-value is 0.0091 and 95% Confidence Interval of the difference is (-37.3%, -5.3%).
- The percent Improvement Index demonstrated no significant difference between treatment groups at week 16, LS Mean improvement was -44.41% (SE4.51) in the leflunomide group and -52.87% (SE4.39) in the methotrexate group.
- JRA DOI $\geq 30\%$ responder rate was requested to add as a primary efficacy variable by agency, because this variable is one of the most commonly used efficacy variable. In fact, percent Improvement Index is rarely used as a primary efficacy variable.
- Secondary analyses showed similar results with primary analyses. All the secondary efficacy variables at week 16 showed in favor of methotrexate compared with leflunomide except CHAQ, and some of them showed significant differences. At week 4, 8, 12, and 16, majority of them showed in favor of methotrexate.
- Since both primary efficacy variables showed in favor of methotrexate, and one of them showed significant difference, we cannot conclude that the efficacy of leflunomide is as good as the efficacy of methotrexate in this study.

5.2 Conclusions and Recommendations

This NDA supplement failed to establish the efficacy of leflunomide comparing to methotrexate in treatment of JRA. The efficacy results demonstrated statistical superiority of methotrexate for DOI 30% responder rate which is one of two co-primary efficacy variables. The other co-primary efficacy variable, % Improvement Index, showed in favor of methotrexate but the difference was not statistically significant.

APPENDIX

Tables

Table 1. Dosage Calculations for leflunomide and leflunomide placebo

Subject's Weight (kg)	Loading dose	Loading dose mg/day	Maintenance dose mg/day
< 20	1 x 100 mg	1 tablet (100 mg) on one day	Average of 5 mg (one 10 mg tablet every other day) for 16 wks
20 -40	2 x 100 mg	1 tablet (100 mg) per day for two consecutive days	10 mg (one 10 mg tablet daily) for 16 wks
> 40	3 x 100 mg	1 tablet (100 mg) per day for three consecutive days	20 mg (two 10 mg tablets daily) for 16 wks

Table 2. Analysis population and subjects disposition

Population	Leflunomide	Methotrexate	Total
Screened			103
Randomized	47	47	94
Safety	47	47	94
Intent to Treat	47	47	94
Completer	42 (89.4%)	44 (93.6%)	86 (91.5%)
Reason for discontinuation			
Adverse Event	3 (6.4%)	1 (2.1%)	4 (4.3%)
Lack of efficacy	1 (2.1%)	1(2.1%)	2 (2.1%)
Lost to follow-up	0	1 (2.1%)	1 (1.1%)
Other	1 (2.1%)	0	1 (1.1%)

Table 3. Demographics of patients

Demographic	Treatment Group		P-value
	leflunomide (N=47)	Methotrexate (N=47)	
Age (years)			
Mean (SD)	10.1 (4.0)	10.2 (3.8)	0.9310
Median	11	11	
Range	3 – 17	3 – 17	
Age group			
< 12 years	27 (57.4%)	27 (57.4%)	0.9495
≥ 12 years	20 (42.6%)	20 (42.6%)	
Sex			
Male	12 (25.5%)	13 (27.7%)	0.6930
Female	35 (74.5%)	34 (72.3%)	

Table 4. JRA DOI 30% Responder Rate; Primary efficacy variable, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	responder / ITT	%	responder / ITT	%	%	95% CI	
4	22/44	50.0	17/42	40.5	9.5	(-11.4, 30.5)	0.3728 ^{lef}
8	29/47	61.7	32/47	68.1	-6.4	(-25.7, 12.9)	0.5159 ^{mtx}
12	32/47	68.1	40/47	85.1	-17.0	(-33.8, -0.2)	0.0467 ^{mtx*}
16	32/47	68.1	42/47	89.4	-21.3	(-37.3, -5.3)	0.0091 ^{mtx*}

a: Z-test using difference of rates with normal approximation

lef: favor of leflunomide in rate

mtx: favor of methotrexate in rate

* : statistical significant (<0.05)

Table 5. Percent Improvement Index; Primary efficacy variable, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	44	-25.56 (3.817)	42	-26.62 (3.837)	1.06	(-9.27, 11.39)	0.8388 ^{mtx}
8	47	-31.26 (3.941)	47	-35.51 (3.843)	4.25	(-6.51, 15.01)	0.4343 ^{mtx}
12	47	-38.63 (4.311)	47	-44.85 (4.203)	6.22	(-5.55, 17.98)	0.2966 ^{mtx}
16	47	-44.41 (4.513)	47	-52.87 (4.399)	8.46	(-3.86, 20.77)	0.1758 ^{mtx}

a: ANOVA with factors of treatment group and country

mtx: favor of methotrexate in rate

Table 6. JRA DOI 50% responder rate; ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	responder / ITT	%	responder / ITT	%	%	95% CI	
4	17/44	38.6	11/42	26.2	12.4	(-7.2, 32.1)	0.2131 ^{lef}
8	20/47	42.6	19/47	40.4	2.1	(-17.8, 22.1)	0.8341 ^{lef}
12	26/47	55.3	31/47	66.0	-10.6	(-30.3, 9.0)	0.2883 ^{mtx}
16	28/47	59.6	36/47	76.6	-17.0	(-35.6, 1.5)	0.0718 ^{mtx}

a: Z-test using difference of rates with normal approximation

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 7. JRA DOI 70% responder rate; ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	responder / ITT	%	responder / ITT	%	%	95% CI	
4	4/44	9.1	4/42	9.5	-0.4	(-12.7, 11.9)	0.9449 ^{mtx}
8	11/47	23.4	9/47	19.1	4.3	(-12.3, 20.8)	0.6138 ^{lef}
12	18/47	38.3	19/47	40.4	-2.1	(-21.9, 17.6)	0.8328 ^{mtx}
16	20/47	42.6	28/47	59.6	-17.0	(-37.0, 2.9)	0.0939 ^{mtx}

a: Z-test using difference of rates with normal approximation

lef: favor of leflunomide in rate

mtx: favor of methotrexate in rate

Table 8. JRA DOI responder-at-endpoint rate; ITT

Respond- er %	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	responder / ITT	%	responder / ITT	%	%	95% CI	
30%	30/47	63.8	39/47	83.0	-19.1	(-36.6, -1.7)	0.0314 ^{mtx*}
50%	26/47	55.3	35/47	74.5	-19.1	(-38.1, -0.2)	0.0471 ^{mtx*}
70%	18/47	38.3	28/47	59.6	-21.3	(-41.0, -1.5)	0.0347 ^{mtx*}

a: Z-test using difference of rates with normal approximation

mtx: favor of methotrexate in rate

* : statistical significant (<0.05)

Table 9. AUC for JRA DOI 30%, 50%, 70%; ITT

Respond- er %	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
30%	47	1.86 (0.171)	47	2.12 (0.17)	-0.26	(-0.73, 0.20)	0.2670 ^{mtx}
50%	47	1.51 (0.185)	47	1.57 (0.180)	-0.06	(-0.57, 0.44)	0.8021 ^{mtx}
70%	47	0.88 (0.169)	47	0.92 (0.165)	-0.04	(-0.50, 0.42)	0.8665 ^{mtx}

a: Z-test using difference of rates with normal approximation

mtx: favor of methotrexate in LS mean

Table 10. Global assessment by physician; Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-17.5 (2.42)	45	-18.5 (2.41)	1.0	(-5.7, 7.6)	0.7736 ^{mtx}
8	47	-23.7 (2.64)	47	-21.8 (2.60)	-2.0	(-9.2, 5.3)	0.5951 ^{lef}
12	47	-27.6 (2.65)	47	-28.6 (2.61)	1.1	(-6.2, 8.3)	0.7754 ^{mtx}
16	47	-31.5 (2.98)	47	-32.1 (2.94)	0.6	(-7.6, 8.8)	0.8884 ^{mtx}

a: ANOVA with factors of treatment group and site

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 11. Global assessment by patient/parent; Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-13.2 (3.50)	44	-5.6 (3.56)	-7.6	(-17.2, 2.0)	0.1177 ^{lef}
8	47	-13.5 (3.06)	47	-15.3 (2.98)	1.8	(-6.5, 10.1)	0.6682 ^{mtx}
12	47	-15.7 (2.94)	47	-19.4 (2.87)	3.7	(-4.3, 11.7)	0.3599 ^{mtx}
16	47	-15.9 (2.97)	47	-22.0 (2.89)	6.1	(-2.0, 14.2)	0.1359 ^{mtx}

a: ANOVA with factors of treatment group and site

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 12. Number of active joints; Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-4.1 (0.90)	45	-3.7 (0.89)	-0.4	(-2.9, 2.1)	0.7459 ^{lef}
8	47	-6.0 (1.05)	47	-5.9 (1.03)	-0.2	(-3.0, 2.7)	0.9054 ^{lef}
12	47	-7.0 (0.98)	47	-8.2 (0.96)	1.2	(-1.5, 3.8)	0.3844 ^{mtx}
16	47	-8.1 (0.99)	47	-8.9 (0.96)	0.8	(-1.9, 3.5)	0.5671 ^{mtx}

a: ANOVA with factors of treatment group and site

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 13. Number of limited ROM joints; Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-2.9 (0.82)	45	-2.5 (0.81)	-0.4	(-2.6, 1.9)	0.7322 ^{lef}
8	47	-4.4 (0.61)	47	-4.5 (0.60)	0.1	(-1.6, 1.7)	0.9461 ^{mtx}
12	47	-5.2 (0.70)	47	-4.6 (0.69)	-0.5	(-2.4, 1.4)	0.6042 ^{lef}
16	47	-5.2 (0.81)	47	-5.3 (0.79)	0.1	(-2.1, 2.3)	0.9157 ^{mtx}

a: ANOVA with factors of treatment group and site

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 14. CHAQ Disability Index; Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-0.18 (0.068)	45	-0.16 (0.067)	-0.02	(-0.21, 0.16)	0.8248 ^{lef}
8	47	-0.25 (0.068)	47	-0.28 (0.066)	0.02	(-0.16, 0.21)	0.7936 ^{mtx}
12	47	-0.34 (0.069)	47	-0.32 (0.067)	-0.02	(-0.21, 0.17)	0.8479 ^{lef}
16	47	-0.44 (0.075)	47	-0.39 (0.073)	-0.05	(-0.24, 0.15)	0.6060 ^{lef}

a: ANOVA with factors of treatment group and site

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 15. ESR (mm/hr); Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-1.7 (2.02)	45	-6.2 (1.90)	4.5	(-0.8, 9.8)	0.0974 ^{mtx}
8	47	-3.4 (1.58)	47	-6.2 (1.49)	2.9	(-1.4, 7.1)	0.1832 ^{mtx}
12	47	-4.3 (1.63)	47	-6.7 (1.54)	2.5	(-1.9, 6.8)	0.2676 ^{mtx}
16	47	-6.5 (1.28)	47	-7.2 (1.20)	0.8	(-2.7, 4.2)	0.6588 ^{mtx}

a: ANOVA with factors of treatment group and site

mtx: favor of methotrexate in LS mean

Table 16. Pain score; Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-13.1 (3.47)	45	-9.1 (3.42)	-4.1	(-13.5, 5.4)	0.3940 ^{lef}
8	47	-14.9 (2.96)	47	-17.6 (2.87)	2.6	(-5.4, 10.7)	0.5180 ^{mtx}
12	47	-18.0 (3.39)	47	-18.3 (3.30)	0.3	(-8.9, 9.6)	0.9416 ^{mtx}
16	47	-15.9 (3.21)	47	-23.0 (3.12)	7.1	(-1.7, 15.8)	0.1109 ^{mtx}

a: ANOVA with factors of treatment group and site

lef: favor of leflunomide in LS mean

mtx: favor of methotrexate in LS mean

Table 17. CRP (mg/L); Change from baseline, ITT

Visit Week	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
4	45	-4.29 (2.28)	44	-9.35 (2.26)	5.06	(-1.19, 11.31)	0.1109 ^{mtx}
8	47	-2.23 (2.25)	47	-10.45 (2.24)	8.22	(1.99, 14.45)	0.0103 ^{mtx*}
12	47	-5.90 (2.73)	47	-11.24 (2.68)	5.34	(-2.17, 12.84)	0.1608 ^{mtx}
16	47	-3.86 (2.65)	47	-11.43 (2.60)	7.56	(0.26, 14.87)	0.0425 ^{mtx*}

a: ANOVA with factors of treatment group and site

mtx: favor of methotrexate in LS mean

*: statistical significant (<0.05)

Table 18. JRA DOI 30%; logistic regression results by subgroup, ITT

Sub-group	Leflunomide		Methotrexate		Difference Lef - Mtx		P-value ^a
	responder / ITT	%	responder / ITT	%	odds ratio	95% CI	
Sex							
Male	8/12	66.7	12/13	92.3	0.57	(0.04, 8.6)	0.6876
Female	24/35	68.6	30/34	88.2			
Age							
<12 years	18/27	66.7	25/27	92.6	0.37	(0.04, 3.70)	0.3989
≥ 12 years	14/20	70.0	17/20	85.0			
Weight							
≤ 40 kg	16/27	59.3	19/21	90.5	0.24	(0.02, 2.60)	0.2387
> 40 kg	16/20	80.0	23/26	88.5			

a. p-value for subgroup interaction

Table 19. Percent Improvement Index subgroup analyses; ITT

Sub-group	Leflunomide		Methotrexate		Difference Lef – Mtx		P-value ^a
	N	LS Mean (SE)	N	LS Mean (SE)	LS Mean	95% CI	
Sex							
Male	12	-48.67 (8.78)	13	-53.64 (8.35)	4.97	(-19.1, 29.0)	0.7456
Female	35	-43.06 (5.17)	34	-52.64 (5.19)	9.58	(-5.0, 24.2)	
Age							
<12 years	27	-44.82 (5.84)	27	-57.50 (5.64)	12.68	(-3.5, 28.9)	0.4224
≥ 12 years	20	-42.96 (6.88)	20	-45.76 (6.92)	2.81	(-15.7, 21.3)	
Weight							
≤ 20 kg	8	-46.29 (11.55)	8	-66.92 (10.59)	20.63	(-10.3, 51.5)	0.6626
20-40 kg	19	-41.83 (7.06)	13	-49.45 (8.32)	7.63	(-14.5, 29.8)	
> 40 kg	20	-46.25 (6.93)	26	-50.86 (6.10)	4.61	(-12.7, 22.0)	

a. p-value for subgroup interaction

Figures

Figure 1. JRA DOI 30% Responder Rate over time





Figure 2. % Improvement Index (LS Mean)



**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Suktae Choi
3/5/04 02:21:17 PM
BIOMETRICS

Stan Lin
3/5/04 03:58:34 PM
UNKNOWN