



(11) **EP 2 336 318 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**22.06.2011 Bulletin 2011/25**

(51) Int Cl.:  
**C12N 15/113<sup>(2010.01)</sup> A61K 48/00<sup>(2006.01)</sup>**  
**C07H 21/00<sup>(2006.01)</sup>**

(21) Application number: **10180483.9**

(22) Date of filing: **13.11.2003**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PT RO SE SI SK TR**

(30) Priority: **13.11.2002 US 426234 P**  
**15.05.2003 PCT/US03/15493**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:  
**03789763.4 / 1 569 695**

(71) Applicant: **Genzyme Corporation**  
**Cambridge, MA 02142 (US)**

(72) Inventors:  
• **Crooke, Rosanne**  
**Carlsbad, CA 92009 (US)**  
• **Graham, Mark**  
**San Clemente, CA 92672 (US)**

• **Lemonidis-Tarbet, Kristina**  
**Oceanside, CA 92057 (US)**  
• **Dobie, Kenneth W.**  
**Delmar, CA 92014 (US)**  
• **Freier, Susan M.**  
**San Diego, California 92122 (US)**

(74) Representative: **Hallybone, Huw George**  
**Carpmaels & Ransford**  
**One Southampton Row**  
**London**  
**WC1B 5HA (GB)**

Remarks:

This application was filed on 27-09-2010 as a divisional application to the application mentioned under INID code 62.

(54) **Antisense modulation of apolipoprotein B expression**

(57) Antisense compounds, compositions and methods are provided for modulating the expression of apolipoprotein B. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding apolipoprotein B. Methods of

using these compounds for modulation of apolipoprotein B expression and for treatment of diseases associated with expression of apolipoprotein B are provided.

**EP 2 336 318 A1**

**Description**

**[0001]** This application is a continuation-in-part of PCT application US03/15493, filed on May 15, 2003, which claims priority to U.S. provisional Application Serial No: 60/426,234, filed November 13, 2002, and which is a continuation-in-part of U.S. Application Serial No. 10/147,196 filed May 15, 2002 (Attorney Docket No. ISPH-0664) which is a continuation-in-part of U.S. Application Serial No. 10/135,985 filed April 30, 2002 (Attorney Docket No. ISPH-0663) which is a continuation-in-part of U.S. Application Serial No. 09/920,033 filed August 1, 2001 (Attorney Docket No ISPH-0592).

**FIELD OF THE INVENTION**

**[0002]** The present invention provides compositions and methods for modulating the expression of apolipoprotein B. In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding apolipoprotein B. Such compounds have been shown to modulate the expression of apolipoprotein B.

**BACKGROUND OF THE INVENTION**

**[0003]** Lipoproteins are globular, micelle-like particles that consist of a non-polar core of acylglycerols and cholesteryl esters surrounded by an amphiphilic coating of protein, phospholipid and cholesterol. Lipoproteins have been classified into five broad categories on the basis of their functional and physical properties: chylomicrons, which transport dietary lipids from intestine to tissues; very low density lipoproteins (VLDL); intermediate density lipoproteins (IDL); low density lipoproteins (LDL); all of which transport triacylglycerols and cholesterol from the liver to tissues; and high density lipoproteins (HDL), which transport endogenous cholesterol from tissues to the liver.

**[0004]** Lipoprotein particles undergo continuous metabolic processing and have variable properties and compositions. Lipoprotein densities increase without decreasing particle diameter because the density of their outer coatings is less than that of the inner core. The protein components of lipoproteins are known as apolipoproteins. At least nine apolipoproteins are distributed in significant amounts among the various human lipoproteins.

**[0005]** Apolipoprotein B (also known as ApoB, apolipoprotein B-100; ApoB-100, apolipoprotein B-48; ApoB-48 and Ag(x) antigen), is a large glycoprotein that serves an indispensable role in the assembly and secretion of lipids and in the transport and receptor-mediated uptake and delivery of distinct classes of lipoproteins. The importance of apolipoprotein B spans a variety of functions, from the absorption and processing of dietary lipids to the regulation of circulating lipoprotein levels (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193). This latter property underlies its relevance in terms of atherosclerosis susceptibility, which is highly correlated with the ambient concentration of apolipoprotein B-containing lipoproteins (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193).

**[0006]** Two forms of apolipoprotein B exist in mammals. ApoB-100 represents the full-length protein containing 4536 amino acid residues synthesized exclusively in the human liver (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193). A truncated form known as ApoB-48 is colinear with the amino terminal 2152 residues and is synthesized in the small intestine of all mammals (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193).

**[0007]** ApoB-100 is the major protein component of LDL and contains the domain required for interaction of this lipoprotein species with the LDL receptor. In addition, ApoB-100 contains an unpaired cysteine residue which mediates an interaction with apolipoprotein(a) and generates another distinct atherogenic lipoprotein called Lp(a) (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193).

**[0008]** In humans, ApoB-48 circulates in association with chylomicrons and chylomicron remnants and these particles are cleared by a distinct receptor known as the LDL-receptor-related protein (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193). ApoB-48 can be viewed as a crucial adaptation by which dietary lipid is delivered from the small intestine to the liver, while ApoB-100 participates in the transport and delivery of endogenous plasma cholesterol (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193).

**[0009]** The basis by which the common structural gene for apolipoprotein B produces two distinct protein isoforms is a process known as RNA editing. A site specific cytosine-to-uracil editing reaction produces a UAA stop codon and translational termination of apolipoprotein B to produce ApoB-48 (Davidson and Shelness, *Annu. Rev. Nutr.*, 2000, 20, 169-193).

**[0010]** Apolipoprotein B was cloned in 1985 (Law et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1985, 82, 8340-8344) and mapped to chromosome 2p23-2p24 in 1986 (Deeb et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1986, 83, 419-422).

**[0011]** Disclosed and claimed in US patent 5,786,206 are methods and compositions for determining the level of low density lipoproteins (LDL) in plasma which include isolated DNA sequences encoding epitope regions of apolipoprotein B-100 (Smith et al., **1998**).

**[0012]** Transgenic mice expressing human apolipoprotein B and fed a high-fat diet were found to develop high plasma cholesterol levels and displayed an 11-fold increase in atherosclerotic lesions over non-transgenic littermates (Kim and Young, *J. Lipid Res.*, 1998, 39, 703-723; Nishina et al., *J. Lipid Res.*, 1990, 31, 859-869).

**[0013]** In addition, transgenic mice expressing truncated forms of human apolipoprotein B have been employed to identify the carboxyl-terminal structural features of ApoB-100 that are required for interactions with apolipoprotein(a) to generate the Lp(a) lipoprotein particle and to investigate structural features of the LDL receptor-binding region of ApoB-100 (Kim and Young, *J. Lipid Res.*, 1998, 39, 703-723; McCormick et al., *J. Biol. Chem.*, 1997, 272, 23616-23622).

**[0014]** Apolipoprotein B knockout mice (bearing disruptions of both ApoB-100 and ApoB-48) have been generated which are protected from developing hypercholesterolemia when fed a high-fat diet (Farese et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1995, 92, 1774-1778; Kim and Young, *J. Lipid Res.*, 1998, 39, 703-723). The incidence of atherosclerosis has been investigated in mice expressing exclusively ApoB-100 or ApoB-48 and susceptibility to atherosclerosis was found to be dependent on total cholesterol levels. Whether the mice synthesized ApoB-100 or ApoB-48 did not affect the extent of the atherosclerosis, indicating that there is probably no major difference in the intrinsic atherogenicity of ApoB-100 versus ApoB-48 (Kim and Young, *J. Lipid Res.*, 1998, 39, 703-723; Veniant et al., *J. Clin. Invest.*, 1997, 100, 180-188).

**[0015]** Elevated plasma levels of the ApoB-100-containing lipoprotein Lp(a) are associated with increased risk for atherosclerosis and its manifestations, which may include hypercholesterolemia (Seed et al., *N. Engl. J. Med.*, 1990, 322, 1494-1499), myocardial infarction (Sandkamp et al., *Clin. Chem.*, 1990, 36, 20-23), and thrombosis (Nowak-Gottl et al., *Pediatrics*, 1997, 99, E11).

**[0016]** The plasma concentration of Lp(a) is strongly influenced by heritable factors and is refractory to most drug and dietary manipulation (Katan and Beynen, *Am. J. Epidemiol.*, 1987, 125, 387-399; Vessby et al., *Atherosclerosis*, 1982, 44, 61-71). Pharmacologic therapy of elevated Lp(a) levels has been only modestly successful and apheresis remains the most effective therapeutic modality (Hajjar and Nachman, *Annu. Rev. Med.*, 1996, 47, 423-442).

**[0017]** Disclosed and claimed in US patent 6,156,315 and the corresponding PCT publication WO 99/18986 is a method for inhibiting the binding of LDL to blood vessel matrix in a subject, comprising administering to the subject an effective amount of an antibody or a fragment thereof, which is capable of binding to the amino-terminal region of apolipoprotein B, thereby inhibiting the binding of low density lipoprotein to blood vessel matrix (Goldberg and Pillarisetti, **2000**; Goldberg and Pillarisetti, **1999**).

**[0018]** Disclosed and claimed in US patent 6,096,516 are vectors containing cDNA encoding murine recombinant antibodies which bind to human ApoB-100 for the purpose of for diagnosis and treatment of cardiovascular diseases (Kwak et al., **2000**).

**[0019]** Disclosed and claimed in European patent application EP 911344 published April 28, 1999 (and corresponding to U.S. Patent 6,309,844) is a monoclonal antibody which specifically binds to ApoB-48 and does not specifically bind to ApoB-100, which is useful for diagnosis and therapy of hyperlipidemia and arterial sclerosis (Uchida and Kurano, **1998**).

**[0020]** Disclosed and claimed in PCT publication WO 01/30354 are methods of treating a patient with a cardiovascular disorder, comprising administering a therapeutically effective amount of a compound to said patient, wherein said compound acts for a period of time to lower plasma concentrations of apolipoprotein B or apolipoprotein B-containing lipoproteins by stimulating a pathway for apolipoprotein B degradation (Fisher and Williams, **2001**).

**[0021]** Disclosed and claimed in US patent 5,220,006 is a cloned cis-acting DNA sequence that mediates the suppression of atherogenic apolipoprotein B (Ross et al., **1993**).

**[0022]** Disclosed and claimed in PCT publication WO 01/12789 is a ribozyme which cleaves ApoB-100 mRNA specifically at position 6679 (Chan et al., **2001**).

**[0023]** To date, strategies aimed at inhibiting apolipoprotein B function have been limited to Lp(a) apheresis, antibodies, antibody fragments and ribozymes. However, with the exception of Lp(a) apheresis, these investigative strategies are untested as therapeutic protocols. Consequently, there remains a long felt need for additional agents capable of effectively inhibiting apolipoprotein B function.

**[0024]** Antisense technology is emerging as an effective means of reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic and research applications involving modulation of apolipoprotein B expression.

**[0025]** The present invention provides compositions and methods for modulating apolipoprotein B expression, including inhibition of the alternative isoform of apolipoprotein B, ApoB-48.

## SUMMARY OF THE INVENTION

**[0026]** The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding apolipoprotein B, and which modulate the expression of apolipoprotein B. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of apolipoprotein B in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of apolipoprotein B by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

**[0027]** In particular, the invention provides a compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding apolipoprotein B, wherein said compound specifically hybridizes with and inhibits the expression of a nucleic acid molecule encoding apolipoprotein B, said compound comprising at least 8 contiguous nucleobases of any one of SEQ ID NOs: 127-134, 136, 138-174, 176-317, 319-321, 323-333, 335-339, 341-374, 376-416, 418-500, 502-510, 512-804, 815, 816, 819-821, 824, 825, 827, 828, 830, 831, 833-835, 837-839, 842, 843, and 845-854.

**[0028]** The invention further provides compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding apolipoprotein B, said compound comprising at least 8 contiguous nucleobases of any one of SEQ ID NOs: 127-134, 136, 138-174, 176-317, 319-321, 323-333, 335-339, 341-374, 376-416, 418-500, 502-510, 512-804, 815, 816, 819-821, 824, 825, 827, 828, 830, 831, 833-835, 837-839, 842, 843, and 845-854, said active site being a region in said nucleic acid wherein binding of said compound to said site significantly inhibits apolipoprotein B expression as compared to a control.

**[0029]** The invention also provides a compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding apolipoprotein B, wherein said compound specifically hybridizes with said nucleic acid and inhibits expression of apolipoprotein B, wherein the apolipoprotein B is encoded by a polynucleotide selected from the group consisting of: (a) SEQ ID NO: 3 and (b) a naturally occurring variant apolipoprotein B-encoding polynucleotide that hybridizes to the complement of the polynucleotide of (a) under stringent conditions, said compound comprising at least 8 contiguous nucleobases of any one of SEQ ID NOs: 127-134, 136, 138-174, 176-317, 319-321, 323-333, 335-339, 341-374, 376-416, 418-500, 502-510, 512-804, 815, 816, 819-821, 824, 825, 827, 828, 830, 831, 833-835, 837-839, 842, 843, and 845-854.

**[0030]** In another aspect the invention provides a compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding apolipoprotein B, wherein said compound specifically hybridizes with said nucleic acid and inhibits expression of apolipoprotein B, wherein the apolipoprotein B is encoded by a polynucleotide selected from the group consisting of SEQ ID NO: 3 and SEQ ID NO: 17, said compound comprising at least 8 contiguous nucleobases of any one of SEQ ID NOs: 127-134, 136, 138-174, 176-317, 319-321, 323-333, 335-339, 341-374, 376-416, 418-500, 502-510, 512-804, 815, 816, 819-821, 824, 825, 827, 828, 830, 831, 833-835, 837-839, 842, 843, and 845-854.

**[0031]** The invention also provides a compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding apolipoprotein B, wherein said compound specifically hybridizes with an active site in said nucleic acid and inhibits expression of apolipoprotein B, said compound comprising at least 8 contiguous nucleobases of any one of SEQ ID NOs: 127-134, 136, 138-174, 176-317, 319-321, 323-333, 335-339, 341-374, 376-416, 418-500, 502-510, 512-804, 815, 816, 819-821, 824, 825, 827, 828, 830, 831, 833-835, 837-839, 842, 843, and 845-854, said active site being a region in said nucleic acid wherein binding of said compound to said site significantly inhibits apolipoprotein B expression as compared to a control.

**[0032]** In another aspect the invention provides an oligonucleotide mimetic compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding apolipoprotein B, wherein said compound specifically hybridizes with said nucleic acid and inhibits expression of apolipoprotein B, said compound comprising at least 8 contiguous nucleobases of any one of SEQ ID NOs: 127-134, 136, 138-174, 176-317, 319-321, 323-333, 335-339, 341-374, 376-416, 418-500, 502-510, 512-804, 815, 816, 819-821, 824, 825, 827, 828, 830, 831, 833-835, 837-839, 842, 843, and 845-854.

**[0033]** In another aspect, the invention provides an antisense compound 8 to 50 nucleobases in length, wherein said compound specifically hybridizes with nucleotides 2920-3420 as set forth in SEQ ID NO:3 and inhibits expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM. In preferred embodiments, the antisense compound 8 to 50 nucleobases in length specifically hybridizes with nucleotides 3230-3288 as set forth in SEQ ID NO:3 and inhibits expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM. In another aspect, the compounds inhibits expression of mRNA encoding apolipoprotein B by at least 50%, after 16 to 24 hours in 80% confluent HepG2 cells in culture at a concentration of 150 nM.

**[0034]** In one aspect, the compounds of the invention are targeted to a nucleic acid molecule encoding apolipoprotein B, wherein said compound specifically hybridizes with and inhibits expression of the long form of apolipoprotein B, ApoB-100. In another aspect, the compounds specifically hybridizes with said nucleic acid and inhibits expression of mRNA encoding apolipoprotein B by at least 5% in 80% confluent HepG2 cells in culture at an optimum concentration. In yet another aspect, the compounds inhibits expression of mRNA encoding apolipoprotein B by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, or at least 50%.

**[0035]** In one aspect, the compounds are antisense oligonucleotides, and in one embodiment the compound has a sequence comprising SEQ ID NO: 224, the antisense oligonucleotide hybridizes with a region complementary to SEQ ID NO: 224, the compound comprises SEQ ID NO: 224, the compound consists essentially of SEQ ID NO: 224 or the compound consists of SEQ ID NO: 224.

**[0036]** In another aspect, the compound has a sequence comprising SEQ ID NO: 247, the antisense oligonucleotide hybridizes with a region complementary to SEQ ID NO: 247, the compound comprises SEQ ID NO: 247, the compound consists essentially of SEQ ID NO: 247 or the compound consists of SEQ ID NO: 247.

**[0037]** In another aspect, the compound has a sequence comprising SEQ ID NO: 319, the antisense oligonucleotide

hybridizes with a region complementary to SEQ ID NO: 319, the compound comprises SEQ ID NO: 319, the compound consists essentially of SEQ ID NO: 319 or the compound consists of SEQ ID NO: 319.

**[0038]** In one embodiment, the compounds comprise at least one modified internucleoside linkage, and in another embodiment, the modified internucleoside linkage is a phosphorothioate linkage.

**[0039]** In another aspect, the compounds comprise at least one modified sugar moiety, and in one aspect, the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

**[0040]** In another embodiment, the compounds comprise at least one modified nucleobase, and in one aspect, the modified nucleobase is a 5-methylcytosine.

**[0041]** In yet another aspect, the compounds are chimeric oligonucleotides. Preferred chimeric compounds include those having one or more phosphorothioate linkages and further comprising 2'-methoxyethoxyl nucleotide wings and a ten nucleobase 2'-deoxynucleotide gap.

**[0042]** In another aspect, the compounds specifically hybridizes with and inhibits the expression of a nucleic acid molecule encoding an alternatively spliced form of apolipoprotein B.

**[0043]** The invention also provide compositions comprising a compound of the invention and a pharmaceutically acceptable carrier or diluent. In one aspect, the composition further comprises a colloidal dispersion system, and in another aspect, the compound in the composition is an antisense oligonucleotide. In certain embodiments, the composition comprises an antisense compound of the invention hybridized to a complementary strand. Hybridization of the antisense strand can form one or more blunt ends or one or more overhanging ends. In some embodiments, the overhanging end comprises a modified base.

**[0044]** The invention further provides methods of inhibiting the expression of apolipoprotein B in cells or tissues comprising contacting said cells or tissues with a compound of the invention so that expression of apolipoprotein B is inhibited. Methods are also provided for treating an animal having a disease or condition associated with apolipoprotein B comprising administering to said animal a therapeutically or prophylactically effective amount of a compound of the invention so that expression of apolipoprotein B is inhibited. In various aspects, the condition is associated with abnormal lipid metabolism, the condition is associated with abnormal cholesterol metabolism, the condition is atherosclerosis, the condition is an abnormal metabolic condition, the abnormal metabolic condition is hyperlipidemia, the disease is diabetes, the diabetes is Type 2 diabetes, the condition is obesity, and/or the disease is cardiovascular disease.

**[0045]** The invention also provide methods of modulating glucose levels in an animal comprising administering to said animal a compound of the invention, and in one aspect, the animal is a human. In various embodiments, the glucose levels are plasma glucose levels, the glucose levels are serum glucose levels, and/or the animal is a diabetic animal.

**[0046]** The invention also provides methods of preventing or delaying the onset of a disease or condition associated with apolipoprotein B in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of a compound of the invention. In one aspect, the animal is a human. In other aspects, the condition is an abnormal metabolic condition, the abnormal metabolic condition is hyperlipidemia, the disease is diabetes, the diabetes is Type 2 diabetes, the condition is obesity, the condition is atherosclerosis, the condition involves abnormal lipid metabolism, and/or the condition involves abnormal cholesterol metabolism.

**[0047]** The invention also provides methods of preventing or delaying the onset of an increase in glucose levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of a compound of the invention. In one aspect, the animal is a human. In other aspects, the glucose levels are serum glucose levels, and/or the glucose levels are plasma glucose levels.

**[0048]** The invention also provides methods of modulating serum cholesterol levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of a compound of the invention. In one aspect, the animal is a human.

**[0049]** The invention also provides methods of modulating lipoprotein levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of a compound of the invention. In one aspect, the animal is a human. In other aspects, the lipoprotein is VLDL, the lipoprotein is HDL, and/or the lipoprotein is LDL.

**[0050]** The invention also provides methods of modulating serum triglyceride levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of a compound of the invention. In one aspect, the animal is a human.

**[0051]** The invention also proves use of a compound of the invention for the manufacture of a medicament for the treatment of a disease or condition associated with apolipoprotein B expression, a medicament for the treatment of a condition associated with abnormal lipid metabolism, a medicament for the treatment of a condition associated with abnormal cholesterol metabolism, a medicament for the treatment of atherosclerosis, a medicament for the treatment of hyperlipidemia, a medicament for the treatment of diabetes, a medicament for the treatment of Type 2 diabetes, a medicament for the treatment of obesity, a medicament for the treatment of cardiovascular disease, a medicament for preventing or delaying the onset of increased glucose levels, a medicament for preventing or delaying the onset of increased serum glucose levels, a medicament for preventing or delaying the onset of increased plasma glucose levels, a medicament for the modulation of serum cholesterol levels, a medicament for the modulation of serum lipoprotein

levels, a medicament for the modulation of serum VLDL levels, a medicament for the modulation of serum HDL levels, and/or a medicament for the modulation of serum LDL levels, a medicament for the modulation of serum triglyceride levels.

5 [0052] In another aspect, the invention provides methods of decreasing circulating lipoprotein levels comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. In another aspect, the invention provides methods of reducing lipoprotein transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. The invention also provides methods of reducing lipoprotein absorption/adsorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

10 [0053] In another aspect, the invention contemplates methods of decreasing circulating triglyceride levels comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. Also provided are methods of reducing triglyceride transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. The invention further provides methods of reducing triglyceride absorption/adsorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

15 [0054] In another aspect, the invention provides methods of decreasing circulating cholesterol levels, including cholesteryl esters and/or unesterified cholesterol, comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. Also contemplated are methods of reducing cholesterol transport, including cholesteryl esters and/or unesterified cholesterol, comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. The invention also provides methods of reducing cholesterol absorption/adsorption, including cholesteryl esters and/or unesterified cholesterol, comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

20 [0055] In another aspect, the invention provides methods of decreasing circulating lipid levels comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. The invention also provides methods of reducing lipid transport in plasma comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. In addition, the invention provides methods of reducing lipid absorption/adsorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

25 [0056] The invention further contemplates methods of decreasing circulating dietary lipid levels comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. Also provided are methods of reducing dietary lipid transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, as well as methods of reducing dietary lipid absorption/adsorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

30 [0057] In another aspect, the invention provides methods of decreasing circulating fatty acid levels comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. The invention also provides methods of reducing fatty acid transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. Also contemplated are methods of reducing fatty acid absorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

35 [0058] The invention also provides methods of decreasing circulating acute phase reactants comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. In another aspect, the invention provides methods of reducing acute phase reactants transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, as well as methods of reducing acute phase reactants absorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

40 [0059] In another aspect, the invention provides methods of decreasing circulating chylomicrons comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, methods of reducing chylomicron transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, and methods of reducing chylomicron absorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

45 [0060] The invention further provides methods of decreasing circulating chylomicron remnant particles comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, methods of reducing chylomicron remnant transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, and methods of reducing chylomicron remnant absorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

**[0061]** The invention further contemplates methods of decreasing circulating VLDL, IDL, LDL, and/or HDL comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression. Likewise, the invention provides methods of reducing VLDL, IDL, LDL, and/or HDL transport comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression, in addition to methods of reducing VLDL, IDL, LDL, and/or HDL absorption comprising the step of administering to an individual an amount of a compound of the invention sufficient to reduce apolipoprotein B expression.

**[0062]** In still another aspect, the invention provides methods of treating a condition associated with apolipoprotein B expression comprising the step of administering to an individual an amount of a compound of the invention sufficient to inhibit apolipoprotein B expression, said condition selected from hyperlipoproteinemia, familial type 3 hyperlipoproteinemia (familial dysbetalipoproteinemia), and familial hyperalphalipoproteinemia; hyperlipidemia, mixed hyperlipidemias, multiple lipoprotein-type hyperlipidemia, and familial combined hyperlipidemia; hypertriglyceridemia, familial hypertriglyceridemia, and familial lipoprotein lipase; hypercholesterolemia, familial hypercholesterolemia, polygenic hypercholesterolemia, and familial defective apolipoprotein B; cardiovascular disorders including atherosclerosis and coronary artery disease; peripheral vascular disease; von Gierke's disease (glycogen storage disease, type I); lipodystrophies (congenital and acquired forms); Cushing's syndrome; sexual ateliotic dwarfism (isolated growth hormone deficiency); diabetes mellitus; hyperthyroidism; hypertension; anorexia nervosa; Werner's syndrome; acute intermittent porphyria; primary biliary cirrhosis; extrahepatic biliary obstruction; acute hepatitis; hepatoma; systemic lupus erythematosus; monoclonal gammopathies (including myeloma, multiple myeloma, macroglobulinemia, and lymphoma); endocrinopathies; obesity; nephrotic syndrome; metabolic syndrome; inflammation; hypothyroidism; uremia (hyperurecemia); impotence; obstructive liver disease; idiopathic hypercalcemia; dysglobulinemia; elevated insulin levels; Syndrome X; Dupuytren's contracture; and Alzheimer's disease and dementia.

**[0063]** The invention also provides methods of reducing the risk of a condition comprising the step of administering to an individual an amount of a compound of the invention sufficient to inhibit apolipoprotein B expression, said condition selected from pregnancy; intermittent claudication; gout; and mercury toxicity and amalgam illness.

**[0064]** The invention further provides methods of inhibiting cholesterol particle binding to vascular endothelium comprising the step of administering to an individual an amount of a compound of the invention sufficient to inhibit apolipoprotein B expression, and as a result, the invention also provides methods of reducing the risk of: (i) cholesterol particle oxidization; (ii) monocyte binding to vascular endothelium; (iii) monocyte differentiation into macrophage; (iv) macrophage ingestion of oxidized lipid particles and release of cytokines (including, but limited to IL-1, TNF-alpha, TGF-beta); (v) platelet formation of fibrous fibrofatty lesions and inflammation; (vi) endothelium lesions leading to clots; and (vii) clots leading to myocardial infarction or stroke, also comprising the step of administering to an individual an amount of a compound of the invention sufficient to inhibit apolipoprotein B expression.

**[0065]** The invention also provides methods of reducing hyperlipidemia associated with alcoholism, smoking, use of oral contraceptives, use of glucocorticoids, use of beta-adrenergic blocking agents, or use of isotretinoin (13-cis-retinoic acid) comprising the step of administering to an individual an amount of a compound of the invention sufficient to inhibit apolipoprotein B expression.

**[0066]** In certain aspects, the invention provides an antisense oligonucleotide compound 8 to 50 nucleobases in length comprising at least 8 contiguous nucleotides of SEQ ID NO:247 and having a length from at least 12 or at least 14 to 30 nucleobases.

**[0067]** In a further aspect, the invention provides an antisense oligonucleotide compound 20 nucleobases in length having a sequence of nucleobases as set forth in SEQ ID NO:247 and comprising 5-methylcytidine at nucleobases 2, 3, 5, 9, 12, 15, 17, 19, and 20, wherein every internucleoside linkage is a phosphothioate linkage, nucleobases 1-5 and 16-20 comprise a 2'-methoxyethoxyl modification, and nucleobases 6-15 are deoxynucleotides.

**[0068]** In another aspect, the invention provides a compound comprising a first nucleobase strand, 8 to 50 nucleobases in length and comprising a sequence of at least 8 contiguous nucleobases of the sequence set forth in SEQ ID NO:3, hybridized to a second nucleobase strand, 8 to 50 nucleobases in length and comprising a sequence sufficiently complementary to the first strand so as to permit stable hybridization, said compound inhibiting expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 30% or by at least 50% in 80% confluent HepG2 cells in culture at a concentration of 100 nM.

**[0069]** Further provided is a vesicle, such as a liposome, comprising a compound or composition of the invention

**[0070]** Preferred methods of administration of the compounds or compositions of the invention to an animal are intravenously, subcutaneously, or orally. Administrations can be repeated.

**[0071]** In another aspect, the invention provides a method of reducing lipoprotein(a) secretion by hepatocytes comprising (a) contacting hepatocytes with an amount of a composition comprising a non-catalytic compound 8 to 50 nucleobases in length that specifically hybridizes with mRNA encoding human apolipoprotein B and inhibits expression of the mRNA after 16 to 24 hours by at least 30% or at least 50% in 80% confluent HepG2 cells in culture at a concentration of 150 nM, wherein said amount is effective to inhibit expression of apolipoprotein B in the hepatocytes; and (b) measuring lipoprotein(a) secretion by the hepatocytes.

[0072] The invention further provides a method of a treating a condition associated with apolipoprotein B expression in a primate, such as a human, comprising administering to the primate a therapeutically or prophylactically effective amount of a non-catalytic compound 8 to 50 nucleobases in length that specifically hybridizes with mRNA encoding human apolipoprotein B and inhibits expression of the mRNA after 16 to 24 hours by at least 30% or by at least 50% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.

[0073] The invention provides a method of reducing apolipoprotein B expression in the liver of an animal, comprising administering to the animal between 2 mg/kg and 20 mg/kg of a non-catalytic compound 8 to 50 nucleobases in length that specifically hybridizes with mRNA encoding human apolipoprotein B by at least 30% or by at least 50% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.

[0074] Also provided is a method of making a compound of the invention comprising specifically hybridizing *in vitro* a first nucleobase strand comprising a sequence of at least 8 contiguous nucleobases of the sequence set forth in SEQ ID NO:3 to a second nucleobase strand comprising a sequence sufficiently complementary to said first strand so as to permit stable hybridization.

[0075] The invention further provides use of a compound of the invention in the manufacture of a medicament for the treatment of any and all conditions disclosed herein.

## DETAILED DESCRIPTION OF THE INVENTION

[0076] The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding apolipoprotein B, ultimately modulating the amount of apolipoprotein B produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding apolipoprotein B. As used herein, the terms "target nucleic acid" and "nucleic acid encoding apolipoprotein B" encompass DNA encoding apolipoprotein B, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of apolipoprotein B. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

[0077] It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding apolipoprotein B. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame

(ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function *in vivo*. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding apolipoprotein B, regardless of the sequence(s) of such codons.

[0079] It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to



a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

5 [0080] The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

10 [0081] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

20 [0082] Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

25 [0083] In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of *in vivo* assays or therapeutic treatment, and in the case of *in vitro* assays, under conditions in which the assays are performed.

30 [0084] Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

35 [0085] Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

40 [0086] For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with other antisense compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

45 [0087] Expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds which affect expression patterns.

50 [0088] Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma

and Vilo, FEBS Lett., 2000, 480, 17-24; Celis, et al., FEBS Lett., 2000, 480, 2-16), SAGE (serial analysis of gene expression)(Madden, et al., Drug Discov. Today, 2000, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, Methods Enzymol., 1999, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., Proc. Natl. Acad. Sci. U. S. A., 2000, 97, 1976-81), protein arrays and proteomics (Celis, et al., FEBS Lett., 2000, 480, 2-16; Jungblut, et al., Electrophoresis, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., FEBS Lett., 2000, 480, 2-16; Larsson, et al., J. Biotechnol., 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., Anal. Biochem., 2000, 286, 91-98; Larson, et al., Cytometry, 2000, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, Curr. Opin. Microbiol., 2000, 3, 316-21), comparative genomic hybridization (Carulli, et al., J. Cell Biochem. Suppl., 1998, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, Eur. J. Cancer, 1999, 35, 1895-904) and mass spectrometry methods (reviewed in (To, Comb. Chem. High Throughput Screen, 2000, 3, 235-41).

**[0089]** The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

**[0090]** In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. Thus, this term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages (RNA and DNA) as well as oligonucleotides having non-naturally-occurring portions which function similarly (oligonucleotide mimetics). Oligonucleotide mimetics are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

**[0091]** While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12, about 14, about 20 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression. In preferred embodiments, the antisense compound is non-catalytic oligonucleotide, i.e., is not dependent on a catalytic property of the oligonucleotide for its modulating activity. Antisense compounds of the invention can include double-stranded molecules wherein a first strand is stably hybridized to a second strand.

**[0092]** As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

**[0093]** Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

**[0094]** Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thiono-alkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

**[0095]** Representative United States patents that teach the preparation of the above phosphorus-containing linkages

include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are commonly owned with this application, and each of which is

herein incorporated by reference.

**[0096]** Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

**[0097]** Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

**[0098]** In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups.

**[0099]** The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., Science, 1991, 254, 1497-1500.

**[0100]** Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CHa-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylimino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- [wherein the native phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

**[0101]** Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O- (2-methoxyethyl) or 2'-MOE) (Martin et al., Helv. Chim. Acta, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, also described in examples hereinbelow.

**[0102]** A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH<sub>2</sub>-)<sub>n</sub> group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

**[0103]** Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 2'-allyl (2'-CH<sub>2</sub>-CH=CH<sub>2</sub>), 2'-O-allyl (2'-O-CH<sub>2</sub>-CH=CH<sub>2</sub>) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics

such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

**[0104]** Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl (-C≡C-CH<sub>3</sub>) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyl-adenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

**[0105]** Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

**[0106]** Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugate groups include cholesterols, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a

polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxysterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937). Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

**[0107]** Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

**[0108]** It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

**[0109]** Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

**[0110]** The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

**[0111]** The antisense compounds of the invention are synthesized *in vitro* and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the *in vivo* synthesis of antisense molecules.

**[0112]** The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

**[0113]** The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically

acceptable salts of such prodrugs, and other bioequivalents.

**[0114]** The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

**[0115]** The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

**[0116]** Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," J. of Pharma Sci., 1977 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

**[0117]** For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

**[0118]** The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of apolipoprotein B is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

**[0119]** The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding apolipoprotein B, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding apolipoprotein B can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of apolipoprotein B in a sample may also be prepared.

**[0120]** The present invention also includes pharmaceutical compositions and formulations which include the antisense

compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

**[0121]** Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1-10</sub> alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

**[0122]** Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate,. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactico-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in detail in United States applications 08/886,829 (filed July 1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed February 23, 1999), 09/082,624 (filed May 21, 1998) and 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

**[0123]** Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

**[0124]** Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

**[0125]** The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques

include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

5 [0126] The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

10 [0127] In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### 15 Emulsions

[0128] The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

35 [0129] Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

40 [0130] Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

50 [0131] Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar



solids such as carbon or glyceryl tristearate.

**[0132]** A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

**[0133]** Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

**[0134]** Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

**[0135]** The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

**[0136]** In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

**[0137]** The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

**[0138]** Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprinate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750) decaglycerol decaoleate (DA0750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain

(C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

**[0139]** Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

**[0140]** Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

**[0141]** There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

**[0142]** Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

**[0143]** In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

**[0144]** Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

**[0145]** Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

**[0146]** Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

**[0147]** Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

**[0148]** Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome,

the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

**[0149]** Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

**[0150]** One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

**[0151]** Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (*e.g.* as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

**[0152]** Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P.Pharma. Sci.*, 1994, 4, 6, 466).

**[0153]** Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G<sub>M1</sub>, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765).

**[0154]** Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G<sub>M1</sub>, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen *et al.*, disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G<sub>M1</sub> or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

**[0155]** Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (*e.g.*, PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, *e.g.*, DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

5 [0156] A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

10 [0157] Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

15 [0158] Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in Pharmaceutical Dosage Forms, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

20 [0159] If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

25 [0160] If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

30 [0161] If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

35 [0162] If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

40 [0163] The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in Pharmaceutical Dosage Forms, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

45 [0164] In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

50 [0165] Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

55 [0166] Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., J. Pharm. Phar-

macol., 1988, 40, 252).

**[0167]** Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein (1-monooleoyl-rac-glycerol)" dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprinate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p.92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; El Hariri et al., J. Pharm. Pharmacol., 1992, 44, 651-654).

**[0168]** Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's The Pharmacological Basis of Therapeutics, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Swinyard, Chapter 39 In: Remington's Pharmaceutical Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; Yamamoto et al., J. Pharm. Exp. Ther., 1992, 263, 25; Yamashita et al., J. Pharm. Sci., 1990, 79, 579-583).

**[0169]** Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, J. Chromatogr., 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines)(Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; Buur et al., J. Control Rel., 1990, 14, 43-51).

**[0170]** Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., J. Pharm. Pharmacol., 1987, 39, 621-626).

**[0171]** Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

**[0172]** Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

#### Carriers

**[0173]** Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-

stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5; 115-121; Takakura et al., *Antisense & Nuc1. Acid Drug Dev.*, 1996, 6, 177-183).

#### Excipients

5  
 10  
 15  
 20  
 25  
 30

**[0174]** In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (*e.g.*, pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (*e.g.*, lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (*e.g.*, magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (*e.g.*, starch, sodium starch glycolate, etc.); and wetting agents (*e.g.*, sodium lauryl sulphate, etc.).

**[0175]** Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

**[0176]** Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

**[0177]** Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### 30 Pulsatile Delivery

**[0178]** The compounds of the present invention may also be administered by pulsatile delivery. "Pulsatile delivery" refers to a pharmaceutical formulations that delivers a first pulse of drug combined with a penetration enhancer and a second pulse of penetration enhancer to promote absorption of drug which is not absorbed upon release with the first pulse of penetration enhancer.

**[0179]** One embodiment of the present invention is a delayed release oral formulation for enhanced intestinal drug absorption, comprising:

- 40
- (a) a first population of carrier particles comprising said drug and a penetration enhancer, wherein said drug and said penetration enhancer are released at a first location in the intestine; and
  - (b) a second population of carrier particles comprising a penetration enhancer and a delayed release coating or matrix, wherein the penetration enhancer is released at a second location in the intestine downstream from the first location, whereby absorption of the drug is enhanced when the drug reaches the second location.

45 **[0180]** Alternatively, the penetration enhancer in (a) and (b) is different.

**[0181]** This enhancement is obtained by encapsulating at least two populations of carrier particles. The first population of carrier particles comprises a biologically active substance and a penetration enhancer, and the second (and optionally additional) population of carrier particles comprises a penetration enhancer and a delayed release coating or matrix.

50 **[0182]** A "first pass effect" that applies to orally administered drugs is degradation due to the action of gastric acid and various digestive enzymes. One means of ameliorating first pass clearance effects is to increase the dose of administered drug, thereby compensating for proportion of drug lost to first pass clearance. Although this may be readily achieved with i.v. administration by, for example, simply providing more of the drug to an animal, other factors influence the bioavailability of drugs administered via non-parenteral means. For example, a drug may be enzymatically or chemically degraded in the alimentary canal or blood stream and/or may be impermeable or semipermeable to various mucosal membranes.

55 **[0183]** It is also contemplated that these pharmaceutical compositions are capable of enhancing absorption of biologically active substances when administered via the rectal, vaginal, nasal or pulmonary routes. It is also contemplated that release of the biologically active substance can be achieved in any part of the gastrointestinal tract.

**[0184]** Liquid pharmaceutical compositions of oligonucleotide can be prepared by combining the oligonucleotide with a suitable vehicle, for example sterile pyrogen free water, or saline solution. Other therapeutic compounds may optionally be included.

**[0185]** The present invention also contemplates the use of solid particulate compositions. Such compositions preferably comprise particles of oligonucleotide that are of respirable size. Such particles can be prepared by, for example, grinding dry oligonucleotide by conventional means, for example with a mortar and pestle, and then passing the resulting powder composition through a 400 mesh screen to segregate large particles and agglomerates. A solid particulate composition comprised of an active oligonucleotide can optionally contain a dispersant which serves to facilitate the formation of an aerosol, for example lactose.

**[0186]** In accordance with the present invention, oligonucleotide compositions can be aerosolized. Aerosolization of liquid particles can be produced by any suitable means, such as with a nebulizer. See, for example, U.S. Patent No. 4,501,729. Nebulizers are commercially available devices which transform solutions or suspensions into a therapeutic aerosol mist either by means of acceleration of a compressed gas, typically air or oxygen, through a narrow venturi orifice or by means of ultrasonic agitation. Suitable nebulizers include those sold by Blairex® under the name PARI LC PLUS, PARI DURA-NEB 2000, PARI-BABY Size, PARI PRONEB Compressor with LC PLUS, PARI WALKHALER Compressor/Nebulizer System, PARI LC PLUS Reusable Nebulizer, and PARI LC Jet+® Nebulizer.

**[0187]** Exemplary formulations for use in nebulizers consist of an oligonucleotide in a liquid, such as sterile, pyrogen free water, or saline solution, wherein the oligonucleotide comprises up to about 40% w/w of the formulation. Preferably, the oligonucleotide comprises less than 20% w/w. If desired, further additives such as preservatives (for example, methyl hydroxybenzoate) antioxidants, and flavoring agents can be added to the composition.

**[0188]** Solid particles comprising an oligonucleotide can also be aerosolized using any solid particulate medicament aerosol generator known in the art. Such aerosol generators produce respirable particles, as described above, and further produce reproducible metered dose per unit volume of aerosol. Suitable solid particulate aerosol generators include insufflators and metered dose inhalers. Metered dose inhalers are used in the art and are useful in the present invention.

**[0189]** Preferably, liquid or solid aerosols are produced at a rate of from about 10 to 150 liters per minute, more preferably from about 30 to 150 liters per minute, and most preferably about 60 liters per minute.

**[0190]** Enhanced bioavailability of biologically active substances is also achieved via the oral administration of the compositions and methods of the present invention. The term "bioavailability" refers to a measurement of what portion of an administered drug reaches the circulatory system when a non-parenteral mode of administration is used to introduce the drug into an animal.

**[0191]** Penetration enhancers include, but are not limited to, members of molecular classes such as surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactant molecules. (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p. 92). Carriers are inert molecules that may be included in the compositions of the present invention to interfere with processes that lead to reduction in the levels of bioavailable drug.

#### Other Components

**[0192]** The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

**[0193]** Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

**[0194]** Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more anti-sense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosourea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosourea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycytosine, 4-hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-

FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (*e.g.*, 5-FU and oligonucleotide), sequentially (*e.g.*, 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (*e.g.*, 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

**[0195]** In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

**[0196]** The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on  $EC_{50}$ s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

#### Combination Therapy

**[0197]** The invention also provides methods of combination therapy, wherein one or more compounds of the invention and one or more other therapeutic/prophylactic compounds are administered treat a condition and/or disease state as described herein. In various aspects, the compound(s) of the invention and the therapeutic/prophylactic compound(s) are co-administered as a mixture or administered individually. In one aspect, the route of administration is the same for the compound(s) of the invention and the therapeutic/prophylactic compound(s), while in other aspects, the compound(s) of the invention and the therapeutic/prophylactic compound(s) are administered by a different routes. In one embodiment, the dosages of the compound(s) of the invention and the therapeutic/prophylactic compound(s) are amounts that are therapeutically or prophylactically effective for each compound when administered individually. Alternatively, the combined administration permits use of lower dosages than would be required to achieve a therapeutic or prophylactic effect if administered individually, and such methods are useful in decreasing one or more side effects of the reduced-dose compound.

**[0198]** In one aspect, a compound of the present invention and one or more other therapeutic/prophylactic compound(s) effective at treating a condition are administered wherein both compounds act through the same or different mechanisms. Therapeutic/prophylactic compound(s) include, but are not limited to, bile salt sequestering resins (*e.g.*, cholestyramine, colestipol, and colesevelam hydrochloride), HMGCoA-reductase inhibitors (*e.g.*, lovastatin, cerivastatin, pravastatin, atorvastatin, simvastatin, and fluvastatin), nicotinic acid, fibric acid derivatives (*e.g.*, clofibrate, gemfibrozil, fenofibrate, bezafibrate, and ciprofibrate), probucol, neomycin, dextrothyroxine, plant-stanol esters, cholesterol absorption inhibitors (*e.g.*, ezetimibe), implitapide, inhibitors of bile acid transporters (apical sodium-dependent bile acid transporters), regulators of hepatic CYP7a, estrogen replacement therapeutics (*e.g.*, tamoxifen), and anti-inflammatories (*e.g.*, glucocorticoids).

**[0199]** Accordingly, the invention further provides use of a compound of the invention and one or more other therapeutic/prophylactic compound(s) as described herein in the manufacture of a medicament for the treatment and/or prevention of a disease or condition as described herein.



## Targeted Delivery

**[0200]** In another aspect, methods are provided to target a compound of the invention to a specific tissue, organ or location in the body. Exemplary targets include liver, lung, kidney, heart, and atherosclerotic plaques within a blood vessel. Methods of targeting compounds are well known in the art.

**[0201]** In one embodiment, the compound is targeted by direct or local administration. For example, when targeting a blood vessel, the compound is administered directly to the relevant portion of the vessel from inside the lumen of the vessel, *e.g.*, single balloon or double balloon catheter, or through the adventitia with material aiding slow release of the compound, *e.g.*, a pluronic gel system as described by Simons et al., Nature 359: 67-70 (1992). Other slow release techniques for local delivery of the compound to a vessel include coating a stent with the compound. Methods of delivery of antisense compounds to a blood vessel are disclosed in U.S. Patent No. 6,159,946, which is incorporated by reference in its entirety.

**[0202]** When targeting a particular tissue or organ, the compound may be administered in or around that tissue or organ. For example, U.S. Patent No. 6,547,787, incorporated herein by reference in its entirety, discloses methods and devices for targeting therapeutic agents to the heart. In one aspect, administration occurs by direct injection or by injection into a blood vessel associated with the tissue or organ. For example, when targeting the liver, the compound may be administered by injection or infusion through the portal vein.

**[0203]** In another aspect, methods of targeting a compound are provided which include associating the compound with an agent that directs uptake of the compound by one or more cell types. Exemplary agents include lipids and lipid-based structures such as liposomes generally in combination with an organ- or tissue-specific targeting moiety such as, for example, an antibody, a cell surface receptor, a ligand for a cell surface receptor, a polysaccharide, a drug, a hormone, a hapten, a special lipid and a nucleic acid as described in U. S. Patent No. 6,495,532, the disclosure of which is incorporated herein by reference in its entirety. U.S. Pat. No. 5,399,331, the disclosure of which is incorporated herein by reference in its entirety, describes the coupling of proteins to liposomes through use of a crosslinking agent having at least one maleimido group and an amine reactive function; U.S. Pat. Nos. 4,885,172, 5,059,421 and 5,171,578, the disclosures of which are incorporated herein by reference in their entirety, describe linking proteins to liposomes through use of the glycoprotein streptavidin and coating targeting liposomes with polysaccharides. Other lipid based targeting agents include, for example, micelle and crystalline products as described in U.S. Patent No. 6,217,886, the disclosure of which is incorporated herein by reference in its entirety.

**[0204]** In another aspect, targeting agents include porous polymeric microspheres which are derived from copolymeric and homopolymeric polyesters containing hydrolyzable ester linkages which are biodegradable, as described in U.S. No. Patent 4,818,542, the disclosure of which is incorporated herein by reference in its entirety. Typical polyesters include polyglycolic (PGA) and polylactic (PLA) acids, and copolymers of glycolide and L(-lactide) (PGL), which are particularly suited for the methods and compositions of the present invention in that they exhibit low human toxicity and are biodegradable. The particular polyester or other polymer, oligomer, or copolymer utilized as the microspheric polymer matrix is not critical and a variety of polymers may be utilized depending on desired porosity, consistency, shape and size distribution. Other biodegradable or bioerodable polymers or copolymers include, for example, gelatin, agar, starch, arabinogalactan, albumin, collagen, natural and synthetic materials or polymers, such as, poly( $\epsilon$ -caprolactone), poly( $\epsilon$ -caprolactone-CO-lactic acid), poly( $\epsilon$ -caprolactone-CO-glycolic acid), poly( $\gamma$ -3-hydroxy butyric acid), polyethylene oxide, polyethylene, poly(alkyl-2-cyanoacrylate), (*e.g.*, methyl, ethyl, butyl), hydrogels such as poly(hydroxyethyl methacrylate), polyamides (*e.g.*, polyacrylamide), poly(amino acids) (*i.e.*, L-leucine, L-aspartic acid,  $\beta$ -methyl-L-aspartate,  $\beta$ -benzyl-L-aspartate, glutamic acid), poly(2-hydroxyethyl DL-aspartamide), poly(ester urea), poly(L-phenylalanine/ethylene glycol/1,6-diisocyanatohexane) and poly(methyl methacrylate). The exemplary natural and synthetic polymers suitable for targeted delivery are either readily available commercially or are obtainable by condensation polymerization reactions from the suitable monomers or, comonomers or oligomers.

**[0205]** In still another embodiment, U.S. Patent No. 6,562,864, the disclosure of which is incorporated herein by reference in its entirety, describes catechins, including epi and other carbo-cationic isomers and derivatives thereof, which as monomers, dimers and higher multimers can form complexes with nucleophilic and cationic bioactive agents for use as delivery agents. Catechin multimers have a strong affinity for polar proteins, such as those residing in the vascular endothelium, and on cell/organelle membranes and are particularly useful for targeted delivery of bioactive agents to select sites *in vivo*. In treatment of vascular diseases and disorders, such as atherosclerosis and coronary artery disease, delivery agents include substituted catechin multimers, including amidated catechin multimers which are formed from reaction between catechin and nitrogen containing moieties such as ammonia.

**[0206]** Other targeting strategies of the invention include ADEPT (antibody-directed enzyme prodrug therapy), GDEPT (gene-directed EPT) and VDEPT (virus-directed EPT) as described in U.S. Patent No. 6,433,012, the disclosure of which is incorporated herein by reference in its entirety.

**[0207]** The present invention further provides medical devices and kits for targeted delivery, wherein the device is, for example, a syringe, stent, or catheter. Kits include a device for administering a compound and a container comprising

a compound of the invention. In one aspect, the compound is preloaded into the device. In other embodiments, the kit provides instructions for methods of administering the compound and dosages. U.S. patents describing medical devices and kits for delivering antisense compounds include US Patent Nos. 6,368,356; 6,344,035; 6,344,028; 6,287,285; 6,200,304; 5,824,049; 5,749,915; 5,674,242; 5,670,161; 5,609,629; 5,593,974; and 5,470,307 (all incorporated herein by reference in their entirety).

[0208] While the present invention has been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

## EXAMPLES

### Example 1

#### Nucleoside Phosphoramidites for Oligonucleotide Syntheses Deoxy and 2'-alkoxy amidites

[0209] 2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

[0210] Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., Nucleic Acids Research, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

#### 2'-Fluoro amidites

##### 2'-Fluorodeoxyadenosine amidites

[0211] 2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., J. Med. Chem., 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoro-adenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

##### 2'-Fluorodeoxyguanosine

[0212] The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups.

[0213] Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

##### 2'-Fluorouridine

[0214] Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

##### 2'-Fluorodeoxycytidiae

[0215] 2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

**2'-O-(2-Methoxyethyl) modified amidites**

[0216] 2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., Helvetica Chimica Acta, 1995, 78, 486-504.

**2,2'-An.hydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

[0217] 5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

**2'-O-Methoxyethyl-5-methyluridiae**

[0218] 2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridiae**

[0219] 2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL). The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

[0220] 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridirie**

[0221] A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole

## EP 2 336 318 A1

(90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POC<sub>13</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

### 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

[0222] A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

### N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

[0223] 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

### N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

[0224] N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

### 2'-O-(Aminoxyethyl) nucleoside amidites and 2'-O-(dimethylaminoxyethyl) nucleosides amidites

#### 2'-(Dimethylaminoxyethoxy) nucleosides amidites

[0225] 2'-(Dimethylaminoxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminoxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

### 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine

[0226] O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. *tert*-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (Rf 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10° C. The resulting crystalline product was collected by filtration,

## EP 2 336 318 A1

washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

### 5'-O-*tert*-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine

[0227] In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-*tert*-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

### 2'-O-([2-phthalimidoxy)ethyl]-5'-*t*-butyldiphenylsilyl-5-methyluridine

[0228] 5'-O-*tert*-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-*t*-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

### 5'-O-*tert*-butyldiphenylsilyl-2'-O-([2-formadoximinoxy)ethyl]-5-methyluridine

[0229] 2'-O-([2-phthalimidoxy)ethyl]-5'-*t*-butyldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH<sub>2</sub>Cl<sub>2</sub> and the combined organic phase was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-([2-formadoximinoxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%).

### 5'-O-*tert*-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine

[0230] 5'-O-*tert*-butyldiphenylsilyl-2'-O-([2-formadoximinoxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

**2'-O-(dimethylaminooxyethyl)-5-methyluridine**

[0231] Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-*tert*-butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-(dimethylaminooxyethyl)-5-methyluridine (766mg, 92.5%).

**5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine**

[0232] 2'-O-(dimethylaminooxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine (1.13g, 80%).

**5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

[0233] 5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

**2'-(Aminoxyethoxy) nucleoside amidites**

[0234] 2'-(Aminoxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminoxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

[0235] The 2'-O-aminoxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-([2-phthalimidoxy]ethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

[0236] 2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

**[0237]** 2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2-</sup>, 2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)-ethyl]-5-methyl uridine**

**[0238]** To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylamino-ethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)-ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

**[0239]** Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

**Example 2****Oligonucleotide synthesis**

**[0240]** Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

**[0241]** Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

**[0242]** Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

**[0243]** 3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

**[0244]** Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

**[0245]** Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

**[0246]** 3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

**[0247]** Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

**[0248]** Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

**Example 3****Oligonucleoside Synthesis**

5 **[0249]** Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 10 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

**[0250]** Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

**[0251]** Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

15

**Example 4****PNA Synthesis**

20 **[0252]** Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, Bioorganic & Medicinal Chemistry, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

25 **Example 5****Synthesis of Chimeric Oligonucleotides**

30 **[0253]** Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

35

**[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric Phosphorothioate oligonucleotides**

**[0254]** Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

50

**[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides**

**[0255]** [2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

55



**[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides**

**[0256]** [2'-O-(2-methoxyethyl phosphodiester]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

**[0257]** Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

**Example 6****Oligonucleotide Isolation**

**[0258]** After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., J. Biol. Chem. 1991, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

**Example 7****Oligonucleotide Synthesis - 96 Well Plate Format**

**[0259]** Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

**[0260]** Oligonucleotides were cleaved from support and deprotected with concentrated NH<sub>4</sub>OH at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

**Example 8****Oligonucleotide Analysis - 96 Well Plate Format**

**[0261]** The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

**Example 9****Cell culture and oligonucleotide treatment**

**[0262]** The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell

## EP 2 336 318 A1

types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 7 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

5 **[0263]** T-24 cells:

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

15 **[0264]** For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

**[0265]** A549 cells:

20 The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

25 **[0266]** NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

30 **[0267]** HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

35 **[0268]** HepG2 cells:

40 The human hepatoblastoma cell line HepG2 was obtained from the American Type Culture Collection (Manassas, VA). HepG2 cells were routinely cultured in Eagle's MEM supplemented with 10% fetal calf serum, non-essential amino acids, and 1 mM sodium pyruvate (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

45 **[0269]** For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

**[0270]** AML12 cells:

50 The AML12 (alpha mouse liver 12) cell line was established from hepatocytes from a mouse (CD1 strain, line MT42) transgenic for human TGF alpha. Cells are cultured in a 1:1 mixture of Dulbecco's modified Eagle's medium and Ham's F12 medium with 0.005 mg/ml insulin, 0.005 mg/ml transferrin, 5 ng/ml selenium, and 40 ng/ml dexamethasone, and 90%; 10% fetal bovine serum. For subculturing, spent

medium is removed and fresh media of 0.25% trypsin, 0.03% EDTA solution is added. Fresh trypsin solution (1 to 2 ml) is added and the culture is left to sit at room temperature until the cells detach.

55 **[0271]** Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

**[0272]** For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

## EP 2 336 318 A1

### [0273] Primary mouse hepatocytes:

Primary mouse hepatocytes were prepared from CD-1 mice purchased from Charles River Labs (Wilmington, MA) and were routinely cultured in Hepatocyte Attachment Media (Gibco) supplemented with 10% Fetal Bovine Serum (Gibco/Life Technologies, Gaithersburg, MD), 250nM dexamethasone (Sigma), and 10nM bovine insulin (Sigma). Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 10000 cells/well for use in RT-PCR analysis.

[0274] For Northern blotting or other analyses, cells are plated onto 100 mm or other standard tissue culture plates coated with rat tail collagen (200ug/mL) (Becton Dickinson) and treated similarly using appropriate volumes of medium and oligonucleotide.

### [0275] Hep3B cells:

The human hepatocellular carcinoma cell line Hep3B was obtained from the American Type Culture Collection (Manassas, VA). Hep3B cells were routinely cultured in Dulbecco's MEM high glucose supplemented with 10% fetal calf serum, L-glutamine and pyridoxine hydrochloride (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 24-well plates (Falcon-Primaria #3846) at a density of 50,000 cells/well for use in RT-PCR analysis.

[0276] For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

### [0277] Rabbit primary hepatocytes:

Primary rabbit hepatocytes were purchased from Invitro Technologies (Gaithersburg, MD) and maintained in Dulbecco's modified Eagle's medium (Gibco). When purchased, the cells had been seeded into 96-well plates for use in RT-PCR analysis and were confluent.

[0278] For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly using appropriate volumes of medium and oligonucleotide.

### [0279] HeLa cells:

The human epitheloid carcinoma cell line HeLa was obtained from the American Tissue Type Culture Collection (Manassas, VA). HeLa cells were routinely cultured in DMEM, high glucose (Invitrogen Corporation, Carlsbad, CA) supplemented with 10% fetal bovine serum (Invitrogen Corporation, Carlsbad, CA). Cells were seeded into 24-well plates (Falcon-Primaria #3846) at a density of 50,000 cells/well for use in RT-PCR analysis. Cells were

routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells 96-well plates (Falcon-Primaria #3872) at a density of 5,000 cells/well for use in RT-PCR analysis. For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

### [0280] Human mammary epithelial cells:

Normal human mammary epithelial cells (HMECs) were obtained from the American Type Culture Collection (Manassas VA). HMECs were routinely cultured in DMEM low glucose (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #353872, BD Biosciences, Bedford, MA) at a density of 7000 cells/well for use in RT-PCR analysis. For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

### [0281] Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM™-1 containing 3.75  $\mu$ g/mL LIPOFECTIN™ (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh

medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

[0282] The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCAT-TCTGCCCCAAGGA**, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments. The concentrations of antisense oligonucleotides used herein are from 5 nM to 300 nM.

### Example 10

#### Analysis of oligonucleotide inhibition of apolipoprotein B expression

[0283] Antisense modulation of apolipoprotein B expression can be assayed in a variety of ways known in the art. For example, apolipoprotein B mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)<sup>+</sup> mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions.

[0284] Protein levels of apolipoprotein B can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to apolipoprotein B can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

[0285] Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

### Example 11

#### Poly(A)<sup>+</sup> mRNA isolation

[0286] Poly(A)<sup>+</sup> mRNA was isolated according to Miura et al., Clin. Chem., 1996, 42, 1758-1764. Other methods for poly(A)<sup>+</sup> mRNA isolation are taught in, for example, Ausubel, F.M. et al., Current Protocols in Molecular Biology, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

[0287] Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all

solutions.

### Example 12

#### 5 Total RNA Isolation

[0288] Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 100  $\mu$ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60  $\mu$ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60  $\mu$ L water.

[0289] The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

### Example 13

25

#### Real-time Quantitative PCR Analysis of apolipoprotein B mRNA Levels

[0290] Quantitation of apolipoprotein B mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

[0291] Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

[0292] PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units

## EP 2 336 318 A1

AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 µL total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

5 **[0293]** Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreen™ (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen™ RNA quantification reagent from Molecular Probes. Methods of RNA quantification by RiboGreen™ are taught in Jones, L.J., et al, Analytical Biochemistry, 1998, 265, 10 368-374.

**[0294]** In this assay, 175 µL of RiboGreen™ working reagent (RiboGreen™ reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25µL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

15 **[0295]** Probes and primers to human apolipoprotein B were designed to hybridize to a human apolipoprotein B sequence, using published sequence information (GenBank accession number NM\_000384.1, incorporated herein as SEQ ID NO: 3). For human apolipoprotein B the PCR primers were:

forward primer: TGCTAAAGGCACATATGGCCT (SEQ ID NO: 4)

reverse primer: CTCAGGTTGGACTCTCCATTGAG (SEQ ID NO: 5) and the PCR probe was: FAM-CTTGTCA-  
20 GAGGGATCCTAACACTGGCCG-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For human GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 7)

reverse primer: GAAGATGGTGATGGGATTC (SEQ ID NO: 8) and the PCR probe was: 5' JOE-CAAGCTTC-  
25 CCGTTCTCAGCC-TAMRA 3' (SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

30 **[0296]** Probes and primers to mouse apolipoprotein B were designed to hybridize to a mouse apolipoprotein B sequence, using published sequence information (GenBank accession number M35186, incorporated herein as SEQ ID NO: 10). For mouse apolipoprotein B the PCR primers were:

forward primer: CGTGGGCTCCAGCATTCTA (SEQ ID NO: 11)

reverse primer: AGTCATTCTGCCTTTGCGTC (SEQ ID NO: 12) and the PCR probe was: FAM-CCAATGGTCG-  
35 GGCCTGCTCAA-TAMRA SEQ ID NO: 13) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For mouse GAPDH the PCR primers were:

forward primer: GGCAAATCAACGGCACAGT (SEQ ID NO: 14)

reverse primer: GGGTCTCGCTCCTGGAAGAT (SEQ ID NO:15) and the PCR probe was: 5' JOE-AAGGCCGA-  
40 GAATGGGAAGCTTGTCATC-TAMRA 3' (SEQ ID NO: 16) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

### Example 14

#### Northern blot analysis of apolipoprotein B mRNA levels

45 **[0297]** Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA) and then robed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions.

55 **[0298]** To detect human apolipoprotein B, a human apolipoprotein B specific probe was prepared by PCR using the forward primer TGCTAAAGGCACATATGGCCT (SEQ ID NO: 4) and the reverse primer CTCAGGTTGGACTCTCCATTGAG (SEQ ID NO: 5). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

## EP 2 336 318 A1

**[0299]** To detect mouse apolipoprotein B, a human apolipoprotein B specific probe was prepared by PCR using the forward primer CGTGGGCTCCAGCATTCTA (SEQ ID NO: 11) and the reverse primer AGTCATTTCTGCCTTTGCGTC (SEQ ID NO: 12). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for mouse glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

**[0300]** Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

### Example 15

#### Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

**[0301]** In accordance with the present invention, a series of oligonucleotides was designed to target different regions of the human apolipoprotein B RNA, using published sequence (GenBank accession number NM\_000384.1, incorporated herein as SEQ ID NO: 3). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxy-nucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments in which HepG2 cells were treated with 150 nM of the compounds in Table 1. If present, "N.D." indicates "no data".

Table 1 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
147780	5'UTR	3	1	CCGCAGGTCCCGGTGGGAA T	40	17
147781	5'UTR	3	21	ACCGAGAAGGGCACTCAGC C	35	18
147782	5'UTR	3	71	GCCTCGGCCTCGCGGCCCT G	67	19
147783	Start Codon	3	114	TCCATCGCCAGCTGCGGTG G	N.D.	20
147784	Coding	3	151	CAGCGCCAGCAGCGCCAGC A	70	21
147785	Coding	3	181	GCCCGCCAGCAGCAGCAGC A	29	22
147786	Coding	3	321	CTTGAATCAGCAGTCCCAGG	34	23
147787	Coding	3	451	CTTCAGCAAGGCTTTGCCCT	N.D.	24
147788	Coding	3	716	TTTCTGTTGCCACATTGCC	95	25
147789	Coding	3	911	GGAAGAGGTGTTGCTCCTTG	24	26
147790	Coding	3	951	TGTGCTACCATCCCATACTT	33	27
147791	Coding	3	1041	TCAAATGCGAGGCCCATCTT	N.D.	28
147792	Coding	3	1231	GGACACCTCAATCAGCTGTG	26	29
147793	Coding	3	1361	TCAGGGCCACCAGGTAGGT G	N.D.	30

EP 2 336 318 A1

(continued)

	ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
5	147794	Coding	3	1561	GTAATCTTCATCCCCAGTGC	47	31
	147795	Coding	3	1611	TGCTCCATGGTTTGGCCCAT	N.D.	32
	147796	Coding	3	1791	GCAGCCAGTCGCTTATCTCC	8	33
10	147797	Coding	3	2331	GTATAGCCAAAGTGGTCCAC	N.D.	34
	147798	Coding	3	2496	CCCAGGAGCTGGAGGTCAT G	N.D.	35
	147799	Coding	3	2573	TTGAGCCCTTCCTGATGACC	N.D.	36
15	147800	Coding	3	2811	ATCTGGACCCCACTCCTAGC	N.D.	37
	147801	Coding	3	2842	CAGACCCGACTCGTGGAAGA	38	38
	147802	Coding	3	3367	GCCCTCAGTAGATTCATCAT	N.D.	39
20	147803	Coding	3	3611	GCCATGCCACCCTCTTGGA	N.D.	40
	147804	Coding	3	3791	AACCCACGTGCCGAAAGTC	N.D.	41
	147805	Coding	3	3841	ACTCCAGATGCCTTCTGAA	N.D.	42
25	147806	Coding	3	4281	ATGTGGTAACGAGCCCGAAG	100	43
	147807	Coding	3	4391	GGCGTAGAGACCCATCACAT	25	44
	147808	Coding	3	4641	GTGTTAGGATCCCTCTGACA	N.D.	45
	147809	Coding	3	5241	CCCAGTGATAGCTCTGTGAG	60	46
30	147810	Coding	3	5355	ATTTACGCATATGAGCCCAT	0	47
	147811	Coding	3	5691	CCCTGAACCTTAGCAACAGT	N.D.	48
	147812	Coding	3	5742	GCTGAAGCCAGCCAGCGA T	N.D.	49
35	147813	Coding	3	5891	ACAGCTGCCCAGTATGTTCT	N.D.	50
	147814	Coding	3	7087	CCAATAAGATTTATAACAA	34	51
	147815	Coding	3	7731	TGGCCTACCAGAGACAGGTA	45	52
40	147816	Coding	3	7841	TCATACGTTTAGCCCAATCT	100	53
	147817	Coding	3	7901	GCATGGTCCCAAGGATGGTC	0	54
	147818	Coding	3	8491	AGTGATGGAAGCTGCGATAC	30	55
	147819	Coding	3	9181	ATGAGCATCATGCCTCCAG	N.D.	56
45	147820	Coding	3	9931	GAACACATAGCCGAATGCCG	100	57
	147821	Coding	3	10263	GTGGTGCCCTCTAATTTGTA	N.D.	58
	147822	Coding	3	10631	CCCGAGAAAGAACCGAACCC	N.D.	59
50	147823	Coding	3	10712	TGCCCTGCAGCTTCACTGAA	19	60
	147824	Coding	3	11170	GAAATCCATAAGCTCTTGT	N.D.	61
	147825	Coding	3	12301	AGAAGCTGCCTCTTCTTCCC	72	62
	147826	Coding	3	12401	TCAGGGTGAGCCCTGTGTGT	80	63
55	147827	Coding	3	12471	CTAATGGCCCTTGATAAAC	13	64
	147828	Coding	3	12621	ACGTTATCCTTGAGTCCCTG	12	65



(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
147829	Coding	3	12741	TATATCCCAGGTTTCCCCGG	64	66
147830	Coding	3	12801	ACCTGGGACAGTACCGTCCC	N.D.	67
147831	3'UTR	3	13921	CTGCCTACTGCAAGGCTGGC	0	68
147832	3'UTR	3	13991	AGAGACCTTCCGAGCCCTGG	N.D.	69
147833	3'UTR	3	14101	ATGATACACAATAAAGACTC	25	70

**[0302]** As shown in Table 1, SEQ ID NOs 17, 18, 19, 21, 23, 25, 27, 31, 38, 43, 46, 51, 52, 53, 55, 57, 62, 63 and 66 demonstrated at least 30% inhibition of human apolipoprotein B expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention. As apolipoprotein B exists in two forms in mammals (ApoB-48 and ApoB-100) which are colinear at the amino terminus, antisense oligonucleotides targeting nucleotides 1-6530 hybridize to both forms, while those targeting nucleotides 6531-14121 are specific to the long form of apolipoprotein B.

#### Example 16

#### Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap-Dose Response Study

**[0303]** In accordance with the present invention, a subset of the antisense oligonucleotides in Example 15 were further investigated in dose-response studies. Treatment doses were 50, 150 and 250 nM. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments and are shown in Table 2.

Table 2 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	Percent Inhibition		
	50 nM	150 nM	250 nM
147788	54	63	72
147806	23	45	28
147816	25	81	65
147820	10	0	73

#### Example 17

#### Antisense inhibition of mouse apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

**[0304]** In accordance with the present invention, a series of oligonucleotides was designed to target different regions of the mouse apolipoprotein B RNA, using published sequence (GenBank accession number M35186, incorporated herein as SEQ ID NO: 10). The oligonucleotides are shown in Table 3. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 3 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxy-nucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on mouse apolipoprotein B mRNA levels in primary mouse hepatocytes by quantitative real-time PCR as described in other

**EP 2 336 318 A1**

examples herein. Primary mouse hepatocytes were treated with 150 nM of the compounds in Table 3. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 3 Inhibition of mouse apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
147475	Coding	10	13	ATTGTATGTGAGAGGTGAGG	79	71
147476	Coding	10	66	GAGGAGATTGGATCTTAAGG	13	72
147477	Coding	10	171	CTTCAAATTGGGACTCTCCT	N.D.	73
147478	Coding	10	211	TCCAGGAATTGAGCTTGTGC	78	74
147479	Coding	10	238	TTCAGGACTGGAGGATGAG G	N.D.	75
147480	Coding	10	291	TCTCACCTCATGCTCCATT	54	76
147481	Coding	10	421	TGACTGTCAAGGGTGAGCT G	24	77
147482	Coding	10	461	GTCCAGCCTAGGAACACTCA	59	78
147483	Coding	10	531	ATGTCAATGCCACATGTCCA	N.D.	79
147484	Coding	10	581	TTCATCCGAGAAGTTGGGAC	49	80
147485	Coding	10	601	ATTTGGGACGAATGTATGCC	64	81
147486	Coding	10	711	AGTTGAGGAAGCCAGATTCA	N.D.	82
147487	Coding	10	964	TTCCCAGTCAGCTTTAGTGG	73	83
147488	Coding	10	1023	AGCTTGCTTGTGGGCACG G	72	84
147489	Coding	10	1111	CCTATACTGGCTTCTATGTT	5	85
147490	Coding	10	1191	TGAACTCCGTGTAAGGCAAG	N.D.	86
147491	Coding	10	1216	GAGAAATCCTTCAGTAAGGG	71	87
147492	Coding	10	1323	CAATGGAATGCTTGCTACTG	68	88
147493	Coding	10	1441	GCTTCATTATAGGAGGTGGT	41	89
147494	Coding	10	1531	ACAACTGGGATAGTGTAGCC	84	90
147495	Coding	10	1631	GTTAGGACCAGGGATTGTGA	0	91
147496	Coding	10	1691	ACCATGGAAAACCTGGCAACT	19	92
147497	Coding	10	1721	TGGGAGGAAAAACTTGAATA	N.D.	93
147498	Coding	10	1861	TGGGCAACGATATCTGATTG	0	94
147499	Coding	10	1901	CTGCAGGGCGTCAGTGACA A	29	95
147500	Coding	10	1932	GCATCAGACGTGATGTTCCC	N.D.	96
147501	Coding	10	2021	CTTGGTTAAACTAATGGTGC	18	97
147502	Coding	10	2071	ATGGGAGCATGGAGGTTGG C	16	98
147503	Coding	10	2141	AATGGATGATGAAACAGTGG	26	99
147504	Coding	10	2201	ATCAATGCCTCCTGTTGCAG	N.D.	100
147505	Coding	10	2231	GGAAGTGAGACTTTCTAAGC	76	101

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
147506	Coding	10	2281	AGGAAGGAACTCTTGATATT	58	102
147507	Coding	10	2321	ATTGGCTTCATTGGCAACAC	81	103
147759	Coding	10	1	AGGTGAGGAAGTTGGAATTC	19	104
147760	Coding	10	121	TTGTTCCCTGAAGTTGTTAC	N.D	105
147761	Coding	10	251	G TTCATGGATTCCCTCAGGA	45	106
147762	Coding	10	281	ATGCTCCATTCTCACATGCT	46	107
147763	Coding	10	338	TGCGACTGTGTCTGATTTCC	34	108
147764	Coding	10	541	G TCCCTGAAGATGTCAATGC	97	109
147765	Coding	10	561	AGGCCCAGTTCATGACCCT	59	110
147766	Coding	10	761	GGAGCCCACGTGCTGAGAT T	59	111
147767	Coding	10	801	CGTCCTTGAGCAGTGCCCG A	5	112
147768	Coding	10	1224	CCCATATGGAGAAATCCTTC	24	113
147769	Coding	10	1581	CATGCCTGGAAGCCAGTGT C	89	114
147770	Coding	10	1741	GTGTTGAATCCCTTGAAATC	67	115
147771	Coding	10	1781	GGTAAAGTTGCCCATGGCTG	68	116
147772	Coding	10	1841	GTTATAAAGTCCAGCATTGG	78	117
147773	Coding	10	1931	CATCAGACGTGATGTTCCCT	85	118
147774	Coding	10	1956	TGGCTAGTTTCAATCCCCTT	84	119
147775	Coding	10	2002	CTGTCATGACTGCCCTTTAC	52	120
147776	Coding	10	2091	GCTTGAAGTTCATTGAGAAT	92	121
147777	Coding	10	2291	TTCCTGAGAAAGGAAGGAAC	N.D	122
147778	Coding	10	2331	TCAGATATACATTGGCTTCA	14	123

[0305] As shown in Table 3, SEQ ID Nos 71, 74, 76, 78, 81, 83, 84, 87, 88, 90, 101, 102, 103, 109, 111, 111, 114, 115, 116, 117, 118, 119, 120 and 121 demonstrated at least 50% inhibition of mouse apolipoprotein B expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

**Example 18**

**Antisense inhibition mouse apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap- Dose Response Study**

[0306] In accordance with the present invention, a subset of the antisense oligonucleotides in Example 17 were further investigated in dose-response studies. Treatment doses were 50, 150 and 300 nM. The compounds were analyzed for their effect on mouse apolipoprotein B mRNA levels in primary hepatocytes cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments and are shown in Table 4.

EP 2 336 318 A1

Table 4 Inhibition of mouse apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	Percent Inhibition		
	50 nM	150 nM	300 nM
147483	56	88	89
147764	48	84	90
147769	3	14	28
147776	0	17	44

Example 19

Western blot analysis of apolipoprotein B protein levels

[0307] Western blot analysis (immunoblot analysis) was carried out using standard methods. Cells were harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels were run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to apolipoprotein B was used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands were visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA) or the ECL+ chemiluminescent detection system (Amersham Biosciences, Piscataway, NJ).

Example 20

Effects of antisense inhibition of apolipoprotein B (ISIS 147764) in C57BL/6 mice: Lean animals vs. High Fat Fed animals.

[0308] C57BL/6 mice, a strain reported to be susceptible to hyperlipidemia-induced atherosclerotic plaque formation were used in the following studies to evaluate antisense oligonucleotides as potential lipid lowering compounds in lean versus high fat fed mice.

[0309] Male C57BL/6 mice were divided into two matched groups; (1) wild-type control animals (lean animals) and (2) animals receiving a high fat diet (60% kcal fat). Control animals received saline treatment and were maintained on a normal rodent diet. After overnight fasting, mice from each group were dosed intraperitoneally every three days with saline or 50 mg/kg ISIS 147764 (SEQ ID No: 109) for six weeks. At study termination and forty eight hours after the final injections, animals were sacrificed and evaluated for target mRNA levels in liver, cholesterol and triglyceride levels, liver enzyme levels and serum glucose levels.

[0310] The results of the comparative studies are shown in Table 5.

Table 5 Effects of ISIS 147764 treatment on apolipoprotein B mRNA, cholesterol, lipid, triglyceride, liver enzyme and glucose levels in lean and high fat mice.

Treatment Group	Percent Change								
	mRNA	CHOL	Lipoproteins			TRIG	Liver Enzymes		GLUC
			VLDL	LDL	HDL		AST	ALT	
Lean-control	-73	-63	No change	-64	-44	-34	Slight decrease	No change	No change
High Fat Group	-87	-67	No change	-87	-65	No change	Slight decrease	Slight increase	-28

[0311] It is evident from these data that treatment with ISIS 147764 lowered cholesterol as well as LDL and HDL lipoproteins and serum glucose in both lean and high fat mice and that the effects demonstrated are, in fact, due to the inhibition of apolipoprotein B expression as supported by the decrease in mRNA levels. No significant changes in liver

enzyme levels were observed, indicating that the antisense oligonucleotide was not toxic to either treatment group.

### Example 21

#### Effects of antisense inhibition of apolipoprotein B (ISIS 147764) on High Fat Fed Mice; 6 Week Timecourse Study

[0312] In accordance with the present invention, a 6-week timecourse study was performed to further investigate the effects of ISIS 147764 on lipid and glucose metabolism in high fat fed mice.

[0313] Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of treatment with the antisense oligonucleotide, ISIS 147764. Control animals received saline treatment (50 mg/kg). A subset of animals received a daily oral dose (20 mg/kg) atorvastatin calcium (Lipitor®, Pfizer Inc.). All mice, except atorvastatin-treated animals, were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks. Serum cholesterol and lipoproteins were analyzed at 0, 2 and 6 week interim timepoints. At study termination, animals were sacrificed 48 hours after the final injections and evaluated for levels of target mRNA levels in liver, cholesterol, lipoprotein, triglyceride, liver enzyme (AST and ALT) and serum glucose levels as well as body, liver, spleen and fat pad weights.

### Example 22

#### Effects of antisense inhibition of apolipoprotein B (ISIS 147764) in high fat fed mice- mRNA expression in liver

[0314] Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of ISIS 147764 on mRNA expression. Control animals received saline treatment (50 mg/kg). Mice were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks. At study termination, animals were sacrificed 48 hours after the final injections and evaluated for levels of target mRNA levels in liver. ISIS 147764 showed a dose-response effect, reducing mRNA levels by 15, 75 and 88% at doses of 5, 25 and 50 mg/kg, respectively.

[0315] Liver protein samples collected at the end of the treatment period were subjected to immunoblot analysis using an antibody directed to mouse apolipoprotein B protein (Gladstone Institute, San Francisco, CA). These data demonstrate that treatment with ISIS 147764 decreases apolipoprotein B protein expression in liver in a dose-dependent manner, in addition to reducing mRNA levels.

### Example 23

#### Effects of antisense inhibition of apolipoprotein B (ISIS 147764) on serum cholesterol and triglyceride levels

[0316] Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of ISIS 147764 on serum cholesterol and triglyceride levels. Control animals received saline treatment (50 mg/kg). Mice were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks.

[0317] Serum cholesterol levels were measured at 0, 2 and 6 weeks and this data is shown in Table 6. Values in the table are expressed as percent inhibition and are normalized to the saline control.

[0318] In addition to serum cholesterol, at study termination, animals were sacrificed 48 hours after the final injections and evaluated for triglyceride levels.

[0319] Mice treated with ISIS 147764 showed a reduction in both serum cholesterol (240 mg/dL for control animals and 225, 125 and 110 mg/dL for doses of 5, 25, and 50 mg/kg, respectively) and triglycerides (115 mg/dL for control animals and 125, 150 and 85 mg/dL for doses of 5, 25, and 50 mg/kg, respectively) to normal levels by study end. These data were also compared to the effects of atorvastatin calcium at an oral dose of 20 mg/kg which showed only a minimal decrease in serum cholesterol of 20 percent at study termination.

Table 6 Percent Inhibition of mouse apolipoprotein B cholesterol levels by ISIS 147764

time	Percent Inhibition			
	Saline	5 mg/kg	25 mg/kg	50 mg/kg
0 weeks	0	0	0	0
2 weeks	0	5	12	20

EP 2 336 318 A1

(continued)

time	Percent Inhibition			
	Saline	5 mg/kg	25 mg/kg	50 mg/kg
6 weeks	0	10	45	55

**Example 24**

**Effects of antisense inhibition of apolipoprotein B (ISIS 147764) on lipoprotein levels**

**[0320]** Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of ISIS 147764 on lipoprotein (VLDL, LDL, and HDL) levels. Control animals received saline treatment (50 mg/kg). Mice were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks.

**[0321]** Lipoprotein levels were measured at 0, 2 and 6 weeks and this data is shown in Table 7. Values in the table are expressed as percent inhibition and are normalized to the saline control. Negative values indicate an observed increase in lipoprotein levels.

**[0322]** These data were also compared to the effects of atorvastatin calcium at a daily oral dose of 20 mg/kg at 0, 2 and 6 weeks.

**[0323]** These data demonstrate that at a dose of 50 mg/kg, ISIS 147764 is capable of lowering all categories of serum lipoproteins investigated to a greater extent than atorvastatin.

Table 7 Percent Inhibition of mouse apolipoprotein B lipoprotein levels by ISIS 147764 as compared to atorvastatin

Lipoprotein	Time (weeks)	Percent Inhibition					atorvastatin (20 mg/kg)
		Saline	5 mg/kg	25 mg/kg	50 mg/kg		
VLDL	0	0	0	0	0	0	
	2	0	25	30	40	15	
	6	0	10	-30	15	-5	
LDL	0	0	0	0	0	0	
	2	0	-30	10	40	10	
	6	0	-10	55	90	-10	
HDL	0	0	0	0	0	0	
	2	0	5	10	10	15	
	6	0	10	45	50	20	

**Example 25**

**Effects of antisense inhibition of apolipoprotein B (ISIS 147764) on serum AST and ALT levels**

**[0324]** Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of ISIS 147764 on liver enzyme (AST and ALT) levels. Increased levels of the liver enzymes ALT and AST indicate toxicity and liver damage. Control animals received saline treatment (50 mg/kg). Mice were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks. AST and ALT levels were measured at 6 weeks.

**[0325]** Mice treated with ISIS 147764 showed no significant change in AST levels over the duration of the study compared to saline controls (105, 70 and 80 IU/L for doses of 5, 25 and 50 mg/kg, respectively compared to 65 IU/L for saline control). Mice treated with atorvastatin at a daily oral dose of 20 mg/kg had AST levels of 85 IU/L.

**[0326]** ALT levels were increased by all treatments with ISIS 147764 over the duration of the study compared to saline controls (50, 70 and 100 IU/L for doses of 5, 25 and 50 mg/kg, respectively compared to 25 IU/L for saline control). Mice

treated with atorvastatin at a daily oral dose of 20 mg/kg had AST levels of 40 IU/L.

**Example 26**

5 **Effects of antisense inhibition of apolipoprotein B (ISIS 147764) on serum glucose levels**

[0327] Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of ISIS 147764 on serum glucose levels. Control animals received saline treatment (50 mg/kg). Mice were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks.

10 [0328] At study termination, animals were sacrificed 48 hours after the final injections and evaluated for serum glucose levels. ISIS 147764 showed a dose-response effect, reducing serum glucose levels to 225, 190 and 180 mg/dL at doses of 5, 25 and 50 mg/kg, respectively compared to the saline control of 300 mg/dL. Mice treated with atorvastatin at a daily oral dose of 20 mg/kg had serum glucose levels of 215 mg/dL. These data demonstrate that ISIS 147764 is capable of  
15 reducing serum glucose levels in high fat fed mice.

**Example 27**

20 **Effects of antisense inhibition of apolipoprotein B (ISIS 147764) on body, spleen, liver and fat pad weight**

[0329] Male C57BL/6 mice (n=8) receiving a high fat diet (60% kcal fat) were evaluated over the course of 6 weeks for the effects of ISIS 147764 on body, spleen, liver and fat pad weight. Control animals received saline treatment (50 mg/kg). Mice were dosed intraperitoneally every three days (twice a week), after fasting overnight, with 5, 25, 50 mg/kg ISIS 147764 (SEQ ID No: 109) or saline (50 mg/kg) for six weeks.

25 [0330] At study termination, animals were sacrificed 48 hours after the final injections and body, spleen, liver and fat pad weights were measured. These data are shown in Table 8. values are expressed as percent change in body weight or organ weight compared to the saline-treated control animals. Data from mice treated with atorvastatin at a daily dose of 20 mg/kg are also shown in the table. Negative values indicated a decrease in weight.

30 Table 8 Effects of antisense inhibition of mouse apolipoprotein B on body and organ weight

Tissue	Percent Change			Atorvastatin 20 mg/kg
	Dose			
	5 mg/kg	25 mg/kg	50 mg/kg	
Total Body Wt.	5	5	-4	1
Spleen	10	10	46	10
Liver	18	70	80	15
Fat	10	6	-47	7

[0331] These data show a decrease in fat over the dosage range of ISIS 147764 counterbalanced by an increase in both spleen and liver weight with increased dose to give an overall decrease in total body weight.

45 **Example 28**

**Effects of antisense inhibition of apolipoprotein B (ISIS 147764) in B6.129P-ApoE<sup>tm1Unc</sup> knockout mice: Lean animals vs. High Fat Fed animals.**

50 [0332] B6.129P-ApoE<sup>tm1Unc</sup> knockout mice (herein referred to as ApoE knockout mice) obtained from The Jackson Laboratory (Bar Harbor, ME), are homozygous for the *ApoE<sup>tm1Unc</sup>* mutation and show a marked increase in total plasma cholesterol levels that are unaffected by age or sex. These animals present with fatty streaks in the proximal aorta at 3 months of age. These lesions increase with age and progress to lesions with less lipid but more elongated cells, typical  
55 of a more advanced stage of pre-atherosclerotic lesion.

[0333] The mutation in these mice resides in the apolipoprotein E (ApoE) gene. The primary role of the ApoE protein is to transport cholesterol and triglycerides throughout the body. It stabilizes lipoprotein structure, binds to the low density lipoprotein receptor (LDLR) and related proteins, and is present in a subclass of HDLs, providing them the ability to bind

EP 2 336 318 A1

to LDLR. ApoE is expressed most abundantly in the liver and brain. Female B6.129P-Apoetm1Unc knockout mice (ApoE knockout mice) were used in the following studies to evaluate antisense oligonucleotides as potential lipid lowering compounds.

**[0334]** Female ApoE knockout mice ranged in age from 5 to 7 weeks and were placed on a normal diet for 2 weeks before study initiation. ApoE knockout mice were then fed ad libitum a 60% fat diet, with 0.15% added cholesterol to induce dyslipidemia and obesity. Control animals were maintained on a high-fat diet with no added cholesterol. After overnight fasting, mice from each group were dosed intraperitoneally every three days with saline, 50 mg/kg of a control antisense oligonucleotide (ISIS 29837; TCGATCTCCTTTTATGCCCG; SEQ ID NO. 124) or 5, 25 or 50 mg/kg ISIS 147764 (SEQ ID No: 109) for six weeks.

**[0335]** The control oligonucleotide is a chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines.

**[0336]** At study termination and forty eight hours after the final injections, animals were sacrificed and evaluated for target mRNA levels in liver by RT-PCR methods verified by Northern Blot analysis, glucose levels, cholesterol and lipid levels by HPLC separation methods and triglyceride and liver enzyme levels (performed by LabCorp Preclinical Services; San Diego, CA). Data from ApoE knockout mice treated with atorvastatin at a daily dose of 20 mg/kg are also shown in the table for comparison.

**[0337]** The results of the comparative studies are shown in Table 9. Data are normalized to saline controls.

Table 9

Effects of ISIS 147764 treatment on apolipoprotein B mRNA, cholesterol, glucose, lipid, triglyceride and liver enzyme levels in ApoE knockout mice.						
		Percent Inhibition				
		Dose				
		Control	5 mg/kg	25 mg/kg	50 mg/kg	atorvastatin (20 mg/kg)
mRNA		0	2	42	70	10
Glucose	Glucose Levels (mg/dL)					
		225	195	209	191	162
Cholesterol	Cholesterol Levels (mg/dL)					
		1750	1630	1750	1490	938
Lipoprotein	Lipoprotein Levels (mg/dL)					
	HDL	51	49	62	61	42
	LDL	525	475	500	325	250
	VLDL	1190	1111	1194	1113	653
Liver Enzymes	Liver Enzyme Levels (IU/L)					
	AST	55	50	60	85	75
	ALT	56	48	59	87	76

**[0338]** It is evident from these data that treatment with ISIS 147764 lowered glucose and cholesterol as well as all lipoproteins investigated (HDL, LDL and VLDL) in ApoE knockout mice. Further, these decreases correlated with a decrease in both protein and RNA levels of apolipoprotein B, demonstrating an antisense mechanism of action. No significant changes in liver enzyme levels were observed, indicating that the antisense oligonucleotide was not toxic to either treatment group.



## Example 29

**Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap: Additional Oligonucleotides**

**[0339]** In accordance with the present invention, another series of oligonucleotides was designed to target different regions of the human apolipoprotein B RNA, using published sequence (GenBank accession number NM\_000384.1, incorporated herein as SEQ ID NO: 3). The oligonucleotides are shown in Table 10. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 10 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments in which HepG2 cells were treated with 150 nM of the compounds in Table 10. If present, "N.D." indicates "no data".

Table 10 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
270985	5'UTR	3	199	TTCTCTTCGGCCCTGGCGC	75	124
270986	coding	3	299	CTCCACTGGAACCTCTCAGCC	0	125
270987	exon: exon junction	3	359	CCTCCAGCTCAACCTTGCAG	0	126
270988	coding	3	429	GGGTTGAAGCCATACACCTC	6	127
270989	exon: exon junction	3	509	CCAGCTTGAGCTCATACCTG	64	128
270990	coding	3	584	CCCTCTTGATGTTCCAGGATG	42	129
270991	coding	3	669	GAGCAGTTTCCATACACGGT	21	130
270992	coding	3	699	CCCTTCCTCGTCTTGACGGT	8	131
270993	coding	3	756	TTGAAGCGATCACACTGCCC	69	132
270994	coding	3	799	GCCTTTGATGAGAGCAAGTG	51	133
270995	coding	3	869	TCCTCTTAGCGTCCAGTGTG	40	134
270996	coding	3	1179	CCTCTCAGCTCAGTAACCAG	0	135
270997	coding	3	1279	GCACTGAGGCTGTCCACACT	24	136
270998	coding	3	1419	CGCTGATCCCTCGCCATGTT	1	137
270999	coding	3	1459	GTTGACCGCGTGGCTCAGCG	76	138
271000	coding	3	1499	GCAGCTCCTGGGTCCCTGTA	22	139
271001	coding	3	1859	CCCATGGTAGAATTTGGACA	53	140
271002	exon: exon junction	3	2179	AATCTCGATGAGGTCAGCTG	48	141
271003	coding	3	2299	GACACCATCAGGAACCTGAC	46	142
271004	coding	3	2459	GCTCCTCTCCCAAGATGCGG	10	143
271005	coding	3	2518	GGCACCCATCAGAAGCAGCT	32	144
271006	coding	3	2789	AGTCCGGAATGATGATGCCC	42	145

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
271007	coding	3	2919	CTGAGCAGCTTGACTGGTCT	26	146
271008	coding	3	3100	CCCGGTCAGCGGATAGTAGG	37	147
271010	exon: exon junction	3	3449	TGTCACAACCTTAGGTGGCCC	57	148
271011	coding	3	3919	GTCTGGCAATCCCATGTTCT	51	149
271012	coding	3	4089	CCCACAGACTTGAAGTGGAG	55	150
271013	coding	3	4579	GAACTGCCCATCAATCTTGA	19	151
271014	coding	3	5146	CCCAGAGAGGCCAAGCTCTG	54	152
271015	coding	3	5189	TGTGTTCCCTGAAGCGGCCA	43	153
271016	coding	3	5269	ACCCAGAATCATGGCCTGAT	19	154
271017	coding	3	6049	GGTGCCTGTCTGCTCAGCTG	30	155
271018	coding	3	6520	ATGTGAACTTGTCTCTCCC	44	156
271019	coding	3	6639	TATGTCTGCAGTTGAGATAG	15	157
271020	coding	3	6859	TTGAATCCAGGATGCAGTAC	35	158
271021	coding	3	7459	GAGTCTCTGAGTCACCTCAC	38	159
271022	coding	3	7819	GATAGAATATTGCTCTGCAA	100	160
271023	coding	3	7861	CCCTTGCTCTACCAATGCTT	44	161
271025	coding	3	8449	TCCATTCCCTATGTCAGCAT	16	162
271026	coding	3	8589	GACTCCTTCAGAGCCAGCGG	39	163
271027	coding	3	8629	CCCATGCTCCGTTCTCAGGT	26	164
271028	coding	3	8829	CGCAGGTCAGCCTGACTAGA	98	165
271030	coding	3	9119	CAGTTAGAACACTGTGGCCC	52	166
271031	coding	3	10159	CAGTGTGATGACACTTGATT	49	167
271032	coding	3	10301	CTGTGGCTAACTTCAATCCC	22	168
271033	coding	3	10349	CAGTACTGTTATGACTACCC	34	169
271034	coding	3	10699	CACTGAAGACCGTGTGCTCT	35	170
271035	coding	3	10811	TCGTA CTGTGCTCCCAGAGG	23	171
271036	coding	3	10839	AAGAGGCCCTCTAGCTGTAA	95	172
271037	coding	3	11039	AAGACCCAGAATGAATCCGG	23	173
271038	coding	3	11779	GTCTACCTCAAAGCGTGCAG	29	174
271039	coding	3	11939	TAGAGGCTAACGTACCATCT	4	175
271041	coding	3	12149	CCATATCCATGCCACGGTG	37	176
271042	coding	3	12265	AGTTTCCTCATCAGATTCCC	57	177
271043	coding	3	12380	CCCAGTGGTACTTGTTGACA	68	178
271044	coding	3	12526	CCCAGTGGTGCCACTGGCTG	22	179
271045	coding	3	12579	GTCAACAGTTCCTGGTACAG	19	180
271046	coding	3	12749	CCCTAGTGTATATCCCAGGT	61	181

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
271048	coding	3	13009	CTGAAGATTACGTAGCACCT	7	182
271049	coding	3	13299	GTCCAGCCAACTATACTTGG	54	183
271050	coding	3	13779	CCTGGAGCAAGCTTCATGTA	42	184
281586	exon: exon junction	3	229	TGGACAGACCAGGCTGACAT	80	185
281587	coding	3	269	ATGTGTACTTCCGGAGGTGC	77	186
281588	coding	3	389	TCTTCAGGATGAAGCTGCAG	80	187
281589	coding	3	449	TCAGCAAGGCTTTGCCCTCA	90	188
281590	coding	3	529	CTGCTTCCCTTCTGGAATGG	84	189
281591	coding	3	709	TGCCACATTGCCCTTCTCTCG	90	190
281592	coding	3	829	GCTGATCAGAGTTGACAAGG	56	191
281593	coding	3	849	TACTGACAGGACTGGCTGCT	93	192
281594	coding	3	889	GATGGCTTCTGCCACATGCT	74	193
281595	coding	3	1059	GATGTGGATTTGGTGCTCTC	76	194
281596	coding	3	1199	TGACTGCTTCATCACTGAGG	77	195
281597	coding	3	1349	GGTAGGTGACCACATCTATC	36	196
281598	coding	3	1390	TCGCAGCTGCTGTGCTGAGG	70	197
281599	exon: exon junction	3	1589	TTCCAATGACCCGCAGAATC	74	198
281600	coding	3	1678	GATCATCAGTGATGGCTTTG	52	199
281601	coding	3	1699	AGCCTGGATGGCAGCTTTCT	83	200
281602	coding	3	1749	GTCTGAAGAAGAACCTCCTG	84	201
281603	coding	3	1829	TATCTGCCTGTGAAGGACTC	82	202
281604	coding	3	1919	CTGAGTTCAAGATATTGGCA	78	203
281605	exon: exon junction	3	2189	CTTCCAAGCCAATCTCGATG	82	204
281606	coding	3	2649	TGCAACTGTAATCCAGCTCC	86	205
281607	exon: exon junction	3	2729	CCAGTTCAGCCTGCATGTTG	84	206
281608	coding	3	2949	GTAGAGACCAAATGTAATGT	62	207
281609	coding	3	3059	CGTTGGAGTAAGCGCCTGAG	70	208
281610	exon: exon junction	3	3118	CAGCTCTAATCTGGTGTCCC	69	209
281611	coding	3	3189	CTGTCCTCTCTCTGGAGCTC	93	210
281612	coding	3	3289	CAAGGTCATACTCTGCCGAT	83	211
281613	coding	3	3488	GTATGGAAATAACACCCTTG	70	212
281614	coding	3	3579	TAAGCTGTAGCAGATGAGTC	63	213
281615	coding	3	4039	TAGATCTCTGGAGGATTTGC	81	214

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO	
5	281616	coding	3	4180	GTCTAGAACACCCAGGAGAG	66	215
	281617	coding	3	4299	ACCACAGAGTCAGCCTTCAT	89	216
	281618	coding	3	4511	AAGCAGACATCTGTGGTCCC	90	217
10	281619	coding	3	4660	CTCTCCATTGAGCCGGCCAG	96	218
	281620	coding	3	4919	CCTGATATTCAGAACGCAGC	89	219
	281621	coding	3	5009	CAGTGCCTAAGATGTCAGCA	53	220
15	281622	coding	3	5109	AGCACCAGGAGACTACACTT	88	221
	281623	coding	3	5212	CCCATCCAGACTGAATTTTG	59	222
	281624	coding	3	5562	GGTTCTAGCCGTAGTTTCCC	75	223
	281625	coding	3	5589	AGGTTACCAGCCACATGCAG	94	224
20	281626	coding	3	5839	ATGTGCATCGATGGTCATGG	88	225
	281627	coding	3	5869	CCAGAGAGCGAGTTTCCCAT	82	226
	281628	coding	3	5979	CTAGACACGAGATGATGACT	81	227
25	281629	coding	3	6099	TCCAAGTCTGGCTGTATTC	83	228
	281630	coding	3	6144	CGTCCAGTAAGCTCCACGCC	82	229
	281631	coding	3	6249	TCAACGGCATCTCTCATCTC	88	230
	281632	coding	3	6759	TGATAGTGCTCATCAAGACT	75	231
30	281633	coding	3	6889	GATTCTGATTTGGTACTTAG	73	232
	281634	coding	3	7149	CTCTCGATTAACTCATGGAC	81	233
	281635	coding	3	7549	ATACACTGCAACTGTGGCCT	89	234
35	281636	coding	3	7779	GCAAGAGTCCACCAATCAGA	68	235
	281637	coding	3	7929	AGAGCCTGAAGACTGACTTC	74	236
	281638	coding	3	8929	TCCCTCATCTGAGAATCTGG	66	237
	281640	coding	3	10240	CAGTGCATCAATGACAGATG	87	238
40	281641	coding	3	10619	CCGAACCCTTGACATCTCCT	72	239
	281642	coding	3	10659	GCCTCACTAGCAATAGTTCC	59	240
	281643	coding	3	10899	GACATTTGCCATGGAGAGAG	61	241
45	281644	coding	3	11209	CTGTCTCCTACCAATGCTGG	26	242
	281645	exon: exon junction	3	11979	TCTGCACTGAAGTCACGGTG	78	243
	281646	coding	3	12249	TCCCGGACCCTCAACTCAGT	76	244
50	281648	3'UTR	3	13958	GCAGGTCCAGTTCATATGTG	81	245
	281649	3'UTR	3	14008	GCCATCCTTCTGAGTTCAGA	76	246
	301012	exon: exon junction	3	3249	GCCTCAGTCTGCTTCGCACC	87	247
55	301013	5'UTR	3	3	CCCCGCAGGTCCCGGTGGG A	82	248

EP 2 336 318 A1

(continued)

	ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
5	301014	5'UTR	3	6	CAGCCCCGCAGGTCCCGGTG	88	249
	301015	5'UTR	3	23	CAACCGAGAAGGGCACTCAG	53	250
	301016	5'UTR	3	35	CCTCAGCGGCAGCAACCGAG	62	251
10	301017	5'UTR	3	36	TCCTCAGCGGCAGCAACCGA	47	252
	301018	5'UTR	3	37	CTCCTCAGCGGCAGCAACCG	45	253
	301019	5'UTR	3	39	GGCTCCTCAGCGGCAGCAAC	70	254
15	301020	5'UTR	3	43	GGCGGGCTCCTCAGCGGCA G	85	255
	301021	5'UTR	3	116	GGTCCATCGCCAGCTGCGGT	89	256
	301022	Start Codon	3	120	GGCGGGTCCATCGCCAGCTG	69	257
20	301023	Stop Codon	3	13800	TAGAGGATGATAGTAAGTTC	69	258
	301024	3'UTR	3	13824	AAATGAAGATTTCTTTTAAA	5	259
25	301025	3'UTR	3	13854	TATGTGAAAGTTCAATTGGA	76	260
	301026	3'UTR	3	13882	ATATAGGCAGTTTGAATTTT	57	261
	301027	3'UTR	3	13903	GCTCACTGTATGGTTTTATC	89	262
	301028	3'UTR	3	13904	GGCTCACTGTATGGTTTTAT	93	263
30	301029	3'UTR	3	13908	GGCTGGCTCACTGTATGGTT	90	264
	301030	3'UTR	3	13909	AGGCTGGCTCACTGTATGGT	90	265
	301031	3'UTR	3	13910	AAGGCTGGCTCACTGTATGG	90	266
35	301032	3'UTR	3	13917	CTACTGCAAGGCTGGCTCAC	63	267
	301033	3'UTR	3	13922	ACTGCCTACTGCAAGGCTGG	77	268
	301034	3'UTR	3	13934	TGCTTATAGTCTACTGCCTA	88	269
	301035	3'UTR	3	13937	TTCTGCTTATAGTCTACTGC	82	270
40	301036	3'UTR	3	13964	TTTGGTGCAGGTCCAGTTCA	88	271
	301037	3'UTR	3	13968	CAGCTTTGGTGCAGGTCCAG	90	272
	301038	3'UTR	3	13970	GCCAGCTTTGGTGCAGGTCC	86	273
45	301039	3'UTR	3	13974	TGGTGCCAGCTTTGGTGCAG	73	274
	301040	3'UTR	3	13978	GCCCTGGTGCCAGCTTTGGT	74	275
	301041	3'UTR	3	13997	GAGTTCAGAGACCTTCCGAG	85	276
	301042	3'UTR	3	14012	AAATGCCATCCTTCTGAGTT	81	277
50	301043	3'UTR	3	14014	AAAAATGCCATCCTTCTGAG	81	278
	301044	3'UTR	3	14049	AAAATAACTCAGATCCTGAT	76	279
	301045	3'UTR	3	14052	AGCAAATAACTCAGATCCT	90	280
55	301046	3'UTR	3	14057	AGTTTAGCAAATAACTCAG	80	281
	301047	3'UTR	3	14064	TCCCCAAGTTTAGCAAAT	56	282

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
301048	3'UTR	3	14071	TTCCTCCTCCCCCAAGTTTA	67	283
301217	3'UTR	3	14087	AGACTCCATTTATTTGTTCC	81	284

**Example 30**

**Antisense inhibition of apolipoprotein B - Gene walk**

**[0340]** In accordance with the present invention, a "gene walk" was conducted in which another series of oligonucleotides was designed to target the regions of the human apolipoprotein B RNA (GenBank accession number NM\_000384.1, incorporated herein as SEQ ID NO: 3) which are near the target site of SEQ ID Nos 224 or 247. The oligonucleotides are shown in Table 11. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 11 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Treatment doses were 50 nM and 150 nM and are indicated in Table 11. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 11 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Gene walk

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB 150 nM	% INHIB 50 nM	SEQ ID NO
308589	exon: exon junction	3	3230	CTTCTGCTTGAGTTACAAAC	94	20	285
308590	exon: exon junction	3	3232	ACCTTCTGCTTGAGTTACAA	98	26	286
308591	exon: exon junction	3	3234	GCACCTTCTGCTTGAGTTAC	92	76	287
308592	exon: exon junction	3	3236	TCGCACCTTCTGCTTGAGTT	96	49	288
308593	exon: exon junction	3	3238	CTTCGCACCTTCTGCTTGAG	80	41	289
308594	exon: exon junction	3	3240	TGCTTCGCACCTTCTGCTTG	88	57	290
308595	exon: exon junction	3	3242	TCTGCTTCGCACCTTCTGCT	82	60	291
308596	exon: exon junction	3	3244	AGTCTGCTTCGCACCTTCTG	94	81	292

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB 150 nM	% INHIB 50 nM	SEQ ID NO
308597	exon: exon junction	3	3246	TCAGTCTGCTTCGCACCTTC	91	66	293
308598	exon: exon junction	3	3248	CCTCAGTCTGCTTCGCACCT	85	59	294
308599	exon: exon junction	3	3250	AGCCTCAGTCTGCTTCGCA C	94	79	295
308600	coding	3	3252	GTAGCCTCAGTCTGCTTCG C	89	72	296
308601	coding	3	3254	TGGTAGCCTCAGTCTGCTTC	91	63	297
308602	coding	3	3256	CATGGTAGCCTCAGTCTGCT	92	83	298
308603	coding	3	3258	GTCATGGTAGCCTCAGTCT G	97	56	299
308604	coding	3	3260	ATGTCATGGTAGCCTCAGTC	90	73	300
308605	coding	3	3262	GAATGTCATGGTAGCCTCA G	81	50	301
308606	coding	3	3264	TTGAATGTCATGGTAGCCTC	97	54	302
308607	coding	3	3266	ATTTGAATGTCATGGTAGCC	77	9	303
308608	coding	3	3268	ATATTTGAATGTCATGGTAG	85	70	304
308609	coding	3	5582	CAGCCACATGCAGCTTCAG G	96	78	305
308610	coding	3	5584	ACCAGCCACATGCAGCTTC A	90	40	306
308611	coding	3	5586	TTACCAGCCACATGCAGCTT	95	59	307
308612	coding	3	5588	GGTTACCAGCCACATGCAG C	90	75	308
308613	coding	3	5590	TAGGTTACCAGCCACATGCA	87	43	309
308614	coding	3	5592	TTTAGGTTACCAGCCACATG	92	74	310
308615	coding	3	5594	CTTTTAGGTTACCAGCCACA	85	45	311
308616	coding	3	5596	TCCTTTTAGGTTACCAGCCA	81	39	312
308617	coding	3	5598	GCTCCTTTTAGGTTACCAGC	87	77	313
308618	coding	3	5600	AGGCTCCTTTTAGGTTACCA	77	61	314
308619	coding	3	5602	GTAGGCTCCTTTTAGGTTAC	74	69	315
308620	coding	3	5604	TGGTAGGCTCCTTTTAGGTT	88	69	316
308621	coding	3	5606	TTTGGTAGGCTCCTTTTAGG	91	56	317

[0341] As shown in Tables 10 and 11, SEQ ID Nos 124, 128, 129, 132, 133, 134, 138, 140, 141, 142, 144, 145, 147, 148, 149, 150, 152, 153, 155, 156, 158, 159, 160, 161, 163, 165, 166, 167, 169, 170, 172, 176, 177, 178, 181, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232,

233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, and 317 demonstrated at least 30% inhibition of human apolipoprotein B expression in this assay and are therefore preferred. More preferred are SEQ ID Nos 224, 247, and 262. The target regions to which these preferred sequences are complementary are herein referred to as "preferred target segments" and are therefore preferred for targeting by compounds of the present invention. These preferred target segments are shown in Table 18. The sequences represent the reverse complement of the preferred antisense compounds shown in Tables 10 and 11. "Target site" indicates the first (5'-most) nucleotide number on the particular target nucleic acid to which the oligonucleotide binds. Also shown in Table 18 is the species in which each of the preferred target segments was found.

**Example 31**

**Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap: Targeting GenBank Accession number M14162.1**

[0342] In accordance with the present invention, another series of oligonucleotides was designed to target different regions of the human apolipoprotein B RNA, using published sequence (GenBank accession number M14162.1, incorporated herein as SEQ ID NO: 318). The oligonucleotides are shown in Table 12. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 12 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments in which HepG2 cells were treated with 150 nM of the compounds in Table 12. If present, "N.D." indicates "no data".

Table 12 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
271009	coding	318	3121	GCCTCAGTCTGCTTCGCGCC	75	319
271024	coding	318	8031	GCTCACTGTTTCAGCATCTGG	27	320
271029	coding	318	8792	TGAGAATCTGGGCGAGGCC C	N.D.	321
271040	coding	318	11880	GTCCTTCATATTTGCCATCT	0	322
271047	coding	318	12651	CCTCCCTCATGAACATAGTG	32	323
281639	coding	318	9851	GACGTCAGAACCTATGATGG	38	324
281647	coding	318	12561	TGAGTGAGTCAATCAGCTTC	73	325

**Example 32**

**Antisense Inhibition of human apolipoprotein B - Gene walk targeting GenBank Accession number M14162.1**

[0343] In accordance with the present invention, a "gene walk" was conducted in which another series of oligonucleotides was designed to target the regions of the human apolipoprotein B RNA (GenBank accession number M14162.1, incorporated herein as SEQ ID NO: 318) which are near the target site of SEQ ID NO: 319. The oligonucleotides are shown in Table 13. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 13 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The



## EP 2 336 318 A1

internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Treatment doses were 50 nM and 150 nM and are indicated in Table 13. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 13 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB 150 nM	% INHIB 50 nM	SEQ ID NO
308622	coding	318	3104	GCCTTCTGCTTGAGTTACAA	87	25	326
308623	coding	318	3106	GCGCCTTCTGCTTGAGTTAC	71	62	327
308624	coding	318	3108	TCGCGCCTTCTGCTTGAGTT	89	69	328
308625	coding	318	3110	CTTCGCGCCTTCTGCTTGAG	83	64	329
308626	coding	318	3116	AGTCTGCTTCGCGCCTTCTG	94	38	330
308627	coding	318	3118	TCAGTCTGCTTCGCGCCTTC	89	67	331
308628	coding	318	3120	CCTCAGTCTGCTTCGCGCCT	92	61	332
308629	coding	318	3122	AGCCTCAGTCTGCTTCGCGC	95	77	333

[0344] As shown in Tables 12 and 13, SEQ ID Nos 319, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, and 333 demonstrated at least 30% inhibition of human apolipoprotein B expression in this assay and are therefore preferred. More preferred is SEQ ID NO: 319. The target regions to which these preferred sequences are complementary are herein referred to as "preferred target segments" and are therefore preferred for targeting by compounds of the present invention. These preferred target segments are shown in Table 18. The sequences represent the reverse complement of the preferred antisense compounds shown in Tables 12 and 13. "Target site" indicates the first (5'-most) nucleotide number on the particular target nucleic acid to which the oligonucleotide binds. Also shown in Table 18 is the species in which each of the preferred target segments was found.

### Example 33

#### Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Targeting the Genomic sequence

[0345] In accordance with the present invention, another series of oligonucleotides was designed to target different regions of the human apolipoprotein B RNA, using published sequence (the complement of nucleotides 39835 to 83279 of the sequence with GenBank accession number NT\_022227.9, representing a genomic sequence, incorporated herein as SEQ ID NO: 334). The oligonucleotides are shown in Table 14. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 14 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxy-nucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments in which HepG2 cells were treated with 150 nM of the oligonucleotides in Table 14. If present, "N.D." indicates "no data".

EP 2 336 318 A1

Table 14 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
301049	intron: exon junction	334	904	TCTGTAAGACAGGAGAAAGA	41	335
301050	intron: exon junction	334	913	ATTCCTCTTCTGTAAGACA	22	336
301051	exon: intron junction	334	952	GATGCCTTACTTGGACAGAC	27	337
301052	intron	334	1945	AGAAATAGCTCTCCCAAGGA	13	338
301053	intron: exon junction	334	1988	GTCGCATCTTCTAACGTGGG	45	339
301054	exon: intron junction	334	2104	TCCTCCATACCTTGCAGTTG	0	340
301055	intron	334	2722	TGGCTCATGTCTACCATATT	49	341
301056	intron	334	2791	CAGTTGAAATGCAGCTAATG	35	342
301057	intron	334	3045	TGCAGACTAGGAGTGAAAGT	30	343
301058	intron	334	3117	AGGAGGATGTCCTTTTATTG	27	344
301059	intron	334	3290	ATCAGAGCACCAAAGGGAAT	12	345
301060	intron: exon junction	334	3381	CCAGCTCAACCTGAGAATTC	17	346
301061	exon: intron junction	334	3527	CATGACTTACCTGGACATGG	52	347
301062	intron	334	3566	CCTCAGCGGACACACACACA	21	348
301063	intron	334	3603	GTCACATCCGTGCCTGGTGC	41	349
301064	intron	334	3864	CAGTGCCTCTGGGACCCAC	60	350
301065	intron	334	3990	AGCTGCAGTGGCCGATCAGC	50	351
301066	intron	334	4251	GACCTCCCAGCCACGTGGA	61	352
301067	intron	334	4853	TCTGATCACCATACATTACA	45	353
301068	intron	334	5023	ATTTCCCACTGGGTA CTCTC	44	354
301069	intron	334	5055	GGCTGAAGCCCATGCTGACT	44	355
301070	intron	334	5091	GTTGGACAGTCATTCTTTTG	38	356
301071	intron	334	5096	CACTTGTTGGACAGTCATTC	48	357
301072	intron	334	5301	ATTTTAAATTACAGTAGATA	43	358
301073	intron	334	5780	CTGTTCTCCACCCATATCAG	37	359

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
301074	intron: exon junction	334	6353	GAGCTCATACCTGTCCCAGA	75	360
301075	intron	334	6534	TTCAAGGGCCACTGCTATCA	52	361
301076	intron	334	6641	CCAGTATTTACGCCAATCC	36	362
301077	intron	334	6661	GGCAGGAGGAACCTCGGGC A	55	363
301078	intron	334	6721	TTTTAAAATTAGACCCAACC	22	364
301079	intron	334	6727	TGACTGTTTTAAAATTAGAC	20	365
301080	intron	334	6788	CCCAGCAAACACAGGTGAAG	25	366
301081	intron	334	7059	GAGTGTGGTCTTGCTAGTGC	46	367
301082	intron	334	7066	CTATGCAGAGTGTGGTCTTG	41	368
301083	intron	334	7189	AGAAGATGCAACCACATGTA	29	369
301084	intron: exon junction	334	7209	ACACGGTATCCTATGGAGGA	49	370
301085	exon: intron junction	334	7365	TGGGACTTACCATGCCTTTG	11	371
301086	intron	334	7702	GGTTTTGCTGCCCTACATCC	30	372
301087	intron	334	7736	ACAAGGAGTCCTTGTGCAGA	40	373
301088	intron	334	8006	ATGTTCACTGAGACAGGCTG	41	374
301089	intron	334	8215	GAAGGTCCATGGTTCATCTG	0	375
301090	intron	334	8239	ATTAGACTGGAAGCATCCTG	39	376
301091	intron	334	8738	GAGATTGGAGACGAGCATTT	35	377
301092	exon: intron junction	334	8881	CATGACCTACTTGTAGGAGA	22	378
301093	intron	334	9208	TGGATTTGGATACACAAGTT	42	379
301094	intron	334	9244	ACTCAATATATATTCATTGA	22	380
301095	intron	334	9545	CAAGGAAGCACACCATGTCA	38	381
301096	intron: exon junction	334	9563	ATACTTATTCCTGGTAACCA	24	382
301097	intron	334	9770	GGTAGCCAGAACACCAGTGT	50	383
301098	intron	334	9776	ACTAGAGGTAGCCAGAACAC	34	384
301099	intron	334	10149	ACCACCTGACATCACAGGTT	24	385
301100	intron	334	10341	TACTGTGACCTATGCCAGGA	55	386
301101	intron	334	10467	GGAGGTGCTACTGTTGACAT	42	387
301102	intron	334	10522	TCCAGACTTGTCTGAGTCTA	47	388

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO	
5	301103	intron	334	10547	TCTAAGAGGTAGAGCTAAAG	7	389
	301104	intron	334	10587	CCAGAGATGAGCAACTTAGG	38	390
	301105	intron	334	10675	GGCCATGTAAATTGCTCATC	7	391
10	301106	intron	334	10831	AAAGAAACTATCCTGTATTC	12	392
	301107	intron: exon junction	334	10946	TTCTTAGTACCTGGAAGATG	23	393
15	301108	exon: intron junction	334	11166	CATTAGATACCTGGACACCT	29	394
	301109	intron	334	11337	GTTTCATGGAACCTCAGCGCA	44	395
20	301110	intron	334	11457	CTGGAGAGCACCTGCAATAG	35	396
	301111	intron	334	11521	TGAAGGGTAGAGAAATCATA	9	397
	301112	exon: intron junction	334	12111	GGAAACTCACTTGTTGACCG	25	398
25	301113	intron	334	12155	AGGTGCAAGATGTTCTCTG	46	399
	301114	intron	334	12162	TGCACAGAGGTGCAAGATGT	16	400
	301115	intron	334	12221	CACAAGAGTAAGGAGCAGAG	39	401
30	301116	intron	334	12987	GATGGATGGTGAGAAATTAC	33	402
	301117	intron	334	13025	TAGACAATTGAGACTCAGAA	39	403
	301118	intron	334	13057	ATGTGCACACAAGGACATAG	33	404
35	301119	intron	334	13634	ACATACAAATGGCAATAGGC	33	405
	301120	intron	334	13673	TAGGCAAAGGACATGAATAG	30	406
	301121	coding	334	14448	TTATGATAGCTACAGAATAA	29	407
40	301122	exon: intron junction	334	14567	CTGAGATTACCCGCAGAATC	32	408
	301123	intron	334	14587	GATGTATGTCATATAAAAGA	26	409
45	301124	intron: exon junction	334	14680	TTCCAATGACCTGCATTGA	48	410
	301125	intron	334	15444	AGGGATGGTCAATCTGGTAG	57	411
	301126	intron	334	15562	GGCTAATAAATAGGGTAGTT	22	412
50	301127	intron	334	15757	TCCTAGAGCACTATCAAGTA	41	413
	301128	intron: exon junction	334	15926	CCTCCTGGTCCTGCAGTCAA	56	414
55	301129	intron	334	16245	CATTTGCACAAGTGTGGTT	35	415
	301130	intron	334	16363	CTGACACACCATGTTATTAT	10	416

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO	
5	301131	intron: exon junction	334	16399	CTTTTTCAGACTAGATAAGA	0	417
10	301132	exon: intron junction	334	16637	TCACACTTACCTCGATGAGG	29	418
	301133	intron	334	17471	AAGAAAATGGCATCAGGTTT	13	419
15	301134	intron: exon junction	334	17500	CCAAGCCAATCTGAGAAAGA	25	420
20	301135	exon: intron junction	334	17677	AAATACACACCTGCTCATGT	20	421
	301136	exon: intron junction	334	17683	CTTCACAAATACACACCTGC	20	422
25	301137	intron	334	18519	AGTGGAAGTTTGGTCTCATT	41	423
	301138	intron	334	18532	TTGCTAGCTTCAAAGTGGAA	44	424
	301139	intron	334	18586	TCAAGAATAAGCTCCAGATC	41	425
	301140	intron	334	18697	GCATACAAGTCACATGAGGT	34	426
30	301141	intron	334	18969	TACAAGGTGTTTCTTAAGAA	38	427
	301142	intron	334	19250	ATGCAGCCAGGATGGGCCTA	54	428
	301143	intron:	334	19340	TTACCATATCCTGAGAGTTT	55	429
35		exon junction					
	301144	intron	334	19802	GCAAAGGTAGAGGAAGGTAT	32	430
	301145	intron	334	19813	AAGGACCTTCAGCAAAGGTA	36	431
40	301146	intron	334	20253	CATAGGAGTACATTTATATA	23	432
	301147	intron	334	20398	ATTATGATAAAAATCAATTTT	19	433
	301148	intron	334	20567	AGAAATTTCACTAGATAGAT	31	434
45	301149	intron	334	20647	AGCATATTTTGATGAGCTGA	44	435
	301150	intron	334	20660	GAAAGGAAGGACTAGCATAT	39	436
	301151	intron: exon junction	334	20772	CCTCTCCAATCTGTAGACCC	28	437
50	301152	intron	334	21316	CTGGATAACTCAGACCTTTG	40	438
	301153	intron	334	21407	AGTCAGAAAACAACCTATTC	11	439
55	301154	intron: exon junction	334	21422	CAGCCTGCATCTATAAGTCA	31	440

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO	
5	301155	exon: intron junction	334	21634	AAAGAATTACCCTCCACTGA	33	441
10	301156	intron	334	21664	TCTTTCAAACCTGGCTAGGCA	39	442
	301157	intron	334	21700	GCCTGGCAAATCTGCAGG	37	443
	301158	intron	334	22032	CTACCTCAAATCAATATGTT	28	444
	301159	intron	334	22048	TGCTTTACCTACCTAGCTAC	36	445
15	301160	intron	334	22551	ACCTTGTGTGTCTCACTCAA	49	446
	301161	intron	334	22694	ATGCATTCCCTGACTAGCAC	34	447
	301162	intron	334	22866	CATCTCTGAGCCCCTTACCA	24	448
20	301163	intron	334	22903	GCTGGGCATGCTCTCTCCCC	51	449
	301164	intron	334	22912	GCTTTTCGAGCTGGGCATGC	55	450
	301165	intron	334	23137	ACTCCTTTCTATACCTGGCT	47	451
	301166	intron	334	23170	ATTCTGCCTCTTAGAAAGTT	38	452
25	301167	intron	334	23402	CCAAGCCTCTTACTGGGCT	29	453
	301168	intron	334	23882	CACTCATGACCAGACTAAGA	35	454
	301169	intron	334	23911	ACCTCCCAGAAGCCTTCCAT	22	455
30	301170	intron	334	24184	TTCATATGAAATCTCCTACT	40	456
	301171	intron	334	24425	TATTTAATTTACTGAGAAAC	7	457
	301172	intron: exon junction	334	24559	TAATGTGTTGCTGGTGAAGA	35	458
35	301173	exon: intron junction	334	24742	CATCTCTAACCTGGTGTCCC	21	459
40	301174	intron	334	24800	GTGCCATGCTAGGTGGCCAT	37	460
	301175	intron	334	24957	AGCAAATTGGGATCTGTGCT	29	461
	301176	intron	334	24991	TCTGGAGGCTCAGAAACATG	57	462
	301177	intron	334	25067	TGAAGACAGGGAGCCACCTA	40	463
45	301178	intron	334	25152	AGGATTCCCAAGACTTTGGA	38	464
	301179	intron: exon junction	334	25351	CAGCTCTAATCTAAAGACAT	22	465
50	301180	exon: intron junction	334	25473	GAATACTCACCTTCTGCTTG	6	466
	301181	intron	334	26047	ATCTCTCTGTCCTCATCTTC	28	467
55	301182	intron	334	26749	CCAACTCCCCCTTTCTTTGT	37	468
	301183	intron	334	26841	TCTGGGCCAGGAAGACACGA	68	469

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
301184	intron	334	27210	TATTGTGTGCTGGGCACTGC	52	470
301185	intron: exon junction	334	27815	TGCTTCGCACCTGGACGAGT	51	471
301186	exon: intron junction	334	28026	CCTTCTTTACCTTAGGTGGC	37	472
301187	intron	334	28145	GCTCTCTCTGCCACTCTGAT	47	473
301188	intron	334	28769	AACTTCTAAAGCCAACATTC	27	474
301189	intron: exon junction	334	28919	TGTGTCACAACCTATGGTAAA	63	475
301190	exon: intron junction	334	29095	AGACACATACCATAATGCCA	22	476
301191	intron: exon junction	334	29204	TTCTCTTCATCTGAAAATAC	21	477
301192	intron	334	29440	TGAGGATGTAATTAGCACTT	27	478
301193	intron: exon junction	334	29871	AGCTCATTGCCTACAAAATG	31	479
301194	intron	334	30181	GTTCTCATGTTTACTAATGC	40	480
301195	intron	334	30465	GAATTGAGACAACCTTGATTT	26	481
301196	intron: exon junction	334	30931	CCGGCCATCGCTGAAATGAA	54	482
301197	exon: intron junction	334	31305	CATAGCTCACCTTGACATT	28	483
301198	intron	334	31325	CGGTGCACCCTTTACCTGAG	28	484
301199	intron: exon junction	334	31813	TCTCCAGATCCTAACATAAA	19	485
301200	intron	334	39562	TTGAATGACACTAGATTTTC	37	486
301201	intron	334	39591	AAAATCCATTTTCTTTAAAG	12	487
301202	intron	334	39654	CAGCTCACACTTATTTTAAA	7	488
301203	intron: exon junction	334	39789	GTTCCCAAACTGTATAGGA	36	489
301204	exon: intron junction	334	39904	AGCTCCATACTGAAGTCCTT	37	490

EP 2 336 318 A1

(continued)

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
301205	intron	334	39916	CAATTCAATAAAAAGCTCCAT	31	491
301206	intron	334	39938	GTTTTCAAAAAGGTATAAGGT	28	492
301207	intron: exon junction	334	40012	TTCCCATTCCCTGAAAGCAG	13	493
301208	exon: intron junction	334	40196	TGGTATTTACCTGAGGGCTG	21	494
301209	intron	334	40412	ATAAATAATAGTGCTGATGG	39	495
301210	intron	334	40483	CTATGGCTGAGCTTGCCTAT	33	496
301211	intron	334	40505	CTCTCTGAAAAATATACCCT	17	497
301212	intron	334	40576	TTGATGTATCTCATCTAGCA	41	498
301213	intron	334	40658	TAGAACCATGTTTGGTCTTC	35	499
301214	intron	334	40935	TTTCTCTTTATCACATGCC	29	500
301215	intron	334	41066	TATAGTACACTAAAACCTCA	1	501
301216	intron: exon junction	334	41130	CTGGAGAGGACTAAACAGAG	49	502

[0346] As shown in Table 14, SEQ ID Nos 335, 339, 341, 342, 343, 347, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 367, 368, 370, 372, 373, 374, 376, 377, 379, 381, 383, 384, 386, 387, 388, 390, 395, 396, 399, 401, 402, 403, 404, 405, 406, 408, 410, 411, 413, 414, 415, 423, 424, 425, 426, 427, 428, 429, 430, 431, 434, 435, 436, 438, 440, 441, 442, 443, 445, 446, 447, 449, 450, 451, 452, 454, 456, 458, 460, 462, 463, 464, 468, 469, 470, 471, 472, 473, 475, 479, 480, 482, 486, 489, 490, 491, 495, 496, 498, 499, and 502 demonstrated at least 30% inhibition of human apolipoprotein B expression in this assay and are therefore preferred. The target regions to which these preferred sequences are complementary are herein referred to as "preferred target segments" and are therefore preferred for targeting by compounds of the present invention. These preferred target segments are shown in Table 18. The sequences represent the reverse complement of the preferred antisense compounds shown in Table 14. "Target site" indicates the first (5'-most) nucleotide number on the particular target nucleic acid to which the oligonucleotide binds. Also shown in Table 18 is the species in which each of the preferred target segments was found.

**Example 34**

**Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Targeting GenBank accession number AI249040.1**

[0347] In accordance with the present invention, another series of oligonucleotides was designed to target different regions of the human apolipoprotein B RNA, using published sequence (the complement of the sequence with GenBank accession number AI249040.1, incorporated herein as SEQ ID NO: 503). The oligonucleotides are shown in Table 15. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 15 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MQE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments in which HepG2 cells were treated with 150 nM of the oligonucleotides in Table 15. If present, "N.D." indicates "no data".



Table 15 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQIDNO
301218	3'UTR	503	484	ACATTTTATCAATGCCCTCG	23	504
301219	3'UTR	503	490	GCCAGAACATTTTATCAATG	35	505
301220	3'UTR	503	504	AGAGGTTTTGCTGTGCCAG A	51	506
301221	3'UTR	503	506	CTAGAGGTTTTGCTGTGCCA	61	507
301222	3'UTR	503	507	TCTAGAGGTTTTGCTGTGCC	14	508
301223	3'UTR	503	522	AATCACACTATGTGTTCTAG	26	509
301224	3'UTR	503	523	AAATCACACTATGTGTTCTA	33	510
301225	3'UTR	503	524	TAAATCACACTATGTGTTCT	3	511
301226	3'UTR	503	526	CTTAAATCACACTATGTGTT	39	512
301227	3'UTR	503	536	TATTCTGTTACTTAAATCAC	23	513

**[0348]** As shown in Table 15, SEQ ID Nos 505, 506, 507, 510, and 512 demonstrated at least 30% inhibition of human apolipoprotein B expression in this assay and are therefore preferred. The target regions to which these preferred sequences are complementary are herein referred to as "preferred target segments" and are therefore preferred for targeting by compounds of the present invention. These preferred target segments are shown in Table 18. The sequences represent the reverse complement of the preferred antisense compounds shown in Table 15. "Target site" indicates the first (5'-most) nucleotide number on the particular target nucleic acid to which the oligonucleotide binds. Also shown in Table 18 is the species in which each of the preferred target segments was found.

### Example 35

#### Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Variation in position of the gap

**[0349]** In accordance with the present invention, a series of antisense compounds was designed to target different regions of the human apolipoprotein B RNA, using published sequences (GenBank accession number NM\_000384.1, incorporated herein as SEQ ID NO: 3). The compounds are shown in Table 16. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the compound binds. All compounds in Table 16 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length. The "gap" region consists of 2'-deoxynucleotides, which is flanked on one or both sides (5' and 3' directions) by "wings" composed of 2'-methoxyethyl (2'-MOE)nucleotides. The number of 2'-MOE nucleotides on either side of the gap varies such that the total number of 2'-MOE nucleotides always equals 10 and the total length of the chimeric oligonucleotide is 20 nucleotides. The exact structure of each oligonucleotide is designated in Table 16 as the "gap structure" and the 2'-deoxynucleotides are in bold type. A designation of 8-10-2, for instance, indicates that the first (5'-most) 8 nucleotides and the last (3'-most) 2 nucleotides are 2'-MOE nucleotides and the 10 nucleotides in the gap are 2'-deoxynucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels by quantitative real-time PCR as described in other examples herein. Data, shown in Table 16, are averages from three experiments in which HepG2 cells were treated with the antisense oligonucleotides of the present invention at doses of 50 nM and 150 nM. If present, "N.D." indicates "no data".

EP 2 336 318 A1

Table 16 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a variable deoxy gap

ISIS #	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB 150 nM	% INHIB 50 nM	gap structure	SEQ ID NO
308631	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	94	74	0-10-10	224
308632	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	97	41	0-10-10	247
308634	3	5589	AGGTTACCAGCCACATGCAG	67	45	10-10-0	224
308635	3	3249	GCCTCAGTCTGCTTCGCACC	93	69	10-10-0	247
308637	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	95	79	1-10-9	224
308638	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	94	91	1-10-9	247
308640	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	96	76	2-10-8	224
308641	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	89	77	2-10-8	247
308643	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	96	56	3-10-7	224
308644	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	93	71	3-10-7	247
308646	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	76	50	4-10-6	224
308647	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	86	53	4-10-6	247
308649	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	91	68	6-10-4	224
308650	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	94	74	6-10-4	247
308652	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	95	73	7-10-3	224
308653	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	89	73	7-10-3	247
308655	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	83	84	8-10-2	224
308656	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	97	37	8-10-2	247
308658	3	5589	<b>AGGTTACCAGCCACATGCAG</b>	78	86	9-10-1	224
308659	3	3249	<b>GCCTCAGTCTGCTTCGCACC</b>	93	70	9-10-1	247
308660	3	3254	<b>TGGTAGCCTCAGTCTGCTTC</b>	92	72	2-10-8	514
308662	3	3254	<b>TGGTAGCCTCAGTCTGCTTC</b>	83	76	8-10-2	514

[0350] As shown in Table 16, SEQ ID Nos 224, 247, and 514 demonstrated at least 30% inhibition of human apolipoprotein B expression in this assay at both doses. These data suggest that the oligonucleotides are effective with a number of variations in the gap placement. The target regions to which these preferred sequences are complementary are herein referred to as "preferred target segments" and are therefore preferred for targeting by compounds of the present invention. These preferred target segments are shown in Table 18. The sequences represent the reverse complement of the preferred antisense compounds shown in Table 16. "Target site" indicates the first (5'-most) nucleotide number on the particular target nucleic acid to which the oligonucleotide binds. Also shown in Table 18 is the species in which each of the preferred target segments was found.

**Example 36**

**Antisense inhibition of human apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Variation in position of the gap of SEQ ID Nos: 319 and 515**

[0351] In accordance with the present invention, a series of antisense compounds was designed based on SEQ ID Nos 319 and 515, with variations in the gap structure. The compounds are shown in Table 17. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the compound binds. All compounds in Table 17 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length. The "gap" region consists of 2'-deoxynucleotides, which is flanked on one or both sides (5' and 3' directions) by "wings" composed of 2'-methoxyethyl (2'-MOE)

EP 2 336 318 A1

nucleotides. The number of 2'-MOE nucleotides on either side of the gap varies such that the total number of 2'-MOE nucleotides always equals 10 and the total length of the chimeric oligonucleotide is 20 nucleotides. The exact structure of each oligonucleotide is designated in Table 17 as the "gap structure" and the 2'-deoxynucleotides are in bold type. A designation of 8-10-2, for instance, indicates that the first (5'-most) 8 nucleotides and the last (3'-most) 2 nucleotides are 2'-MOE nucleotides and the 10 nucleotides in the gap are 2'-deoxynucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human apolipoprotein B mRNA levels by quantitative real-time PCR as described in other examples herein. Data, shown in Table 17, are averages from three experiments in which HepG2 cells were treated with the antisense oligonucleotides of the present invention at doses of 50 nM and 150 nM. If present, "N.D." indicates "no data".

Table 17 Inhibition of human apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a variable deoxy gap

ISIS #	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB 150 nM	% INHIB 50 nM	gap structure	SEQ ID NO
308630	318	3121	<b>GCCTCAGTCTGCTTCGCGCC</b>	89	69	0-10-10	319
308633	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	83	66	10-10-0	319
308636	318	3121	<b>GCCTCAGTCTGCTTCGCGCC</b>	91	81	1-10-9	319
308639	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	94	86	2-10-8	319
308642	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	95	85	3-10-7	319
308645	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	98	57	4-10-6	319
308648	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	89	78	6-10-4	319
308651	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	88	87	7-10-3	319
308654	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	90	81	8-10-2	319
308657	318	3121	GCCTCAGTCT <b>GCTTCGCGCC</b>	78	61	9-10-1	319
308661	318	3116	<b>AGTCTGCTTCGCGCCTTCTG</b>	91	70	2-10-8	515
308663	318	3116	<b>AGTCTGCTTCGCGCCTTCTG</b>	84	44	8-10-2	515

[0352] As shown in Table 17, SEQ ID Nos 319 and 515 demonstrated at least 44% inhibition of human apolipoprotein B expression in this assay for either dose. These data suggest that the compounds are effective with a number of variations in gap placement. The target regions to which these preferred sequences are complementary are herein referred to as "preferred target segments" and are therefore preferred for targeting by compounds of the present invention. These preferred target segments are shown in Table 18. The sequences represent the reverse complement of the preferred antisense compounds shown in Table 17. "Target site" indicates the first (5'-most) nucleotide number on the particular target nucleic acid to which the oligonucleotide binds. Also shown in Table 18 is the species in which each of the preferred target segments was found.

Table 18 Sequence and position of preferred target segments identified in apolipoprotein B.

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
187342	3	199	GCGCCAGGGCCGAAGAGGAA	124	<i>H. sapiens</i>	516
187346	3	509	CAGGTATGAGCTCAAGCTGG	128	<i>H. sapiens</i>	517
187347	3	584	CATCCTGAACATCAAGAGGG	129	<i>H. sapiens</i>	518
187350	3	756	GGCAGTGTGATCGCTTCAA	132	<i>H. sapiens</i>	519
187351	3	799	CACTTGCTCTCATCAAAGGC	133	<i>H. sapiens</i>	520

EP 2 336 318 A1

(continued)

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
187352	3	869	CACACTGGACGCTAAGAGGA	134	<i>H. sapiens</i>	521
187356	3	1459	CGCTGAGCCACGCGGTCAAC	138	<i>H. sapiens</i>	522
187358	3	1859	TGTCCAAATTCTACCATGGG	140	<i>H. sapiens</i>	523
187359	3	2179	CAGCTGACCTCATCGAGATT	141	<i>H. sapiens</i>	524
187360	3	2299	GTCAAGTTCCTGATGGTGTGTC	142	<i>H. sapiens</i>	525
187362	3	2518	AGCTGCTTCTGATGGGTGCC	144	<i>H. sapiens</i>	526
187363	3	2789	GGGCATCATCATTCCGGACT	145	<i>H. sapiens</i>	527
187365	3	3100	CCTACTATCCGCTGACCGGG	147	<i>H. sapiens</i>	528
187367	3	3449	GGGCCACCTAAGTTGTGACA	148	<i>H. sapiens</i>	529
187368	3	3919	AGAACATGGGATTGCCAGAC	149	<i>H. sapiens</i>	530
187369	3	4089	CTCCACTTCAAGTCTGTGGG	150	<i>H. sapiens</i>	531
187371	3	5146	CAGAGCTTGGCCTCTCTGGG	152	<i>H. sapiens</i>	532
187372	3	5189	TGGCCGCTTCAGGGAACACA	153	<i>H. sapiens</i>	533
187374	3	6049	CAGCTGAGCAGACAGGCACC	155	<i>H. sapiens</i>	534
187375	3	6520	GGGAGAGACAAGTTTCACAT	156	<i>H. sapiens</i>	535
187377	3	6859	GTA CTGCATCCTGGATTCAA	158	<i>H. sapiens</i>	536
187378	3	7459	GTGAGGTGACTCAGAGACTC	159	<i>H. sapiens</i>	537
187379	3	7819	TTGCAGAGCAATATTCTATC	160	<i>H. sapiens</i>	538
187380	3	7861	AAGCATTGGTAGAGCAAGGG	161	<i>H. sapiens</i>	539
187383	3	8589	CCGCTGGCTCTGAAGGAGTC	163	<i>H. sapiens</i>	540
187385	3	8829	TCTAGTCAGGCTGACCTGCG	165	<i>H. sapiens</i>	541
187387	3	9119	GGGCCACAGTGTCTAACTG	166	<i>H. sapiens</i>	542
187388	3	10159	AATCAAGTGTCATCACACTG	167	<i>H. sapiens</i>	543
187390	3	10349	GGGTAGTCATAACAGTACTG	169	<i>H. sapiens</i>	544
187391	3	10699	AGAGCACACGGTCTTCAGTG	170	<i>H. sapiens</i>	545
187393	3	10839	TTACAGCTAGAGGGCCTCTT	172	<i>H. sapiens</i>	546
187398	3	12149	CACCGTGGGCATGGATATGG	176	<i>H. sapiens</i>	547
187399	3	12265	GGGAATCTGATGAGGAAACT	177	<i>H. sapiens</i>	548
187400	3	12380	TGTCAACAAGTACCACTGGG	178	<i>H. sapiens</i>	549
187403	3	12749	ACCTGGGATATACTAGGG	181	<i>H. sapiens</i>	550
187406	3	13299	CCAAGTATAGTTGGCTGGAC	183	<i>H. sapiens</i>	551
187407	3	13779	TACATGAAGCTTGCTCCAGG	184	<i>H. sapiens</i>	552
197724	3	229	ATGTCAGCCTGGTCTGTCCA	185	<i>H. sapiens</i>	553
197725	3	269	GCACCTCCGGAAGTACACAT	186	<i>H. sapiens</i>	554
197726	3	389	CTGCAGCTTCATCCTGAAGA	187	<i>H. sapiens</i>	555
197727	3	449	TGAGGGCAAAGCCTTGCTGA	188	<i>H. sapiens</i>	556

EP 2 336 318 A1

(continued)

	SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
5	197728	3	529	CCATTCCAGAAGGGAAGCAG	189	<i>H. sapiens</i>	557
	197729	3	709	CGAGGAAGGGCAATGTGGCA	190	<i>H. sapiens</i>	558
10	197730	3	829	CCTTGTCAACTCTGATCAGC	191	<i>H. sapiens</i>	559
	197731	3	849	AGCAGCCAGTCCTGTGAGTA	192	<i>H. sapiens</i>	560
	197732	3	889	AGCATGTGGCAGAAGCCATC	193	<i>H. sapiens</i>	561
15	197733	3	1059	GAGAGCACCAAATCCACATC	194	<i>H. sapiens</i>	562
	197734	3	1199	CCTCAGTGATGAAGCAGTCA	195	<i>H. sapiens</i>	563
	197735	3	1349	GATAGATGTGGTCACCTACC	196	<i>H. sapiens</i>	564
	197736	3	1390	CCTCAGCACAGCAGCTGCGA	197	<i>H. sapiens</i>	565
20	197737	3	1589	GATTCTGCGGGTCATTGGAA	198	<i>H. sapiens</i>	566
	197738	3	1678	CAAAGCCATCACTGATGATC	199	<i>H. sapiens</i>	567
	197739	3	1699	AGAAAGCTGCCATCCAGGCT	200	<i>H. sapiens</i>	568
25	197740	3	1749	CAGGAGGTTCTTCTTCAGAC	201	<i>H. sapiens</i>	569
	197741	3	1829	GAGTCCTTACAGGCAGATA	202	<i>H. sapiens</i>	570
	197742	3	1919	TGCCAATATCTTGAAGTCTAG	203	<i>H. sapiens</i>	571
	197743	3	2189	CATCGAGATTGGCTTGGAAAG	204	<i>H. sapiens</i>	572
30	197744	3	2649	GGAGCTGGATTACAGTTGCA	205	<i>H. sapiens</i>	573
	197745	3	2729	CAACATGCAGGCTGAACTGG	206	<i>H. sapiens</i>	574
	197746	3	2949	ACATTACATTTGGTCTCTAC	207	<i>H. sapiens</i>	575
35	197747	3	3059	CTCAGGCGCTTACTCCAACG	208	<i>H. sapiens</i>	576
	197748	3	3118	GGGACACCAGATTAGAGCTG	209	<i>H. sapiens</i>	577
	197749	3	3189	GAGCTCCAGAGAGAGGACAG	210	<i>H. sapiens</i>	578
	197750	3	3289	ATCGGCAGAGTATGACCTTG	211	<i>H. sapiens</i>	579
40	197751	3	3488	CAAGGGTGTATTTCATAC	212	<i>H. sapiens</i>	580
	197752	3	3579	GACTCATCTGCTACAGCTTA	213	<i>H. sapiens</i>	581
	197753	3	4039	GCAAATCCTCCAGAGATCTA	214	<i>H. sapiens</i>	582
45	197754	3	4180	CTCTCCTGGGTGTTCTAGAC	215	<i>H. sapiens</i>	583
	197755	3	4299	ATGAAGGCTGACTCTGTGGT	216	<i>H. sapiens</i>	584
	197756	3	4511	GGGACCACAGATGTCTGCTT	217	<i>H. sapiens</i>	585
	197757	3	4660	CTGGCCGGCTCAATGGAGAG	218	<i>H. sapiens</i>	586
50	197758	3	4919	GCTGCGTTCTGAATATCAGG	219	<i>H. sapiens</i>	587
	197759	3	5009	TGCTGACATCTTAGGCACTG	220	<i>H. sapiens</i>	588
	197760	3	5109	AAGTGTAGTCTCCTGGTGCT	221	<i>H. sapiens</i>	589
55	197761	3	5212	CAAAATTCAGTCTGGATGGG	222	<i>H. sapiens</i>	590
	197762	3	5562	GGGAAACTACGGCTAGAACC	223	<i>H. sapiens</i>	591
	197763	3	5589	CTGCATGTGGCTGGTAACCT	224	<i>H. sapiens</i>	592

EP 2 336 318 A1

(continued)

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
197764	3	5839	CCATGACCATCGATGCACAT	225	<i>H. sapiens</i>	593
197765	3	5869	ATGGGAAACTCGCTCTCTGG	226	<i>H. sapiens</i>	594
197766	3	5979	AGTCATCATCTCGTGTCTAG	227	<i>H. sapiens</i>	595
197767	3	6099	GAATACAGCCAGGACTTGGGA	228	<i>H. sapiens</i>	596
197768	3	6144	GGCGTGGAGCTTACTGGACG	229	<i>H. sapiens</i>	597
197769	3	6249	GAGATGAGAGATGCCGTTGA	230	<i>H. sapiens</i>	598
197770	3	6759	AGTCTTGATGAGCACTATCA	231	<i>H. sapiens</i>	599
197771	3	6889	CTAAGTACCAAATCAGAATC	232	<i>H. sapiens</i>	600
197772	3	7149	GTCCATGAGTTAATCGAGAG	233	<i>H. sapiens</i>	601
197773	3	7549	AGGCCACAGTTGCAGTGTAT	234	<i>H. sapiens</i>	602
197774	3	7779	TCTGATTGGTGGACTCTTGC	235	<i>H. sapiens</i>	603
197775	3	7929	GAAGTCAGTCTTCAGGCTCT	236	<i>H. sapiens</i>	604
197776	3	8929	CCAGATTCTCAGATGAGGGA	237	<i>H. sapiens</i>	605
197778	3	10240	CATCTGTCATTGATGCACTG	238	<i>H. sapiens</i>	606
197779	3	10619	AGGAGATGTCAAGGGTTCGG	239	<i>H. sapiens</i>	607
197780	3	10659	GGAECTATTGCTAGTGAGGC	240	<i>H. sapiens</i>	608
197781	3	10899	CTCTCTCCATGGCAAATGTC	241	<i>H. sapiens</i>	609
197783	3	11979	CACCGTGACTTCAGTGCAGA	243	<i>H. sapiens</i>	610
197784	3	12249	ACTGAGTTGAGGGTCCGGGA	244	<i>H. sapiens</i>	611
197786	3	13958	CACATATGAACTGGACCTGC	245	<i>H. sapiens</i>	612
197787	3	14008	TCTGAACTCAGAAGGATGGC	246	<i>H. sapiens</i>	613
216825	3	3249	GGTGCGAAGCAGACTGAGGC	247	<i>H. sapiens</i>	614
216826	3	3	TCCCACCGGGACCTGCGGGG	248	<i>H. sapiens</i>	615
216827	3	6	CACCGGGACCTGCGGGGCTG	249	<i>H. sapiens</i>	616
216828	3	23	CTGAGTGCCCTTCTCGGTTG	250	<i>H. sapiens</i>	617
216829	3	35	CTCGGTTGCTGCCGCTGAGG	251	<i>H. sapiens</i>	618
216830	3	36	TCCGTTGCTGCCGCTGAGGA	252	<i>H. sapiens</i>	619
216831	3	37	CGGTTGCTGCCGCTGAGGAG	253	<i>H. sapiens</i>	620
216832	3	39	GTTGCTGCCGCTGAGGAGCC	254	<i>H. sapiens</i>	621
216833	3	43	CTGCCGCTGAGGAGCCCGCC	255	<i>H. sapiens</i>	622
216834	3	116	ACCGCAGCTGGCGATGGACC	256	<i>H. sapiens</i>	623
216835	3	120	CAGCTGGCGATGGACCCGCC	257	<i>H. sapiens</i>	624
216836	3	13800	GAACTTACTATCATCTCTA	258	<i>H. sapiens</i>	625
216838	3	13854	TCCAATTGAACTTTACATA	260	<i>H. sapiens</i>	626
216839	3	13882	AAAATTCAAACCTGCCTATAT	261	<i>H. sapiens</i>	627
216840	3	13903	GATAAAACCATACAGTGAGC	262	<i>H. sapiens</i>	628

EP 2 336 318 A1

(continued)

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
216841	3	13904	ATAAAACCATACAGTGAGCC	263	<i>H. sapiens</i>	629
216842	3	13908	AACCATACAGTGAGCCAGCC	264	<i>H. sapiens</i>	630
216843	3	13909	ACCATACAGTGAGCCAGCCT	265	<i>H. sapiens</i>	631
216844	3	13910	CCATACAGTGAGCCAGCCTT	266	<i>H. sapiens</i>	632
216845	3	13917	GTGAGCCAGCCTTGCAGTAG	267	<i>H. sapiens</i>	633
216846	3	13922	CCAGCCTTGCAGTAGGCAGT	268	<i>H. sapiens</i>	634
216847	3	13934	TAGGCAGTAGACTATAAGCA	269	<i>H. sapiens</i>	635
216848	3	13937	GCAGTAGACTATAAGCAGAA	270	<i>H. sapiens</i>	636
216849	3	13964	TGAACTGGACCTGCACCAAA	271	<i>H. sapiens</i>	637
216850	3	13968	CTGGACCTGCACCAAAGCTG	272	<i>H. sapiens</i>	638
216851	3	13970	GGACCTGCACCAAAGCTGGC	273	<i>H. sapiens</i>	639
216852	3	13974	CTGCACCAAAGCTGGCACCA	274	<i>H. sapiens</i>	640
216853	3	13978	ACCAAAGCTGGCACCAGGGC	275	<i>H. sapiens</i>	641
216854	3	13997	CTCGGAAGGTCTCTGAACTC	276	<i>H. sapiens</i>	642
216855	3	14012	AACTCAGAAGGATGGCATT	277	<i>H. sapiens</i>	643
216856	3	14014	CTCAGAAGGATGGCATT	278	<i>H. sapiens</i>	644
216857	3	14049	ATCAGGATCTGAGTTAT	279	<i>H. sapiens</i>	645
216858	3	14052	AGGATCTGAGTTATTTGCT	280	<i>H. sapiens</i>	646
216859	3	14057	CTGAGTTATTTGCTAACT	281	<i>H. sapiens</i>	647
216860	3	14064	ATTTTGCTAACTTGGGGGA	282	<i>H. sapiens</i>	648
216861	3	14071	TAACTTGGGGGAGGAGGAA	283	<i>H. sapiens</i>	649
217030	3	14087	GGAACAAATAAATGGAGTCT	284	<i>H. sapiens</i>	650
224316	3	3230	GTTTGTAACTCAAGCAGAAG	285	<i>H. sapiens</i>	651
224317	3	3232	TTGTAACTCAAGCAGAAGGT	286	<i>H. sapiens</i>	652
224318	3	3234	GTA ACTCAAGCAGAAGGTGC	287	<i>H. sapiens</i>	653
224319	3	3236	AACTCAAGCAGAAGGTGCGA	288	<i>H. sapiens</i>	654
224320	3	3238	CTCAAGCAGAAGGTGCGAAG	289	<i>H. sapiens</i>	655
224321	3	3240	CAAGCAGAAGGTGCGAAGCA	290	<i>H. sapiens</i>	656
224322	3	3242	AGCAGAAGGTGCGAAGCAGA	291	<i>H. sapiens</i>	657
224323	3	3244	CAGAAGGTGCGAAGCAGACT	292	<i>H. sapiens</i>	658
224324	3	3246	GAAGGTGCGAAGCAGACTGA	293	<i>H. sapiens</i>	659
224325	3	3248	AGGTGCGAAGCAGACTGAGG	294	<i>H. sapiens</i>	660
224326	3	3250	GTGCGAAGCAGACTGAGGCT	295	<i>H. sapiens</i>	661
224327	3	3252	GCGAAGCAGACTGAGGCTAC	296	<i>H. sapiens</i>	662
224328	3	3254	GAACCAGACTGAGGCTACCA	297	<i>H. sapiens</i>	663
224329	3	3256	AGCAGACTGAGGCTACCATG	298	<i>H. sapiens</i>	664

EP 2 336 318 A1

(continued)

	SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
5	224330	3	3258	CAGACTGAGGCTACCATGAC	299	<i>H. sapiens</i>	665
	224331	3	3260	GACTGAGGCTACCATGACAT	300	<i>H. sapiens</i>	666
10	224332	3	3262	CTGAGGCTACCATGACATTC	301	<i>H. sapiens</i>	667
	224333	3	3264	GAGGCTACCATGACATTCAA	302	<i>H. sapiens</i>	668
	224334	3	3266	GGCTACCATGACATTCAAAT	303	<i>H. sapiens</i>	669
15	224335	3	3268	CTACCATGACATTCAAATAT	304	<i>H. sapiens</i>	670
	224336	3	5582	CCTGAAGCTGCATGTGGCTG	305	<i>H. sapiens</i>	671
	224337	3	5584	TGAAGCTGCATGTGGCTGGT	306	<i>H. sapiens</i>	672
	224338	3	5586	AAGCTGCATGTGGCTGGTAA	307	<i>H. sapiens</i>	673
20	224339	3	5588	GCTGCATGTGGCTGGTAACC	308	<i>H. sapiens</i>	674
	224340	3	5590	TGCATGTGGCTGGTAACCTA	309	<i>H. sapiens</i>	675
	224341	3	5592	CATGTGGCTGGTAACCTAAA	310	<i>H. sapiens</i>	676
25	224342	3	5594	TGTGGCTGGTAACCTAAAAG	311	<i>H. sapiens</i>	677
	224343	3	5596	TGGCTGGTAACCTAAAAGGA	312	<i>H. sapiens</i>	678
	224344	3	5598	GCTGGTAACCTAAAAGGAGC	313	<i>H. sapiens</i>	679
	224345	3	5600	TGGTAACCTAAAAGGAGCCT	314	<i>H. sapiens</i>	680
30	224346	3	5602	GTAACCTAAAAGGAGCCTAC	315	<i>H. sapiens</i>	681
	224347	3	5604	AACCTAAAAGGAGCCTACCA	316	<i>H. sapiens</i>	682
	224348	3	5606	CCTAAAAGGAGCCTACCAA	317	<i>H. sapiens</i>	683
35	187366	318	3121	GGCGCGAAGCAGACTGAGGC	319	<i>H. sapiens</i>	684
	187404	318	12651	CACTATGTTTCATGAGGGAGG	323	<i>H. sapiens</i>	685
	197777	318	9851	CCATCATAGGTTCTGACGTC	324	<i>H. sapiens</i>	686
	197785	318	12561	GAAGCTGATTGACTCACTCA	325	<i>H. sapiens</i>	687
40	224349	318	3104	TTGTAACTCAAGCAGAAGGC	326	<i>H. sapiens</i>	688
	224350	318	3106	GTAACTCAAGCAGAAGGCGC	327	<i>H. sapiens</i>	689
	224351	318	3108	AACTCAAGCAGAAGGCGCGA	328	<i>H. sapiens</i>	690
45	224352	318	3110	CTCAAGCAGAAGGCGCGAAG	329	<i>H. sapiens</i>	691
	224353	318	3116	CAGAAGGCGCGAAGCAGACT	330	<i>H. sapiens</i>	692
	224354	318	3118	GAAGGCGCGAAGCAGACTGA	331	<i>H. sapiens</i>	693
	224355	318	3120	AGGCGCGAAGCAGACTGAGG	332	<i>H. sapiens</i>	694
50	224356	318	3122	GCGCGAAGCAGACTGAGGCT	333	<i>H. sapiens</i>	695
	224328	3	3254	GAAGCAGACTGAGGCTACCA	514	<i>H. sapiens</i>	696
	224353	318	3116	CAGAAGGCGCGAAGCAGACT	515	<i>H. sapiens</i>	697
55	216862	334	904	TCTTTCTCCTGTCTTACAGA	335	<i>H. sapiens</i>	698
	216866	334	1988	CCCACGTTAGAAGATGCGAC	339	<i>H. sapiens</i>	699
	216868	334	2722	AATATGGTAGACATGAGCCA	341	<i>H. sapiens</i>	700



EP 2 336 318 A1

(continued)

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
216869	334	2791	CATTAGCTGCATTTCAACTG	342	<i>H. sapiens</i>	701
216870	334	3045	ACTTTCCTCCTAGTCTGCA	343	<i>H. sapiens</i>	702
216874	334	3527	CCATGTCCAGGTAAGTCATG	347	<i>H. sapiens</i>	703
216876	334	3603	GCACCAGGCACGGATGTGAC	349	<i>H. sapiens</i>	704
216877	334	3864	GTGGGGTCCCAGAGGCACTG	350	<i>H. sapiens</i>	705
216878	334	3990	GCTGATCGGCCACTGCAGCT	351	<i>H. sapiens</i>	706
216879	334	4251	TCCACGTGGCTGGGAGGTC	352	<i>H. sapiens</i>	707
216880	334	4853	TGTAATGTATGGTGATCAGA	353	<i>H. sapiens</i>	708
216881	334	5023	GAGAGTACCCAGTGGGAAAT	354	<i>H. sapiens</i>	709
216882	334	5055	AGTCAGCATGGGCTTCAGCC	355	<i>H. sapiens</i>	710
216883	334	5091	CAAAAGAATGACTGTCCAAC	356	<i>H. sapiens</i>	711
216884	334	5096	GAATGACTGTCCAACAAGTG	357	<i>H. sapiens</i>	712
216885	334	5301	TATCTACTGTAATTTAAAAT	358	<i>H. sapiens</i>	713
216886	334	5780	CTGATATGGGTGGAGAACAG	359	<i>H. sapiens</i>	714
216887	334	6353	TCTGGACAGGTATGAGCTC	360	<i>H. sapiens</i>	715
216888	334	6534	TGATAGCAGTGGCCCTTGAA	361	<i>H. sapiens</i>	716
216889	334	6641	GGATTGGCGTGAATACTGG	362	<i>H. sapiens</i>	717
216890	334	6661	TGCCCGAGGTTCTCCTGCC	363	<i>H. sapiens</i>	718
216894	334	7059	GCACTAGCAAGACCACACTC	367	<i>H. sapiens</i>	719
216895	334	7066	CAAGACCACACTCTGCATAG	368	<i>H. sapiens</i>	720
216897	334	7209	TCCTCCATAGGATACCGTGT	370	<i>H. sapiens</i>	721
216899	334	7702	GGATGTAGGGCAGCAAACC	372	<i>H. sapiens</i>	722
216900	334	7736	TCTGCACAAGGACTCCTTGT	373	<i>H. sapiens</i>	723
216901	334	8006	CAGCCTGTCTCAGTGAACAT	374	<i>H. sapiens</i>	724
216903	334	8239	CAGGATGCTTCCAGTCTAAT	376	<i>H. sapiens</i>	725
216904	334	8738	AAATGCTCGTCTCCAATCTC	377	<i>H. sapiens</i>	726
216906	334	9208	AACTTGTGTATCCAAATCCA	379	<i>H. sapiens</i>	727
216908	334	9545	TGACATGGTGTGCTTCCTTG	381	<i>H. sapiens</i>	728
216910	334	9770	ACACTGGTGTCTGGCTACC	383	<i>H. sapiens</i>	729
216911	334	9776	GTGTTCTGGCTACCTCTAGT	384	<i>H. sapiens</i>	730
216913	334	10341	TCCTGGCATAGGTCACAGTA	386	<i>H. sapiens</i>	731
216914	334	10467	ATGTCAACAGTAGCACCTCC	387	<i>H. sapiens</i>	732
216915	334	10522	TAGACTCAGACAAGTCTGGA	388	<i>H. sapiens</i>	733
216917	334	10587	CCTAAGTTGCTCATCTCTGG	390	<i>H. sapiens</i>	734
216922	334	11337	TGCGCTGAGTTCATGAAAC	395	<i>H. sapiens</i>	735
216923	334	11457	CTATTGCAGGTGCTCTCCAG	396	<i>H. sapiens</i>	736

EP 2 336 318 A1

(continued)

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
216926	334	12155	CAGAGGAACATCTTGACCT	399	<i>H. sapiens</i>	737
216928	334	12221	CTCTGCTCCTTACTCTTGTG	401	<i>H. sapiens</i>	738
216929	334	12987	GTAATTTCTCACCATCCATC	402	<i>H. sapiens</i>	739
216930	334	13025	TTCTGAGTCTCAATTGTCTA	403	<i>H. sapiens</i>	740
216931	334	13057	CTATGTCCTTGTGTGCACAT	404	<i>H. sapiens</i>	741
216932	334	13634	GCCTATTGCCATTTGTATGT	405	<i>H. sapiens</i>	742
216933	334	13673	CTATTCATGTCCTTGCCTA	406	<i>H. sapiens</i>	743
216935	334	14567	GATTCTGCGGGTAATCTCAG	408	<i>H. sapiens</i>	744
216937	334	14680	TCAATGCAGGTCATTGGAAA	410	<i>H. sapiens</i>	745
216938	334	15444	CTACCAGATTGACCATCCCT	411	<i>H. sapiens</i>	746
216940	334	15757	TACTTGATAGTGCTCTAGGA	413	<i>H. sapiens</i>	747
216941	334	15926	TTGACTGCAGGACCAGGAGG	414	<i>H. sapiens</i>	748
216942	334	16245	AACAAACACTTGTGCAAATG	415	<i>H. sapiens</i>	749
216950	334	18519	AATGAGACCAAACCTCCACT	423	<i>H. sapiens</i>	750
216951	334	18532	TTCCACTTTGAAGCTAGCAA	424	<i>H. sapiens</i>	751
216952	334	18586	GATCTGGAGCTTATTCCTGA	425	<i>H. sapiens</i>	752
216953	334	18697	ACCTCATGTGACTTGTATGC	426	<i>H. sapiens</i>	753
216954	334	18969	TTCTTAAGAAACACCTTGTA	427	<i>H. sapiens</i>	754
216955	334	19250	TAGGCCATCCTGGCTGCAT	428	<i>H. sapiens</i>	755
216956	334	19340	AAACTCTCAGGATATGGTAA	429	<i>H. sapiens</i>	756
216957	334	19802	ATACCTTCCTCTACCTTTGC	430	<i>H. sapiens</i>	757
216958	334	19813	TACCTTTGCTGAAGGTCCTT	431	<i>H. sapiens</i>	758
216961	334	20567	ATCTATCTAGTGAAATTTCT	434	<i>H. sapiens</i>	759
216962	334	20647	TCAGCTCATCAAAATATGCT	435	<i>H. sapiens</i>	760
216963	334	20660	ATATGCTAGTCCTTCCTTTC	436	<i>H. sapiens</i>	761
216965	334	21316	CAAAGGTCTGAGTTATCCAG	438	<i>H. sapiens</i>	762
216967	334	21422	TGACTTATAGATGCAGGCTG	440	<i>H. sapiens</i>	763
216968	334	21634	TCAGTGGAGGGTAATTCCTT	441	<i>H. sapiens</i>	764
216969	334	21664	TGCCTAGCCAGTTTGAAAGA	442	<i>H. sapiens</i>	765
216970	334	21700	CCTGCAGAATTTTGCCAGGC	443	<i>H. sapiens</i>	766
216972	334	22048	GTAGCTAGGTAGGTAAAGCA	445	<i>H. sapiens</i>	767
216973	334	22551	TTGAGTGAGACACACAAGGT	446	<i>H. sapiens</i>	768
216974	334	22694	GTGCTAGTCAGGGAATGCAT	447	<i>H. sapiens</i>	769
216976	334	22903	GGGGAGAGAGCATGCCCAGC	449	<i>H. sapiens</i>	770
216977	334	22912	GCATGCCCAGCTGCGAAAGC	450	<i>H. sapiens</i>	771
216978	334	23137	AGCCAGGTATAGAAAGGAGT	451	<i>H. sapiens</i>	772

EP 2 336 318 A1

(continued)

SITE ID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID NO	ACTIVE IN	SEQ ID NO
216979	334	23170	AACTTTCTAAGAGGCAGAAT	452	<i>H. sapiens</i>	773
216981	334	23882	TCTTAGTCTGGTCATGAGTG	454	<i>H. sapiens</i>	774
216983	334	24184	AGTAGGAGATTTTCATATGAA	456	<i>H. sapiens</i>	775
216985	334	24559	TCTTCACCAGCAACACATTA	458	<i>H. sapiens</i>	776
216987	334	24800	ATGGCCACCTAGCATGGCAC	460	<i>H. sapiens</i>	777
216989	334	24991	CATGTTTCTGAGCCTCCAGA	462	<i>H. sapiens</i>	778
216990	334	25067	TAGGTGGCTCCCTGTCTTCA	463	<i>H. sapiens</i>	779
216991	334	25152	TCCAAAGTCTTGGGAATCCT	464	<i>H. sapiens</i>	780
216995	334	26749	ACAAAGAAAGGGGGAGTTGG	468	<i>H. sapiens</i>	781
216996	334	26841	TCGTGTCTTCTGGCCCAGA	469	<i>H. sapiens</i>	782
216997	334	27210	GCAGTGCCCAGCACACAATA	470	<i>H. sapiens</i>	783
216998	334	27815	ACTCGTCCAGGTGCGAAGCA	471	<i>H. sapiens</i>	784
216999	334	28026	GCCACCTAAGGTAAAGAAGG	472	<i>H. sapiens</i>	785
217000	334	28145	ATCAGAGTGGCAGAGAGAGC	473	<i>H. sapiens</i>	786
217002	334	28919	TTTACCATAGTTGTGACACA	475	<i>H. sapiens</i>	787
217006	334	29871	CATTTTGTAGGCAATGAGCT	479	<i>H. sapiens</i>	788
217007	334	30181	GCATTAGTAAACATGAGAAC	480	<i>H. sapiens</i>	789
217009	334	30931	TTCATTTAGCGATGGCCGG	482	<i>H. sapiens</i>	790
217013	334	39562	GAAAATCTAGTGTCATTCAA	486	<i>H. sapiens</i>	791
217016	334	39789	TCCTATACAGTTTTGGGAAC	489	<i>H. sapiens</i>	792
217017	334	39904	AAGGACTTCAGTATGGAGCT	490	<i>H. sapiens</i>	793
217018	334	39916	ATGGAGCTTTTATTGAATTG	491	<i>H. sapiens</i>	794
217022	334	40412	CCATCAGCACTATTATTTAT	495	<i>H. sapiens</i>	795
217023	334	40483	ATAGGCAAGCTCAGCCATAG	496	<i>H. sapiens</i>	796
217025	334	40576	TGCTAGATGAGATACATCAA	498	<i>H. sapiens</i>	797
217026	334	40658	GAAGACCAAACATGGTTCTA	499	<i>H. sapiens</i>	798
217029	334	41130	CTCTGTTTAGTCCTCTCCAG	502	<i>H. sapiens</i>	799
217032	503	490	CATTGATAAAATGTTCTGGC	505	<i>H. sapiens</i>	800
217033	503	504	TCTGGCACAGCAAAACCTCT	506	<i>H. sapiens</i>	801
217034	503	506	TGGCACAGCAAAACCTCTAG	507	<i>H. sapiens</i>	802
217037	503	523	TAGAACACATAGTGTGATTT	510	<i>H. sapiens</i>	803
217039	503	526	AACACATAGTGTGATTTAAG	512	<i>H. sapiens</i>	804

**[0353]** As these "preferred target segments" have been found by experimentation to be open to, and accessible for, hybridization with the antisense compounds of the present invention, one of skill in the art will recognize or be able to ascertain, using no more than routine experimentation, further embodiments of the invention that encompass other compounds that specifically hybridize to these preferred target segments and consequently inhibit the expression of apolipoprotein B.

**[0354]** According to the present invention, antisense compounds include antisense oligomeric compounds, antisense oligonucleotides, ribozymes, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other short oligomeric compounds which hybridize to at least a portion of the target nucleic acid.

### 5 Example 37

#### Antisense inhibition of human apolipoprotein B expression - dose response of oligonucleotides

**[0355]** In accordance with the present invention, 12 oligonucleotides described in Examples 29 and 31 were further investigated in a dose response study. The control oligonucleotides used in this study were ISIS 18076 (SEQ ID NO: 805) and ISIS 13650 (SEQ ID NO: 806).

**[0356]** All compounds in this study, including the controls, were chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotides. All cytidine residues are 5-methylcytidines.

**[0357]** In the dose-response experiment, with mRNA levels as the endpoint, HepG2 cells were treated with the antisense oligonucleotides or the control oligonucleotides at doses of 37, 75, 150, and 300 nM oligonucleotide. Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from two experiments with mRNA levels in the treatment groups being normalized to an untreated control group. The data are shown in Table 19.

Table 19 Inhibition of apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Dose Response

ISIS #	Dose				SEQ ID NO
	37 nM	75 nM	150 nM	300 nM	
	% inhibition				
271009	82	91	94	96	319
281625	62	76	84	94	224
301014	75	90	96	98	249
301027	80	90	95	96	262
301028	70	79	85	92	263
301029	54	67	79	85	264
301030	64	75	87	92	265
301031	61	82	92	96	266
301034	73	87	93	97	269
301036	67	83	92	95	271
301037	73	85	89	96	272
301045	77	86	94	98	280

### Example 38

#### Antisense inhibition of human apolipoprotein B expression - dose response - Lower dose range

**[0358]** In accordance with the present invention, seven oligonucleotides described in Examples 29, 31, 35, and 36 were further investigated in a dose response study. The control oligonucleotides used in this study were ISIS 18076 (SEQ ID NO: 805), ISIS 13650 (SEQ ID NO: 806), and ISIS 129695 (SEQ ID NO: 807).

**[0359]** All compounds in this study, including the controls, were chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotides. All cytidine residues are 5-methylcytidines.

**[0360]** In the dose-response experiment, with mRNA levels as the endpoint, HepG2 cells were treated with the anti-sense oligonucleotides or the control oligonucleotides at doses of 12.5, 37, 75, 150, and 300 nM oligonucleotide. Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from two experiments with mRNA levels in the treatment groups being normalized to an untreated control group. The data are shown in Table 20.

Table 20 Inhibition of apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap - Dose Response

ISIS #	Dose					SEQ ID #
	12.5	37 nM	75 nM	150 nM	300 nM	
	% inhibition					
271009	67	86	92	94	95	319
281625	44	66	83	85	94	224
301012	63	79	90	92	95	247
308638	42	73	91	96	97	247
308642	59	84	91	97	98	319
308651	57	76	84	90	88	319
308658	29	61	73	78	90	224

### Example 39

#### RNA Synthesis

**[0361]** In general, RNA synthesis chemistry is based on the selective incorporation of various protecting groups at strategic intermediary reactions. Although one of ordinary skill in the art will understand the use of protecting groups in organic synthesis, a useful class of protecting groups includes silyl ethers. In particular bulky silyl ethers are used to protect the 5'-hydroxyl in combination with an acid-labile orthoester protecting group on the 2'-hydroxyl. This set of protecting groups is then used with standard solid-phase synthesis technology. It is important to lastly remove the acid labile orthoester protecting group after all other synthetic steps. Moreover, the early use of the silyl protecting groups during synthesis ensures facile removal when desired, without undesired deprotection of 2' hydroxyl.

**[0362]** Following this procedure for the sequential protection of the 5'-hydroxyl in combination with protection of the 2'-hydroxyl by protecting groups that are differentially removed and are differentially chemically labile, RNA oligonucleotides were synthesized.

**[0363]** RNA oligonucleotides are synthesized in a stepwise fashion. Each nucleotide is added sequentially (3'- to 5'-direction) to a solid support-bound oligonucleotide. The first nucleoside at the 3'-end of the chain is covalently attached to a solid support. The nucleotide precursor, a ribonucleoside phosphoramidite, and activator are added, coupling the second base onto the 5'-end of the first nucleoside. The support is washed and any unreacted 5'-hydroxyl groups are capped with acetic anhydride to yield 5'-acetyl moieties. The linkage is then oxidized to the , more stable and ultimately desired P(V) linkage. At the end of the nucleotide addition cycle, the 5'-silyl group is cleaved with fluoride. The cycle is repeated for each subsequent nucleotide.

**[0364]** Following synthesis, the methyl protecting groups on the phosphates are cleaved in 30 minutes utilizing 1 M disodium-2-carbamoyl-2-cyanoethylene-1,1-dithiolate trihydrate ( $S_2Na_2$ ) in DMF. The deprotection solution is washed from the solid support-bound oligonucleotide using water. The support is then treated with 40% methylamine in water for 10 minutes at 55 °C. This releases the RNA oligonucleotides into solution, deprotects the exocyclic amines, and modifies the 2'- groups. The oligonucleotides can be analyzed by anion exchange HPLC at this stage.

**[0365]** The 2'-orthoester groups are the last protecting groups to be removed. The ethylene glycol monoacetate orthoester protecting group developed by Dharmacon Research, Inc. (Lafayette, CO), is one example of a useful orthoester protecting group which, has the following important properties. It is stable to the conditions of nucleoside phosphoramidite synthesis and oligonucleotide synthesis. However, after oligonucleotide synthesis the oligonucleotide is treated with methylamine which not only cleaves the oligonucleotide from the solid support but also removes the acetyl groups from the orthoesters. The resulting 2-ethyl-hydroxyl substituents on the orthoester are less electron withdrawing than the acetylated precursor. As a result, the modified orthoester becomes more labile to acid-catalyzed hydrolysis. Specifically, the rate of cleavage is approximately 10 times faster after the acetyl groups are removed. Therefore, this

orthoester possesses sufficient stability in order to be compatible with oligonucleotide synthesis and yet, when subsequently modified, permits deprotection to be carried out under relatively mild aqueous conditions compatible with the final RNA oligonucleotide product.

**[0366]** Additionally, methods of RNA synthesis are well known in the art (Scaringe, S. A. Ph.D. Thesis, University of Colorado, 1996; Scaringe, S. A., et al., J. Am. Chem. Soc., 1998, 120, 11820-11821; Matteucci, M. D. and Caruthers, M. H. J. Am. Chem. Soc., 1981, 103, 3185-3191; Beaucage, S. L. and Caruthers, M. H. Tetrahedron Lett., 1981, 22, 1859-1862; Dahl, B. J., et al., Acta Chem. Scand., 1990, 44, 639-641; Reddy, M. P., et al., Tetrahedron Lett., 1994, 25, 4311-4314; Wincott, F. et al., Nucleic Acids Res., 1995, 23, 2677-2684; Griffin, B. E., et al., Tetrahedron, 1967, 23, 2301-2313; Griffin, B. E., et al., Tetrahedron, 1967, 23, 2315-2331).


**[0367]** RNA antisense compounds (RNA oligonucleotides) of the present invention can be synthesized by the methods herein or purchased from Dharmacon Research, Inc (Lafayette, CO). Once synthesized, complementary RNA antisense compounds can then be stably annealed by methods known in the art to form double stranded (duplexed) antisense compounds. For example, duplexes can be formed by combining 30 µl of each of the complementary strands of RNA oligonucleotides (50 µM RNA oligonucleotide solution) and 15 µl of 5X annealing buffer (100 mM potassium acetate, 30 mM HEPES-KOH pH 7.4, 2 mM magnesium acetate) followed by heating for 1 minute at 90°C, then 1 hour at 37°C. The resulting duplexed antisense compounds can be used in kits, assays, screens, or other methods to investigate the role of a target nucleic acid.

#### Example 40


##### Design and screening of duplexed antisense compounds targeting apolipoprotein B

**[0368]** In accordance with the present invention, a series of nucleic acid duplexes comprising the antisense compounds of the present invention and their complements are designed to target apolipoprotein B. The nucleobase sequence of the antisense strand of the duplex comprises at least a portion of an oligonucleotide described herein. The ends of the strands may be modified by the addition of one or more natural or modified nucleobases to form an overhang. The sense strand of the dsRNA is then designed and synthesized as the complement of the antisense strand and may also contain modifications or additions to either terminus. For example, in one embodiment, both strands of the dsRNA duplex would be complementary over the central nucleobases, each having overhangs at one or both termini. The antisense and sense strands of the duplex comprise from about 17 to 25 nucleotides, or from about 19 to 23 nucleotides. Alternatively, the antisense and sense strands comprise 20, 21 or 22 nucleotides.

**[0369]** For example, a duplex comprising an antisense strand having the sequence CGAGAGGCGGACGGGACCG and having a two-nucleobase overhang of deoxythymidine(dT) would have the following structure:



**[0370]** In another embodiment, a duplex comprising an antisense strand having the same sequence CGAGAGGCGGACGGGACCG may be prepared with blunt ends (no single stranded overhang) as shown:



**[0371]** RNA strands of the duplex can be synthesized by methods disclosed herein or purchased from Dharmacon Research Inc., (Lafayette, CO). Once synthesized, the complementary strands are stably annealed. The single strands are aliquoted and diluted to a concentration of 50 µM. Once diluted, 30 µL of each strand is combined with 15µL of a 5X solution of annealing buffer. The final concentration of said buffer is 100 mM potassium acetate, 30 mM HEPES-KOH pH 7.4, and 2mM magnesium acetate. The final volume is 75 µL. This solution is incubated for 1 minute at 90°C and then centrifuged for 15 seconds. The tube is allowed to sit for 1 hour at 37°C at which time the dsRNA duplexes are used in experimentation. The final concentration of the dsRNA duplex is 20 µM. This solution can be stored frozen (-20°C) and freeze-thawed up to 5 times.

**[0372]** Once prepared, the duplexed antisense compounds are evaluated for their ability to modulate apolipoprotein

B expression.

**[0373]** When cells reached 80% confluency, they are treated with duplexed antisense compounds of the invention. For cells grown in 96-well plates, wells are washed once with 200  $\mu$ L OPTI-MEM-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM-1 containing 12  $\mu$ g/mL LIPOFECTIN (Gibco BRL) and the desired duplex antisense compound at a final concentration of 200 nM. After 5 hours of treatment, the medium is replaced with fresh medium. Cells are harvested 16 hours after treatment, at which time RNA is isolated and target reduction measured by RT-PCR.

#### Example 41

#### Design of phenotypic assays and *in vivo* studies for the use of apolipoprotein B inhibitors

##### *Phenotypic assays*

**[0374]** Once apolipoprotein B inhibitors have been identified by the methods disclosed herein, the compounds are further investigated in one or more phenotypic assays, each having measurable endpoints predictive of efficacy in the treatment of a particular disease state or condition. Phenotypic assays, kits and reagents for their use are well known to those skilled in the art and are herein used to investigate the role and/or association of apolipoprotein B in health and disease. Representative phenotypic assays, which can be purchased from any one of several commercial vendors, include those for determining cell viability, cytotoxicity, proliferation or cell survival (Molecular Probes, Eugene, OR; PerkinElmer, Boston, MA), protein-based assays including enzymatic assays (Panvera, LLC, Madison, WI; BD Biosciences, Franklin Lakes, NJ; Oncogene Research Products, San Diego, CA), cell regulation, signal transduction, inflammation, oxidative processes and apoptosis (Assay Designs Inc., Ann Arbor, MI), triglyceride accumulation (Sigma-Aldrich, St. Louis, MO), angiogenesis assays, tube formation assays, cytokine and hormone assays and metabolic assays (Chemicon International Inc., Temecula, CA; Amersham Biosciences, Piscataway, NJ).

**[0375]** In one non-limiting example, cells determined to be appropriate for a particular phenotypic assay (i.e., MCF-7 cells selected for breast cancer studies; adipocytes for obesity studies) are treated with apolipoprotein B inhibitors identified from the *in vitro* studies as well as control compounds at optimal concentrations which are determined by the methods described above. At the end of the treatment period, treated and untreated cells are analyzed by one or more methods specific for the assay to determine phenotypic outcomes and endpoints.

**[0376]** Phenotypic endpoints include changes in cell morphology over time or treatment dose as well as changes in levels of cellular components such as proteins, lipids, nucleic acids, hormones, saccharides or metals. Measurements of cellular status which include pH, stage of the cell cycle, intake or excretion of biological indicators by the cell, are also endpoints of interest.

**[0377]** Analysis of the genotype of the cell (measurement of the expression of one or more of the genes of the cell) after treatment is also used as an indicator of the efficacy or potency of the apolipoprotein B inhibitors. Hallmark genes, or those genes suspected to be associated with a specific disease state, condition, or phenotype, are measured in both treated and untreated cells.

##### *In vivo studies*

**[0378]** The individual subjects of the *in vivo* studies described herein are warm-blooded vertebrate animals, which includes humans.

**[0379]** The clinical trial is subjected to rigorous controls to ensure that individuals are not unnecessarily put at risk and that they are fully informed about their role in the study.

**[0380]** To account for the psychological effects of receiving treatments, volunteers are randomly given placebo or apolipoprotein B inhibitor. Furthermore, to prevent the doctors from being biased in treatments, they are not informed as to whether the medication they are administering is a apolipoprotein B inhibitor or a placebo. Using this randomization approach, each volunteer has the same chance of being given either the new treatment or the placebo.

**[0381]** Volunteers receive either the apolipoprotein B inhibitor or placebo for eight week period with biological parameters associated with the indicated disease state or condition being measured at the beginning (baseline measurements before any treatment), end (after the final treatment), and at regular intervals during the study period. Such measurements include the levels of nucleic acid molecules encoding apolipoprotein B or apolipoprotein B protein levels in body fluids, tissues or organs compared to pre-treatment levels. Other measurements include, but are not limited to, indices of the disease state or condition being treated, body weight, blood pressure, serum titers of pharmacologic indicators of disease or toxicity as well as ADME (absorption, distribution, metabolism and excretion) measurements.

**[0382]** Information recorded for each patient includes age (years), gender, height (cm), family history of disease state or condition (yes/no), motivation rating (some/moderate/great) and number and type of previous treatment regimens for

the indicated disease or condition.

[0383] Volunteers taking part in this study are healthy adults (age 18 to 65 years) and roughly an equal number of males and females participate in the study. Volunteers with certain characteristics are equally distributed for placebo and apolipoprotein B inhibitor treatment. In general, the volunteers treated with placebo have little or no response to treatment, whereas the volunteers treated with the apolipoprotein B inhibitor show positive trends in their disease state or condition index at the conclusion of the study.

#### Example 42

#### Antisense inhibition of rabbit apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

[0384] In accordance with the present invention, a series of oligonucleotides was designed to target different regions of rabbit apolipoprotein B, using published sequences (GenBank accession number X07480.1, incorporated herein as SEQ ID NO: 808, GenBank accession number M17780.1, incorporated herein as SEQ ID NO: 809, and a sequence was derived using previously described primers (Tanaka, Journ. Biol. Chem., 1993, 268, 12713-12718) representing an mRNA of the rabbit apolipoprotein B, incorporated herein as SEQ ID NO: 810). The oligonucleotides are shown in Table 21. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 21 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on rabbit apolipoprotein B mRNA levels in primary rabbit hepatocytes by quantitative real-time PCR as described in other examples herein. Primary rabbit hepatocytes were treated with 150 nM of the compounds in Table 21. For rabbit apolipoprotein B the PCR primers were:

forward primer: AAGCACCCCAATGTCACC (SEQ ID NO: 811)

reverse primer: GGGATGGCAGAGCCAATGTA (SEQ ID NO: 812) and the PCR probe was: FAM- TCCTGGAT-TCAAGCTTCTATGTGCCTTCA - TAMRA (SEQ ID NO: 813) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 21 Inhibition of rabbit apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
233149	808	1	TGCTTGGAGAAGGTAAGATC	0	814
233150	810	1	GCGTTGTCTCCGATGTTCTG	20	815
233151	809	13	TAATCATTAAGTGTGTGG	20	816
233152	808	22	TCAGCACGTAGCAATGCATT	0	817
233153	808	31	GCCTGATACTCAGCACGTAG	0	818
233154	809	31	CAATTGAATGTAAGTGTGTGG	18	819
233155	808	51	ACCTCAGTGACTTGTAAATCA	47	820
233156	809	51	CACTGGAACTTGTCTCTCC	23	821
233157	809	71	AGTAGTTAGTTTCTCCTTGG	0	822
233159	808	121	TCAGTGCCCAAGATGTCAGC	0	823
233160	810	121	ATTGGAATAATGTATCCAGG	81	824
233161	809	130	TTGGCATTATCCAATGCAGT	28	825
233162	808	151	GTTGCCTTGTGAGCAGCAGT	0	826
233163	810	151	ATTGTGAGTGGAGATACTTC	80	827



EP 2 336 318 A1

(continued)

ISIS #	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
233164	809	171	CATATGTCTGAAGTTGAGAC	8	828
233165	808	181	GTAGATACTCCATTTTGGCC	0	829
233166	810	181	GGATCACATGACTGAATGCT	82	830
233167	808	201	TCAAGCTGGTTGTTGCACTG	28	831
233168	808	211	GGACTGTACCTCAAGCTGGT	0	832
233169	808	231	GCTCATTCTCCAGCATCAGG	14	833
233170	809	251	TTGATCTATAATACTAGCTA	23	834
233172	810	282	ATGGAAGACTGGCAGCTCTA	86	835
233173	808	301	TTGTGTTCTTGAAGCGGCC	3	836
233174	809	301	TGTGCACGGATATGATAACG	21	837
233175	810	306	GACCTTGAGTAGATTCCTGG	90	838
233176	810	321	GAAATCTGGAAGAGAGACCT	62	839
233177	808	331	GTAGCTTTCCCATCTAGGCT	0	840
233178	808	346	GATAACTCTGTGAGGGTAGC	0	841
233179	810	371	ATGTTGCCCATGGCTGGAAT	65	842
233180	809	381	AAGATGCAGTACTACTTCCA	13	843
233181	808	382	GCACCCAGAATCATGGCCT G	0	844
233182	809	411	CTTGATACTTGGTATCCACA	59	845
233183	810	411	CAGTGTAATGATCGTTGATT	88	846
233184	810	431	TAAAGTCCAGCATTGGTATT	69	847
233185	810	451	CAACAATGTCTGATTGGTTA	73	848
233186	810	473	GAAGAGGAAGAAAGGATAT G	60	849
233187	810	481	TGACAGATGAAGAGGAAGAA	66	850
233188	810	500	TTGTACTGTAGTGCATCAAT	74	851
233189	809	511	GCCTCAATCTGTTGTTTCAG	46	852
233190	810	520	ACTTGAGCGTGCCCTCTAAT	69	853
233191	809	561	GAAATGGAATTGTAGTTCTC	31	854

**Example 43**

**Antisense inhibition of rabbit apolipoprotein B expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap-Dose Response Study**

**[0385]** In accordance with the present invention, a subset of the antisense oligonucleotides in Example 42 was further investigated in dose-response studies. Treatment doses were 10, 50, 150 and 300 nM. ISIS 233160 (SEQ ID NO: 824), ISIS 233166 (SEQ ID NO: 830), ISIS 233172 (SEQ ID NO: 835), ISIS 233175 (SEQ ID NO: 838), and ISIS 233183 (SEQ ID NO: 846) were analyzed for their effect on rabbit apolipoprotein B mRNA levels in primary rabbit hepatocytes by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments and are shown in Table 22.

EP 2 336 318 A1

Table 22 Inhibition of rabbit apolipoprotein B mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	Percent Inhibition			
	300 nM	150 nM	50 nM	10 nM
233160	80	74	67	33
233166	73	79	81	66
233172	84	81	76	60
233175	93	90	85	67
233183	80	81	71	30

**Example 44**

**Effects of antisense inhibition of apolipoprotein B in LDLr<sup>-/-</sup> mice - Dose Response**

**[0386]** LDL receptor-deficient mice (LDLr<sup>-/-</sup> mice), a strain that cannot edit the apolipoprotein B mRNA and therefore synthesize exclusively apolipoprotein B-100, have markedly elevated LDL cholesterol and apolipoprotein B-100 levels and develop extensive atherosclerosis.

**[0387]** LDLr<sup>-/-</sup> mice, purchased from Taconic (Germantown, NY) were used to evaluate antisense oligonucleotides for their potential to lower apolipoprotein B mRNA or protein levels, as well as phenotypic endpoints associated with apolipoprotein B. LDLr<sup>-/-</sup> mice were separated into groups of males and females. LDLr<sup>-/-</sup> mice were dosed intraperitoneally twice a week for six weeks with either 10, 25, or 50 mg/kg of ISIS 147764 (SEQ ID NO: 109) or ISIS 270906 (SEQ ID NO: 856) which is a 4 base mismatch of ISIS 147764, or with saline, or 20 mg/kg of Atorvastatin. At study termination animals were sacrificed and evaluated for several phenotypic markers.

**[0388]** ISIS 147764 was able to lower cholesterol, triglycerides, and mRNA levels in a dose-dependent manner in both male and female mice while the 4-base mismatch ISIS 270906 was not able to do this. The results of the study are summarized in Table 23.

Table 23 Effects of ISIS 147764 treatment in male and female LDLr<sup>-/-</sup> mice on apolipoprotein B mRNA, liver enzyme, cholesterol, and triglyceride levels.

ISIS No.	Dose mg/kg	Liver Enzymes IU/L		Lipoproteins mg/dL			TRIG	mRNA % control
		AST	ALT	CHOL	HDL	LDL		
<b>Males</b>								
Saline		68.4	26.6	279.2	125.4	134.7	170.6	100.0
147764	10	57.6	29.8	314.2	150.0	134.7	198.6	61.7
	25	112.6	78.8	185.0	110.6	66.2	104.2	30.7
	50	163.6	156.8	165.6	107.8	51.2	113.4	16.6
270906	50	167.4	348.0	941.0	244.2	541.9	844.8	N.D.
Atorvastatin	20	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	110.9
<b>Females</b>								
Saline		65.0	23.4	265.8	105.8	154.9	121.4	100.0
147764	10	82.0	27.2	269.6	121.0	127.8	140.8	64.2
	25	61.4	32.2	175.8	99.5	68.9	100.4	41.3
	50	134.6	120.4	138.2	92.2	45.9	98.0	18.5
270906	50	96.0	88.6	564.6	200.0	310.0	240.4	N.D.

(continued)

Females								
Atorvastatin	20	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	109.0

**Example 45****Effects of antisense inhibition of apolipoprotein B in Cynomolgus monkeys**

**[0389]** Cynomolgus monkeys fed an atherogenic diet develop atherosclerosis with many similarities to atherosclerosis of human beings. Female Cynomolgus macaques share several similarities in lipoproteins and the cardiovascular system with humans. In addition to these characteristics, there are similarities in reproductive biology. The Cynomolgus female has a 28-day menstrual cycle like that of women. Plasma hormone concentrations have been measured throughout the Cynomolgus menstrual cycle, and the duration of the follicular and luteal phases, as well as plasma estradiol and progesterone concentrations across the cycle, are also remarkably similar to those in women.

**[0390]** Cynomolgus monkeys (male or female) can be used to evaluate antisense oligonucleotides for their potential to lower apolipoprotein B mRNA or protein levels, as well as phenotypic endpoints associated with apolipoprotein B including, but not limited to cardiovascular indicators, atherosclerosis, lipid diseases, obesity, and plaque formation. One study could include normal and induced hypercholesterolemic monkeys fed diets that are normal or high in lipid and cholesterol. Cynomolgus monkeys can be dosed in a variety of regimens, one being subcutaneously with 10-20 mg/kg of the oligomeric compound for 1-2 months. Parameters that may be observed during the test period could include: total plasma cholesterol, LDL-cholesterol, HDL-cholesterol, triglyceride, arterial wall cholesterol content, and coronary intimal thickening.

**Example 46****Sequencing of Cynomolgus monkey (*Macaca fascicularis*) apolipoprotein B preferred target segment**

**[0391]** In accordance with the present invention, a portion of the cynomolgus monkey apolipoprotein B mRNA not available in the art, was amplified. Positions 2920 to 3420 of the human apolipoprotein B mRNA sequence (GenBank accession number NM\_000384.1, incorporated herein as SEQ ID NO: 3) contain the preferred target segment to which ISIS 301012 hybridizes and the corresponding segment of cynomolgus monkey apolipoprotein B mRNA was amplified and sequenced. The site to which ISIS 301012 hybridizes in the human apolipoprotein B was amplified by placing primers at 5' position 2920 and 3' position 3420. The cynomolgus monkey hepatocytes were purchased from In Vitro Technologies (Gaithersburg, MD). The 500 bp fragments were produced using human and cynomolgus monkey 1<sup>o</sup> hepatocyte cDNA and were produced by reverse transcription of purified total RNA followed by 40 rounds of PCR amplification. Following gel purification of the human and cynomolgus amplicons, the forward and reverse sequencing reactions of each product were performed by Retrogen (Invitrogen kit was used to create the single-stranded cDNA and provided reagents for Amplitaq PCR reaction). This cynomolgus monkey sequence is incorporated herein as SEQ ID NO: 855 and is 96% identical to positions 2920 to 3420 of the human apolipoprotein B mRNA.

**Example 47****Effects of antisense inhibition of human apolipoprotein B gene (ISIS 281625 and 301012) in C57BL/6NTac-TgN (APOB100) transgenic mice**

**[0392]** C57BL/6NTac-TgN(APOB100) transgenic mice have the human apolipoprotein B gene "knocked-in". These mice express high levels of human apolipoprotein B100 resulting in mice with elevated serum levels of LDL cholesterol. These mice are useful in identifying and evaluating compounds to reduce elevated levels of LDL cholesterol and the risk of atherosclerosis. When fed a high fat cholesterol diet, these mice develop significant foam cell accumulation underlying the endothelium and within the media, and have significantly more complex atherosclerotic lesions than control animals.

**[0393]** C57BL/6NTac-TgN(APOB100) mice were divided into two groups - one group receiving oligonucleotide treatment and control animals receiving saline treatment. After overnight fasting, mice were dosed intraperitoneally twice a week with saline or 25 mg/kg ISIS 281625 (SEQ ID No: 224) or ISIS 301012 (SEQ ID No: 247) for eight weeks. At study termination and forty eight hours after the final injections, animals were sacrificed and evaluated for target mRNA levels

in liver, cholesterol and triglyceride levels, and liver enzyme levels. In addition, the endogenous mouse apolipoprotein B levels in liver were measured to evaluate any effects of these antisense oligonucleotides targeted to the human apolipoprotein B.

[0394] Upon treatment with either ISIS 281625 or ISIS 301012, the AST and ALT levels were increased, yet did not exceed normal levels (~300 IU/L). Cholesterol levels were slightly increased relative to saline treatment, while triglyceride levels were slightly decreased. Treatment with either of these oligonucleotides targeted to the human apolipoprotein B which is expressed in these mice markedly decreased the mRNA levels of the human apolipoprotein, while the levels of the endogenous mouse apolipoprotein B were unaffected, indicating that these oligonucleotides exhibit specificity for the human apolipoprotein B. The results of the comparative studies are shown in Table 24.

Table 24 Effects of ISIS 281625 and 301012 treatment in mice on apolipoprotein B mRNA, liver enzyme, cholesterol, and triglyceride levels.

		ISIS No.		
		SALINE	281625	301012
Liver Enzymes IU/L				
	AST	70.3	265.8	208.4
	ALT	32.8	363.8	137.4
Lipoproteins mg/dL				
	CHOL	109.5	152.0	145.1
	HDL	67.3	84.6	98.6
	LDL	30.2	49.8	36.6
	TRIG	194.5	171.1	157.8
mRNA % control				
	human mRNA	100.0	45.2	23.7
	mouse mRNA	100.0	111.0	94.6

[0395] Following 2 and 4 weeks of ISIS 301012 treatment, LDL-cholesterol levels were significantly reduced to 22 mg/dL and 17 mg/dL, respectively.

[0396] Apolipoprotein B protein levels in liver were also evaluated at the end of the 8 week treatment period. Liver protein was isolated and subjected to immunoblot analysis using antibodies specific for human or mouse apolipoprotein B protein (US Biologicals, Swampscott, MA and Santa Cruz Biotechnology, Inc., Santa Cruz, CA, respectively). Immunoblot analysis of liver protein samples reveals a reduction in the expression of both forms of human apolipoprotein B , apolipoprotein B-100 and apolipoprotein B-48. Mouse apolipoprotein B levels in liver were not significantly changed, as judged by immunoblot analysis.

[0397] Serum samples were also collected at 2, 4, 6 and 8 weeks and were evaluated for human apolipoprotein B expression by using a human apolipoprotein B specific ELISA kit (ALerCHEK Inc., Portland, ME). Quantitation of serum human apolipoprotein B protein by ELISA revealed that treatment with ISIS 281625 reduced serum human apolipoprotein B protein by 31, 26, 11 and 26% at 2, 4, 6 and 8 weeks, respectively, relative to saline-treated animals. Treatment with ISIS 301012 reduced serum human apolipoprotein B protein by 70, 87, 81 and 41% at 2, 4, 6 and 8 weeks, respectively, relative to saline-treated control animals. Serum from transgenic mice was also subjected to immunoblot analysis using both human and mouse specific apolipoprotein B antibodies (US Biologicals, Swampscott, MA and Santa Cruz Biotechnology, Inc., Santa Cruz, CA, respectively). Immunoblot analysis of serum samples taken from animals shows a similar pattern of human apolipoprotein B expression, with a significant reduction in serum apolipoprotein B protein after 2, 4 and 6 weeks of treatment and a slight reduction at 8 weeks. Mouse apolipoprotein B in serum was not significantly changed, as judged by immunoblot analysis.

**Example 48**

**Effects of ant3.sense inhibition of apolipoprotein B (ISIS 233172, 233175, 281625, 301012, and 301027) in C57BL/6 mice**

**[0398]** C57BL/6 mice, a strain reported to be susceptible to hyperlipidemia-induced atherosclerotic plaque formation were used in the following studies to evaluate the toxicity in mice of several antisense oligonucleotides targeted to human or rabbit apolipoprotein B.

**[0399]** C57BL/6 mice were divided into two groups - one group receiving oligonucleotide treatment and control animals receiving saline treatment. After overnight fasting, mice were dosed intraperitoneally twice a week with saline or 25 mg/kg of one of several oligonucleotides for two weeks. The antisense oligonucleotides used in the present study were ISIS 233172 (SEQ ID NO: 835) and ISIS 233175 (SEQ ID NO: 838), both targeted to rabbit apolipoprotein B, and ISIS 281625 (SEQ ID NO: 224), ISIS 301012 (SEQ ID NO: 247), and ISIS 301027 (SEQ ID NO: 262), targeted to human apolipoprotein B. At study termination and forty eight hours after the final injections, animals were sacrificed and evaluated for liver enzyme levels, body weight, liver weight, and spleen weight.

**[0400]** The levels of liver enzymes in mice were decreased relative to saline treatment for three of the antisense oligonucleotide. However, the rabbit oligonucleotide ISIS 233175 and the human oligonucleotide ISIS 301027 both elicited drastically increased levels of these liver enzymes, indicating toxicity. For all of the oligonucleotides tested, the change in weight of body, liver, and spleen were minor. The results of the comparative studies are shown in Table 25.

Table 25 Effects of antisense oligonucleotides targeted to human or rabbit apolipoprotein B on mouse apolipoprotein B mRNA, liver enzyme, cholesterol, and triglyceride levels.

		SALINE	233172	233175	ISIS No.		
					281625	301012	301027
Liver Enzymes							
AST	IU/L	104.5	94.3	346.7	89.5	50.6	455.3
ALT	IU/L	39.5	43.3	230.2	36.2	21.2	221.3
Weight							
BODY		21.2	21.3	21.5	20.9	21.3	21.2
LIVER		1.1	1.3	1.4	1.2	1.1	1.3
SPLEEN		0.1	0.1	0.1	0.1	0.1	0.1

**Example 49**

**Time course evaluation of oligonucleotide at two different doses**

**[0401]** C57BL/6 mice, a strain reported to be susceptible to hyperlipidemia-induced atherosclerotic plaque formation were used in the following studies to evaluate the toxicity in mice of several antisense oligonucleotides targeted to human apolipoprotein B.

**[0402]** Female C57BL/6 mice were divided into two groups - one group receiving oligonucleotide treatment and control animals receiving saline treatment. After overnight fasting, mice were dosed intraperitoneally twice a week with saline or 25 mg/kg or 50 mg/kg of ISIS 281625 (SEQ ID NO: 224), ISIS 301012 (SEQ ID NO: 247), or ISIS 301027 (SEQ ID NO: 262). After 2 weeks, a blood sample was taken from the tail of the mice and evaluated for liver enzyme. After 4 weeks, and study termination, animals were sacrificed and evaluated for liver enzyme levels.

**[0403]** For ISIS 281625 and ISIS 301012, AST and ALT levels remained close to those of saline at either dose after 2 weeks. After 4 weeks, AST and ALT levels showed a moderate increase over saline treated animals for the lower dose, but a large increase at the higher dose. ISIS 301027, administered at either dose, showed a small increase in AST and ALT levels after 2 weeks and a huge increase in AST and ALT levels after 4 weeks. The results of the studies are summarized in Table 26.

EP 2 336 318 A1

Table 26 AST and ALT levels in mice treated with ISIS 281625, 301012, or 301027 after 2 and 4 weeks

		AST (IU/L)		ALT (IU/L)	
		2 weeks	4 weeks	2 weeks	4 weeks
SALINE		49.6	63.2	22.4	25.2
ISIS No.	Dose (mg/kg)				
281625	25	40.8	75	21.2	31.8
	50	44.4	152.4	30.8	210.4
301012	25	37.2	89.8	22.4	24.8
	50	38.4	107.4	23.2	29.2
301027	25	55.4	537.6	27.2	311.2
	50	64	1884	34.8	1194

Example 50

Effects of antisense inhibition of apolipoprotein B (ISIS 147483 and 147764) in ob/ob mice

[0404] Leptin is a hormone produced by fat that regulates appetite. Deficiencies in this hormone in both humans and non-human animals leads to obesity. ob/ob mice have a mutation in the leptin gene which results in obesity and hyperglycemia. As such, these mice are a useful model for the investigation of obesity and diabetes and treatments designed to treat these conditions.

[0405] Ob/ob mice receiving a high fat, high cholesterol diet (60% kcal fat supplemented with 0.15% cholesterol) were treated with one of several oligonucleotides to evaluate their effect on apolipoprotein B-related phenotypic endpoints in ob/ob mice. After overnight fasting, mice from each group were dosed intraperitoneally twice a week with 50 mg/kg of ISIS 147483 (SEQ ID NO: 79), or 147764 (SEQ ID NO: 109), or the controls ISIS 116847 (SEQ ID NO: 857), or 141923 (SEQ ID NO: 858), or saline for six weeks. At study termination and forty eight hours after the final injections, animals were sacrificed and evaluated for target mRNA levels in liver, cholesterol and triglyceride levels, liver enzyme levels, serum glucose levels, and PTEN levels.

[0406] ISIS 147483 and 147764 were both able to lower apolipoprotein B mRNA levels, as well as glucose, cholesterol, and triglyceride levels. The results of the comparative studies are shown in Table 27.

Table 27 Effects of ISIS 147483 and 147764 treatment in ob/ob mice on apolipoprotein B mRNA, cholesterol, lipid, triglyceride, liver enzyme, glucose, and PTEN levels.

		ISIS No.				
		SALINE	116847	141923	147483	147764
Glucose mg/dL		269.6	135.5	328.5	213.2	209.2
Liver Enzymes						
IU/L	AST	422.3	343.2	329.3	790.2	406.5
	ALT	884.3	607.5	701.7	941.7	835.0
Lipoproteins						
mg/dL	CHOL	431.9	287.5	646.3	250.0	286.3
	TRIG	128.6	196.5	196.5	99.8	101.2
mRNA						
% control						
ApoB		100.0	77.0	100.0	25.2	43.1
PTEN		100.0	20.0	113.6	143.2	115.3

## Example 51

## Antisense inhibition of apolipoprotein B in high fat fed mice: time-dependent effects

**[0407]** In a further embodiment of the invention, the inhibition of apolipoprotein B mRNA in mice was compared to liver oligonucleotide concentration, total cholesterol, LDL-cholesterol and HDL-cholesterol. Male C57B1/6 mice receiving a high fat diet (60% fat) were evaluated over the course of 6 weeks for the effects of treatment with twice weekly intraperitoneal injections of 50 mg/kg ISIS 147764 (SEQ ID NO: 109) or 50 mg/kg of the control oligonucleotide ISIS 141923 (SEQ ID NO: 858). Control animals received saline treatment. Animals were sacrificed after 2 days, 1, 2, 4 and 6 weeks of treatment. Each treatment group at each time point consisted of 8 mice.

**[0408]** Target expression in liver was measured by real-time PCR as described by other examples herein and is expressed as percent inhibition relative to saline treated mice. Total, LDL- and HDL-cholesterol levels were measured by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY) and are presented in mg/dL. Results from saline-treated animals are shown for comparison. Intact oligonucleotide in liver tissue was measured by capillary gel electrophoresis and is presented as micrograms of oligonucleotide per gram of tissue. All results are the average of 8 animals and are shown in Table 28.

Table 28 Correlation between liver drug concentration, apolipoprotein B mRNA expression and serum lipids during ISIS 147764 treatment

	ISIS # days	Treatment period				
		2 days	1 week	2 weeks	4 weeks	6 weeks
% Inhibition apolipoprotein B mRNA	141923	9	4	7	0	0
	147764	50	57	73	82	88
Intact oligonucleotide ug/g	141923	58	61	152	261	631
	147764	85	121	194	340	586
Total cholesterol mg/dL	saline	105	152	144	180	191
	141923	99	146	152	169	225
	147764	101	128	121	75	73
LDL-cholesterol mg/dL	saline	8	32	28	50	46
HDL-cholesterol mg/dL	141923	8	27	27	38	56
	147764	7	19	14	7	7
HDL-cholesterol mg/dL	saline	74	117	114	127	141
	141923	70	116	122	128	166
	147764	76	107	105	66	64

**[0409]** These results illustrate that inhibition of apolipoprotein B mRNA by ISIS 147764 occurred within 2 days of treatment, increased with successive treatments and persisted for 6 weeks of treatment. Quantitation of liver oligonucleotide levels reveals a strong correlation between the extent of target inhibition and liver drug concentration. Furthermore, at 1, 2, 3 and 4 weeks of treatment, a inverse correlation between inhibition of target mRNA and cholesterol levels (total, HDL and LDL) is observed, with cholesterol levels lowering as percent inhibition of apolipoprotein B mRNA becomes greater. Serum samples were subjected to immunoblot analysis using an antibody to detect mouse apolipoprotein B protein (Gladstone Institute, San Francisco, CA). The expression of protein follows the same pattern as that of the mRNA, with apolipoprotein B protein in serum markedly reduced within 48 hours and lowered throughout the 6 week treatment period.

**[0410]** The oligonucleotide treatments described in this example were duplicated to investigate the extent to which

**EP 2 336 318 A1**

effects of ISIS 147764 persist following cessation of treatment. Mice were treated as described, and sacrificed 1, 2, 4, 6 and 8 weeks following the cessation of oligonucleotide treatment. The same parameters were analyzed and the results are shown in Table 29.

5 Table 29 Correlation between liver drug concentration, apolipoprotein B mRNA expression, and serum lipids after cessation of dosing

	ISIS #	Treatment period				
		1 week	2 weeks	4 weeks	6 weeks	8 weeks
10 % Inhibition apolipoprotein B mRNA	141923	15	2	7	11	7
	147764	82	78	49	37	19
15 Intact oligonucleotide ug/g	141923	297	250	207	212	128
	147764	215	168	124	70	43
20 Total cholesterol mg/dL	saline	114	144	195	221	160
	141923	158	139	185	186	151
	147764	69	67	111	138	135
25 LDL-cholesterol mg/dL	saline	21	24	34	37	22
	141923	24	24	32	32	24
	147764	14	14	18	24	21
30 HDL-cholesterol mg/dL	saline	86	109	134	158	117
	141923	121	105	135	136	108
	147764	51	49	79	100	94

35 [0411] These data demonstrate that after termination of oligonucleotide treatment, the effects of ISIS 147764, including apolipoprotein B mRNA inhibition, and cholesterol lowering, persist for up to 8 weeks. Immunoblot analysis demonstrates that apolipoprotein B protein levels follow a pattern similar that observed for mRNA expression levels.

**Example 52**

40 **Effects of antisense inhibition of human apolipoprotein B gene by 301012 in C57BL/6NTac-TgN(APOB100) transgenic mice: dosing study**

45 [0412] C57BL/6NTac-TgN(APOB100) transgenic mice have the human apolipoprotein B gene "knocked-in". These mice express high levels of human apolipoprotein B resulting in mice with elevated serum levels of LDL cholesterol. These mice are useful in identifying and evaluating compounds to reduce elevated levels of LDL cholesterol and the risk of atherosclerosis. When fed a high fat cholesterol diet, these mice develop significant foam cell accumulation underlying the endothelium and within the media, and have significantly more complex atherosclerotic plaque lesions than control animals.

50 [0413] A long-term study of inhibition of human apolipoprotein B by ISIS 301012 in C57BL/6NTac-TgN(APOB100) mice (Taconic, Germantown, NY) was conducted for a 3 month period. Mice were dosed intraperitoneally twice a week with 10 or 25 mg/kg ISIS 301012 (SEQ ID No: 247) for 12 weeks. Saline-injected animals served as controls. Each treatment group comprised 4 animals.

55 [0414] After 2, 4, 6, 8 and 12 weeks of treatment, serum samples were collected for the purpose of measuring human apolipoprotein B protein. Serum protein was quantitated using an ELISA kit specific for human apolipoprotein B (ALerCHEK Inc., Portland, ME). The data are shown in Table 30 and each result represents the average of 4 animals. Data are normalized to saline-treated control animals.



EP 2 336 318 A1

Table 30 Reduction of human apolipoprotein B protein in transgenic mouse serum following ISIS 301012 treatment

Dose of oligonucleotide mg/kg	% Reduction in human apolipoprotein B protein in serum				
	2 weeks	4 weeks	6 weeks	8 weeks	12 weeks
10	76	78	73	42	85
25	80	87	86	47	79

[0415] These data illustrate that following 2, 4, 6 or 12 weeks of treatment with ISIS 301012, the level of human apolipoprotein B protein in serum from transgenic mice is lowered by approximately 80%, demonstrating that in addition to inhibiting mRNA expression, ISIS 301012 effectively inhibits human apolipoprotein B protein expression in mice carrying the human apolipoprotein B transgene. Apolipoprotein B protein in serum was also assessed by immunoblot analysis using an antibody directed to human apolipoprotein B protein (US Biologicals, Swampscott, MA). This analysis shows that the levels human apolipoprotein B protein, both the apolipoprotein B-100 and apolipoprotein B-48 forms, are lowered at 2, 4, 6 and 12 weeks of treatment. Immunoblot analysis using a mouse apolipoprotein B specific antibody (Santa Cruz Biotechnology, Inc., Santa Cruz, CA) reveals no significant change in the expression of the mouse protein in serum.

[0416] At the beginning of the treatment (start) and after 2, 4, 6 and 8 weeks of treatment, serum samples were collected and total, LDL- and HDL-cholesterol levels were measured by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY), and these data are presented in Table 31. Results are presented as mg/dL in serum and represent the average of 4 animals. Results from the saline control animals are also shown.

Table 31 Effects of ISIS 301012 on serum lipids in human apolipoprotein B transgenic mice

	Treatment	Treatment period				
		Start	2 weeks	4 weeks	6 weeks	8 weeks
Total cholesterol mg/dL	Saline	120	110	129	121	126
	10	115	97	111	120	122
	25	107	101	107	124	147
HDL-cholesterol mg/dL	Saline	67	61	69	62	64
	10	70	69	78	72	79
	25	64	73	76	80	91
LDL-cholesterol mg/dL	Saline	39	41	50	45	47
	10	35	20	23	37	33
	25	33	19	19	37	44

[0417] These data demonstrate that LDL-cholesterol is lowered by treatment with 10 or 25 mg/kg of ISIS 147764 during the first 4 weeks of treatment.

[0418] The study was terminated forty eight hours after the final injections in the eighth week of treatment, when animals were sacrificed and evaluated for target mRNA levels in liver, apolipoprotein B protein levels in liver and serum cholesterol and liver enzyme levels. In addition, the expression of endogenous mouse apolipoprotein B levels in liver was measured to evaluate any effects of ISIS 301012 on mouse apolipoprotein B mRNA expression.

[0419] Human and mouse apolipoprotein B mRNA levels in livers of animals treated for 12 weeks were measured by real-time PCR as described herein. Each result represents the average of data from 4 animals. The data were normalized to saline controls and are shown in Table 32.

Table 32 Effects of ISIS 301012 on human and mouse apolipoprotein B mRNA levels in transgenic mice

mRNA species measured	% Inhibition	
	Dose of ISIS 301012	
	10 mg/kg	25 mg/kg

(continued)

	% Inhibition	
	Dose of ISIS 301012	
human apolipoprotein B	65	75
mouse apolipoprotein B	6	6

[0420] These data demonstrate that following 12 weeks of treatment with ISIS 301012, human apolipoprotein B mRNA is reduced by as much as 75% in the livers of transgenic mice, whereas mouse liver apolipoprotein B mRNA was unaffected. Furthermore, ELISA analysis of apolipoprotein B protein in livers of transgenic mice reveals an 80% and 82% reduction in the human protein following 10 and 20 mg/kg ISIS 301012, respectively. Immunoblot analysis using an antibody directed to human apolipoprotein B also demonstrates a reduction in the expression of human apolipoprotein B, both the apolipoprotein B-100 and apolipoprotein B-48 forms, in the livers of transgenic mice. Immunoblot analysis using an antibody directed to mouse apolipoprotein B protein (Santa Cruz Biotechnology, Inc., Santa Cruz, CA) reveals that expression of the mouse protein in liver does not change significantly.

[0421] ALT and AST levels in serum were also measured using the Olympus Clinical Analyzer (Olympus America Inc., Melville, NY) and showed that following treatment with ISIS 301012, the AST and ALT levels were increased, yet did not exceed normal levels (~300 IU/L), indicating a lack of toxicity due to ISIS 301012 treatment.

### Example 53

#### Assessment of in vitro immunostimulatory effects of ISIS 301012

[0422] Immunostimulatory activity is defined by the production of cytokines upon exposure to a proinflammatory agent. In a further embodiment of the invention, ISIS 301012 was tested for immunostimulatory, or proinflammatory, activity. These studies were performed by MDS Pharma Services (Saint Germain sur l'Arbresle, France). Whole blood was collected from naive B6C3F1 mice, which had not been knowingly exposed to viral, chemical or radiation treatment. Cultured blood cells were exposed to 0.5, 5 or 50  $\mu$ M of ISIS 301012 for a period of 14 to 16 hours. Antisense oligonucleotides known to possess proinflammatory activity served as positive controls. Each treatment was performed in triplicate. At the end of the treatment period, supernatants were collected and cytokine analysis was performed using a flow cytometry method with the mouse Inflammation CBA kit (Becton Dickinson, Franklin Lakes, NJ). The results revealed that ISIS 301012 does not stimulate the release of any of the tested cytokines, which were interleukin-12p70 (IL-12p70), tumor necrosis factor-alpha (TNF-alpha), interferon-gamma (IFN-gamma), interleukin-6 (IL-6), macrophage chemoattractant protein-1 (MCP-1) and interleukin-10 (IL-10). Thus, ISIS 301012 does not possess immunostimulatory activity, as determined by the in vitro immunostimulatory assay.

### Example 54

#### Comparative genomic analysis of apolipoprotein B

[0423] In accordance with the present invention, a comparative genomic analysis of apolipoprotein B sequences from human, mouse and monkey was performed and illustrated that apolipoprotein B sequences are conserved across species. The organization of human and mouse apolipoprotein B genes is also highly conserved. The human and mouse genes are comprised of 29 and 26 exons, respectively. The mouse mRNA is approximately 81% homologous to the human sequence. The complete sequence and gene structure of the apolipoprotein B gene in non-human primates have not been identified. However, as illustrated in Example 46, a 500 base pair fragment which contains the ISIS 301012 target sequence exhibits approximately 96% identity to the human sequence.

[0424] The binding site for ISIS 301012 lies within the coding region, within exon 22 of the human apolipoprotein B mRNA. When the ISIS 301012 binding sites from human, mouse and monkey were compared, significant sequence diversity was observed. Although the overall sequence conservation between human and monkey over a 500 nucleotide region was approximately 96%, the ISIS 301012 binding site of the monkey sequence contains 2 mismatches relative to the human sequence. Likewise, though the mouse apolipoprotein B mRNA sequence is approximately 81% homologous to human, within the ISIS 301012 binding site, 5 nucleotides are divergent. The sequence comparisons for the ISIS 301012 binding site for human, mouse and monkey apolipoprotein B sequences are shown in Table 33. Mismatched nucleotides relative to the ISIS 301012 target sequence are underlined.

Table 33 Comparison of ISIS 301012 binding site among human, monkey and mouse apolipoprotein B sequences

Species	# Mismatches	ISIS 301012 target sequence
Human	0	aggtgcaagcagactgagg
Monkey	2	aggtgtaagcagactgagg
Mouse	5	aggagtcagcagctgaag

[0425] The target sequence to which the mouse antisense oligonucleotide ISIS 147764 hybridizes lies within exon 24 of the mouse apolipoprotein B gene. The sequence comparisons for the ISIS 147764 binding site in mouse and human apolipoprotein B sequences are shown in Table 34. Mismatched nucleotides relative to the ISIS 147764 target sequence are underlined.

Table 34 Comparison of ISIS 147764 binding site between mouse and human apolipoprotein B sequences

Species	# Mismatches	ISIS 147764 binding site
Human	5	gcattgacatcttcagggac
Mouse	0	gcatggacttcttgaaa

### Example 55

#### BLAST analysis of ISIS 301012

[0426] In accordance with the present invention, the number of regions in the human genome to which ISIS 301012 will hybridize with perfect complementarity was determined. Percent complementarity of an antisense compound with a region of a target nucleic acid was determined using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., J. Mol. Biol., 1990, 215, 403-410; Zhang and Madden, Genome Res., 1997, 7, 649-656). This analysis assessed sequence complementarity in genomic or pre-mRNA regions and in coding sequences.

[0427] In genomic regions, ISIS 301012 shows perfect sequence complementarity to the apolipoprotein B gene only. No target sequences with one mismatch relative to ISIS 301012 were found. Two mismatches are found between the ISIS 301012 target sequence and the heparanase gene, and 3 mismatches are found between the ISIS 301012 target sequence and 28 unique genomic sites.

[0428] In RNA sequences, perfect sequence complementarity is found between ISIS 301012 and the apolipoprotein B mRNA and three expressed sequence tags that bear moderate similarity to a human apolipoprotein B precursor. A single mismatch is found between ISIS 301012 and an expressed sequence tag similar to the smooth muscle form of myosin light chain.

### Example 56

#### Antisense inhibition of apolipoprotein B in primary human hepatocytes: dose response studies

[0429] In accordance with the present invention, antisense oligonucleotides targeted to human apolipoprotein B were tested in dose response studies in primary human hepatocytes. Pre-plated primary human hepatocytes were purchased from Invitro Technologies (Baltimore, MD). Cells were cultured in high-glucose DMEM (Invitrogen Corporation, Carlsbad, CA) supplemented with 10% fetal bovine serum (Invitrogen Corporation, Carlsbad, CA), 100 units/mL and 100 µg/mL streptomycin (Invitrogen Corporation, Carlsbad, CA).

[0430] Human primary hepatocytes were treated with ISIS 301012 (SEQ ID NO: 247) at 10, 50, 150 or 300 nM. Untreated cells and cells treated with the scrambled control oligonucleotide ISIS 113529 (CTCTTACTGTGCTGTGGACA, SEQ ID NO: 859) served as two groups of control cells. ISIS 113529 is a chimeric oligonucleotide ("gapmer") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidines are 5-methylcytidines.

[0431] Oligonucleotides were introduced into cells through LIPOFECTIN-mediated transfection as described by other examples herein. Cells were harvested both 24 and 48 hours after treatment with oligonucleotide, and both RNA and

**EP 2 336 318 A1**

protein were isolated. Additionally, the culture media from treated cells was collected for ELISA analysis of apolipoprotein B protein secretion.

[0432] Apolipoprotein B mRNA expression was determined by real-time PCR of RNA samples as described by other examples herein. Each result represents 6 experiments. The data are normalized to untreated control cells and are shown in Table 35.

Table 35 Inhibition of apolipoprotein B mRNA by antisense oligonucleotides in human primary hepatocytes

		% Inhibition of apolipoprotein B mRNA	
		ISIS #	
Dose of oligonucleotide	Treatment (hours)	301012	113529
10 nM	24	65	N.D.
	48	33	N.D.
50 nM	24	75	N.D.
	48	48	N.D.
150 nM	24	90	16
	48	78	5
300 nM	24	89	10
	48	72	18

[0433] These data demonstrate that ISIS 301012 inhibits apolipoprotein B expression in a dose-dependent manner in human primary hepatocytes.

[0434] Apolipoprotein B protein secreted from into the cultured cell media was measured in the samples treated with 50 and 150 nM of oligonucleotide, using a target protein specific ELISA kit (ALerCHEK Inc., Portland, ME). Each result represents 3 experiments. The data are normalized to untreated control cells and are shown in Table 36.

Table 36 Inhibition of apolipoprotein B protein secretion from human primary hepatocytes by ISIS 301012

		% Change in apolipoprotein B protein secretion	
		ISIS #	
Dose	Treatment (hours)	301012	113529
150 nM	24	-57	+6
	48	-75	+4
300 nM	24	-41	-2
	48	-48	-5

[0435] Protein samples from 50, 150 and 300 nM doses after 24 hours and 150 and 300 nM doses after 48 hours were subjected to immunoblot analysis as described by other examples herein, using a human apolipoprotein B protein specific antibody purchased from US Biological (Swampscott, MA). Immunoblot analysis further demonstrates that apolipoprotein B protein in human hepatocytes is reduced in a dose-dependent manner following antisense oligonucleotide treatment with ISIS 301012.

[0436] An additional experiment was performed to test the effects of ISIS 271009 (SEQ ID NO: 319), ISIS 281625 (SEQ ID NO: 224) and ISIS 301027 (SEQ ID NO: 262) on human apolipoprotein B mRNA in human primary hepatocytes. Cells were cultured as described herein and treated with 5, 10, 50 or 150 nM of ISIS 271009, ISIS 281625 or ISIS 301027 for a period of 24 hours. The control oligonucleotides ISIS 13650 (SEQ ID NO: 806) and ISIS 113529 (SEQ ID NO: 859) were used at 50 or 150 nM. Human apolipoprotein B mRNA expression was evaluated by real-time PCR as described by other examples herein. Apolipoprotein B protein secreted into the cultured cell media was measured in the samples treated with 50 and 150 nM of oligonucleotide, using a target protein specific ELISA kit (ALerCHEK Inc., Portland, ME).

[0437] The data, shown in Table 37, represent the average 2 experiments and are normalized to untreated control cells. Where present, a "+" indicates that gene expression was increased.

Table 37 Antisense inhibition of human apolipoprotein B mRNA by ISIS 271009, ISIS 281625 and ISIS 301027

	Oligonucleotide dose	ISIS 271009	ISIS 281625	ISIS 301027	ISIS 13650	ISIS 113529
% Inhibition of apolipoprotein B mRNA expression	5 nM	+4	8	11	N.D.	N.D.
	10 nM	5	22	37	N.D.	N.D.
	50 nM	52	49	50	38	0
	150 nM	81	52	70	26	14
% Inhibition of apolipoprotein B protein secretion	50 nM	17	18	21	N.D.	N.D.
	150 nM	32	18	32	+18	+1

[0438] These data demonstrate that ISIS 271009, ISIS 281625 and ISIS 301027 inhibit apolipoprotein B mRNA expression in a dose-dependent manner in human primary hepatocytes. ISIS 271009 and ISIS 301027 inhibit the secretion of apolipoprotein B protein from cells in a dose-dependent manner.

#### Example 57

#### Effects of apolipoproteinB-100 antisense oligonucleotides on apolipoprotein(a) expression

[0439] Lipoprotein(a) [Lp(a)] contains two disulfide-linked distinct proteins, apolipoprotein(a) and apolipoprotein B (Rainwater and Kammerer, J. Exp. Zool., 1998, 282, 54-61). In accordance with the present invention, antisense oligonucleotides targeted to apolipoprotein B were tested for effects on the expression of the apolipoprotein(a) component of the lipoprotein(a) particle in primary human hepatocytes.

[0440] Primary human hepatocytes (Invitro Technologies, Baltimore, MD), cultured and transfected as described herein, were treated with 5, 10, 50 or 150 nM of ISIS 271009 (SEQ ID NO: 319), 281625 (SEQ ID NO: 224), 301012 (SEQ ID NO: 247) or 301027 (SEQ ID NO: 262). Cells were also treated with 50 or 150 nM of the control oligonucleotides ISIS 113529 (SEQ ID NO: 859) or ISIS 13650 (SEQ ID NO: 806). Untreated cells served as a control. Following 24 hours of oligonucleotide treatment, apolipoprotein(a) mRNA expression was measured by quantitative real-time PCR as described in other examples herein.

[0441] Probes and primers to human apolipoprotein(a) were designed to hybridize to a human apolipoprotein(a) sequence, using published sequence information (GenBank accession number NM-005577.1, incorporated herein as SEQ ID NO: 860). For human apolipoprotein(a) the PCR primers were:

forward primer: CAGCTCCTTATTGTTATACGAGGGA (SEQ ID NO: 861)

reverse primer: TGCCTCTGAGCATTGCGT (SEQ ID NO: 862) and the PCR probe was: FAM-CCCGGTGTCAG-GTGGGAGTACTGC-TAMRA (SEQ ID NO: 863) where FAM is the fluorescent dye and TAMRA is the quencher dye.

[0442] Data are the average of three experiments and are expressed as percent inhibition relative to untreated controls. The results are shown in Table 38. A "+" or "-" preceding the number indicates that apolipoprotein(a) expression was increased or decreased, respectively, following treatment with antisense oligonucleotides.

Table 38 Effects of apolipoprotein B antisense oligonucleotides on apolipoprotein(a) expression

Oligonucleotide Dose	% Change in apolipoprotein(a) mRNA expression following antisense inhibition of apolipoprotein B					
	ISIS #					
	271009	281625	301012	301027	13650	113529
5 nM	+70	-9	+34	-16	N.D.	N.D.
10 nM	+31	-23	+86	-45	N.D.	N.D.

(continued)

	% Change in apolipoprotein(a) mRNA expression following antisense inhibition of apolipoprotein B					
Oligonucleotide Dose	ISIS #					
	271009	281625	301012	301027	13650	113529
50 nM	+25	-34	+30	-39	-68	+14
150 nM	-47	+32	+38	-43	-37	-9

[0443] These results illustrate that ISIS 301012 did not inhibit the expression of apolipoprotein(a) in human primary hepatocytes. ISIS 271009 inhibited apolipoprotein(a) expression at the highest dose. ISIS 281625 and ISIS 301027 decreased the levels of apolipoprotein(a) mRNA.

#### Example 58

#### Inhibition of lipoprotein(a) particle secretion with antisense oligonucleotides targeted to apolipoprotein B-100

[0444] In accordance with the present invention, the secretion of lipoprotein(a) particles, which are comprised of one apolipoprotein(a) molecule covalently linked to one apolipoprotein B molecule, was evaluated in primary human hepatocytes treated with antisense oligonucleotides targeted to the apolipoprotein B component of lipoprotein(a).

[0445] Primary human hepatocytes (InVitro Technologies, Baltimore, MD), cultured and transfected as described herein, were treated for 24 hours with 50 or 150 nM of ISIS 271009 (SEQ ID NO: 319), 281625 (SEQ ID NO: 224), 301012 (SEQ ID NO: 247) or 301027 (SEQ ID NO: 262). Cells were also treated with 150 nM of the control oligonucleotides ISIS 113529 (SEQ ID NO: 859) or ISIS 13650 (SEQ ID NO: 806). Untreated cells served as a control. Following 24 hours of oligonucleotide treatment, the amount of lipoprotein(a) in the culture medium collected from the treated cells was measured using a commercially available ELISA kit (ALerCHEK Inc., Portland, ME). The results are the average of three experiments and are expressed as percent change in lipoprotein(a) secretion relative to untreated controls. The data are shown in Table 39. A "+" or "-" preceding the number indicates that lipoprotein(a) particle secretion was increased or decreased, respectively, following treatment with antisense oligonucleotides targeted to apolipoprotein B.

Table 39 Inhibition of lipoprotein(a) particle secretion with antisense oligonucleotides targeted to apolipoprotein B

	% Change in lipoprotein(a) secretion					
Oligonucleotide Dose	ISIS #					
	271009	281625	301012	301027	13650	113529
50 nM	-25	-26	-27	-33	N.D.	N.D.
150 nM	-42	-24	-37	-44	+14	+14

[0446] These data demonstrate that antisense inhibition of apolipoprotein B, a component of the lipoprotein(a) particle, can reduce the secretion of lipoprotein(a) from human primary hepatocytes. In addition, this reduction in lipoprotein(a) secretion is not necessarily concomitant with a decrease in apolipoprotein(a) mRNA expression, as shown in Example 57.

#### Example 59

#### Mismatched and truncated derivatives of ISIS 301012

[0447] As demonstrated herein, ISIS 301012 (SEQ ID NO: 247) reduces apolipoprotein B mRNA levels in cultured human cell lines as well as in human primary hepatocytes. In a further embodiment of the invention, a study was performed using nucleotide sequence derivatives of ISIS 301012. A series of oligonucleotides containing from 1 to 7 base mismatches, starting in the center of the ISIS 301012 sequence, was designed. This series was designed to introduce the consecutive loss of Watson-Crick base pairing between ISIS 301012 and its target mRNA sequence. These compounds are shown in Table 40. The antisense compounds with mismatched nucleotides relative to ISIS 301012 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxy-nucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed

## EP 2 336 318 A1

of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide.

[0448] An additional derivative of ISIS 301012 was designed, comprising the ISIS 301012 sequence with 2'MOE nucleotides throughout the oligonucleotide (uniform 2'-MOE). This compound is 20 nucleotides in length, with phosphorothioate linkages throughout the oligonucleotide. This compound is also shown in Table 40.

[0449] HepG2 cells were treated with 50 or 150 nM of the compounds in Table 40 for a 24 hour period, after which RNA was isolated and target expression was measured by real-time PCR as described herein. Untreated cells served as controls. The results are shown in Tables 40 and are normalized to untreated control samples.

Table 40 Effects of ISIS 301012 mismatched oligonucleotides and a uniform 2'MOE oligonucleotide on apolipoprotein B expression in HepG2 cells

ISIS #	SEQUENCE	# Mismatches	% Change in apolipoprotein B mRNA expression		SEQ ID NO
			Dose of oligonucleotide		
			50	150	
301012	GCCTCAGTCTGCTTCGCAC C	0	-44	-75	247
<b>Mismatch Series, chimeric oligonucleotides</b>					
332770	GCCTCAGTCTTC <u>T</u> TCGCACC	1	+7	-22	864
332771	GCCTCAGTCTT <u>A</u> TCGCACC	2	+37	+37	865
332772	GCCTCAGT <u>A</u> T <u>A</u> TCGCACC	3	+99	+84	866
332773	GCCTC <u>A</u> T <u>A</u> T <u>A</u> TCGCACC	4	+75	+80	867
332774	GCCTC <u>A</u> T <u>A</u> T <u>A</u> T <u>A</u> GCACC	5	+62	+66	868
332775	GCCTC <u>A</u> T <u>A</u> T <u>A</u> T <u>A</u> T <u>A</u> CACC	6	-1	+10	869
332776	GCCT <u>A</u> <u>A</u> T <u>A</u> T <u>A</u> T <u>A</u> T <u>A</u> CACC	7	+10	+20	870
<b>Uniform 2'-MOE oligonucleotide</b>					
332769	GCCTCAGTCTGCTTCGCAC C	0	-11	-14	247

[0450] The results of treatment of HepG2 cells with the compounds in Table 40 reveals that none of the compounds displays the dose-dependent inhibition observed following treatment with the parent ISIS 301012 sequence. ISIS 332770, which has only a single thymidine to cytosine substitution in the center of the oligonucleotide, was 3-fold less potent than ISIS 301012. Further nucleotide substitutions abrogated antisense inhibition of apolipoprotein B expression.

[0451] Phosphorothioate chimeric oligonucleotides are metabolized in vivo predominantly by endonucleolytic cleavage. In accordance with the present invention, a series of oligonucleotides was designed by truncating the ISIS 301012 sequence in 1 or 2 base increments from the 5' and/or 3' end. The truncated oligonucleotides represent the possible products that result from endonucleolytic cleavage. These compounds are shown in Table 41. The compounds in Table 41 are chimeric oligonucleotides ("gapmers") of varying lengths, composed of a central "gap" region consisting of 2'-deoxynucleotides, which is flanked on both ends by 2'-methoxyethyl (2'-MOE)nucleotides. The exact structure of each chimeric oligonucleotide is designated in Table 41 as the "chimera structure". For example, a designation of 4-10-4 indicates that the first 4 (5' most) and last 4 (3' most) nucleotides are 2'-MOE nucleotides, and the 10 nucleotides in the gap are 2'-deoxynucleotides. 2'-MOE nucleotides are indicated by bold type. The internucleoside (backbone) linkages are phosphodiester (P=O) between underscored nucleotides; all other internucleoside linkages are phosphorothioate (P=S).

[0452] These compounds were tested for their ability to reduce the expression of apolipoprotein B mRNA. HepG2 cells were treated with 10, 50 or 150 nM of each antisense compound in Table 41 for a 24 hour period, after which RNA was isolated and target expression was measured by real-time PCR as described herein. Untreated cells served as controls. The results are shown in Tables 41 and are normalized to untreated control samples.

EP 2 336 318 A1

Table 41 Effect of ISIS 301012 truncation mutants on apolipoprotein B expression in HepG2 cells

ISIS #	Target SEQ ID NO	Target Site	SEQUENCE	Chimeric structure	% Change in apolipoprotein B mRNA expression			SEQ ID NO
					Dose of oligonucleotide			
					10	50	150	
301012	3	3249	GCCTCAGTCTGCTTCGCACC	5~10~5	-51	-72	-92	247
331022	3	3249	GCCTCAGTCTGCTTCGCAC	5~10~4	-33	-49	-87	871
332777	3	3249	GCCTCAGTCTGCTTCGCA	5~10~3	-27	-53	-80	872
332778	3	3249	GCCTCAGTCTGCTTC	5~10~0	-11	-20	-58	873
332780	3	3248	CCTCAGTCTGCTTCGCAC	4~10~4	-3	-43	-74	874
332781	3	3247	CTCAGTCTGCTTCGCA	3~10~3	-9	-35	-60	875
332782	3	3246	TCAGTCTGCTTCGC	2~10~2	-16	-16	-69	876
332784	3	3249	GCCTCAGTCT	5~5~0	+12	-1	+7	877
332785	3	3238	GCTTCGCACC	0~5~5	+5	-2	-4	878

[0453] The results in Table 41 illustrate that inhibition of apolipoprotein B is dependent upon sequence length, as well as upon sequence complementarity and dose, as demonstrated in Table 41, but truncated versions of ISIS 301012 are to a certain degree capable of inhibiting apolipoprotein B mRNA expression.

Example 60

Design and screening of dsRNAs targeting human apolipoprotein B

[0454] In accordance with the present invention, a series of nucleic acid duplexes comprising the antisense compounds of the present invention and their complements were designed to target apolipoprotein B and are shown in Table 42. All compounds in Table 42 are oligoribonucleotides 20 nucleotides in length with phosphodiester internucleoside linkages (backbones) throughout the compound. The compounds were prepared with blunt ends. Table 41 shows the antisense strand of the dsRNA, and the sense strand is synthesized as the complement of the antisense strand. These sequences are shown to contain uracil (U) but one of skill in the art will appreciate that uracil (U) is generally replaced by thymine (T) in DNA sequences. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the compound binds. A subset of the compounds in Table 42 are the RNA equivalents of DNA antisense oligonucleotides described herein, and, where applicable, this is noted by the ISIS # of the DNA oligonucleotide in the column "RNA equivalent of ISIS #".

Table 42 dsRNAs targeted to human apolipoprotein B

ISIS #	Region	Target SEQ ID NO	Target Site	Sequence	SEQ ID NO	RNA equivalent of ISIS #
42855	coding	3	3249	GCCUCAGUCUGCUUCGCACC	247	301012
342856	3' UTR	3	13903	GCUCACUGUAUGGUUUUAUC	262	301027
342857	coding	3	5589	AGGUUACCAGCCACAUGCAG	224	308361
342858	coding	3	669	GAGCAGUUCCAUAACACGGU	130	270991
342859	coding	3	1179	CCUCUCAGCUCAGUAACCAG	135	270996
342860	coding	3	2331	GUAUAGCCAAAGUGGUCCAC	34	147797
342861	coding	3	3579	UAAGCUGUAGCAGAUGAGUC	213	281614
342862	5' UTR	3	6	CAGCCCCGCAGGUCCCGGUG	249	301014



EP 2 336 318 A1

(continued)

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

ISIS #	Region	Target SEQ ID NO	Target Site	Sequence	SEQ ID NO	RNA equivalent of ISIS #
342863	5' UTR	3	116	GGUCCAUCGCCAGCUGCGGU	256	301021
342864	3' UTR	3	13910	AAGGCUGGCUCACUGUAUGG	266	301031
342865	3' UTR	3	13970	GCCAGCUUUGGUGCAGGUCC	273	301038
342866	coding	3	426	UUGAAGCCAUCACCCUCUUU	879	none
342867	coding	3	3001	UGACCAGGACUGCCUGUUCU	880	none
342868	coding	3	5484	GAAUAGGGCUGUAGCUGUAA	881	none
342869	coding	3	6662	UAUACUGAUCAAUUGUAUC	882	none
342870	coding	3	8334	UGGAAUUCUGGUAUGUGAAG	883	none
342871	coding	3	9621	AAAUCAAUGAUUGCUUUGU	883	none
342872	coding	3	10155	GUGAUGACACUUGAUUUAAA	885	none
342873	coding	3	12300	GAAGCUGCCUCUUCUCCCA	886	none
342874	coding	3	13629	GAGAGUUGGUCUGAAAAAUC	887	none

**[0455]** The dsRNA compounds in Table 42 were tested for their effects on human apolipoprotein mRNA in HepG2 cells. HepG2 cells were treated with 100 nM of dsRNA compounds mixed with 5 µg/mL LIPOFECTIN (Invitrogen Corporation, Carlsbad, CA) for a period of 16 hours. In the same experiment, HepG2 cells were also treated with 150 nM of subset of the antisense oligonucleotides described herein mixed with 3.75 µg/mL LIPOFECTIN; these compounds are listed in Table 43. Control oligonucleotides included ISIS 18078 (GTGCGCGGAGCCCGAAATC, SEQ ID NO: 888). ISIS 18078 is a chimeric oligonucleotide ("gapmer") 20 nucleotides in length, composed of a central "gap" region consisting of 9 2'-deoxynucleotides, which is flanked on the 5' and 3' ends by a five-nucleotide "wing" and a six-nucleotide "wing", respectively. The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidines are 5-methylcytidines.

**[0456]** The duplex of ISIS 263188 (CUUCUGGCAUCCGGUUUAGTT, SEQ ID NO: 889) and its complement was also used as a control. ISIS 263188 is an oligoribonucleotide 21 nucleotides in length with the 2 nucleotides on the 3' end being oligodeoxyribonucleotides (TT) and with phosphodiester internucleoside linkages (backbones) throughout the compound.

**[0457]** Cells were treated for 4 hours, after which human apolipoprotein B mRNA expression was measured as described by examples herein. Results were normalized to untreated control cells, which were not treated with LIPOFECTIN or oligonucleotide. Data are the average of 4 experiments and are presented in Table 43.

Table 43 Inhibition of apolipoprotein B mRNA by dsRNAs in HepG2 cells

ISIS #	Dose	% Inhibition	SEQ ID #
342855	100 nM	53	247
342856	100 nM	34	262
342857	100 nM	55	224
342858	100 nM	44	130
342859	100 nM	23	135
342860	100 nM	34	34
342861	100 nM	42	213
342862	100 nM	16	249
342863	100 nM	34	256
342864	100 nM	53	266

EP 2 336 318 A1

(continued)

ISIS #	Dose	% Inhibition	SEQ ID #
342865	100 nM	50	273
342866	100 nM	12	879
342867	100 nM	26	880
342868	100 nM	36	881
342869	100 nM	78	882
342870	100 nM	71	883
342871	100 nM	9	883
342872	100 nM	2	885
342873	100 nM	53	886
342874	100 nM	73	887
281625	150 nM	79	224
301012	150 nM	77	247
301014	150 nM	88	249
301021	150 nM	67	256
301027	150 nM	79	262
301028	150 nM	85	263
301029	150 nM	77	264
301030	150 nM	70	265
301031	150 nM	73	266
301037	150 nM	80	272
301038	150 nM	84	273
301045	150 nM	77	280
263188	150 nM	26	888
18078	150 nM	13	889

**Example 61**

**Antisense inhibition of apolipoprotein B in Cynomolgus monkey primary hepatocytes**

**[0458]** As demonstrated in Example 46, the region containing the target site to which ISIS 301012 hybridizes shares 96% identity with the corresponding region of Cynomolgus monkey apolipoprotein B mRNA sequence. ISIS 301012 contains two mismatched nucleotides relative to the Cynomolgus monkey apolipoprotein B mRNA sequence to which it hybridizes. In a further embodiment of the invention, oligonucleotides were designed to target regions of the monkey apolipoprotein B mRNA, using the partial Cynomolgus monkey apolipoprotein B sequence described herein (SEQ ID NO: 855) and an additional portion of Cynomolgus monkey apolipoprotein B RNA sequence, incorporated herein as SEQ ID NO: 890. The target site indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. For ISIS 326358 (GCCTCAGTCTGCTTTACACC, SEQ ID NO: 891) the target site is nucleotide 168 of SEQ ID NO: 855 and for ISIS 315089 (AGATTACCAGCCATATGCAG, SEQ ID NO: 892) the target site is nucleotide 19 of SEQ ID NO: 890. ISIS 326358 and ISIS 315089 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. ISIS 326358 and ISIS 315089 are the Cynomolgus monkey equivalents of the human apolipoprotein B antisense oligonucleotides ISIS 301012 (SEQ ID NO: 247) and ISIS 281625 (SEQ ID NO: 224), respectively.

EP 2 336 318 A1

[0459] Antisense inhibition by ISIS 301012 was compared to that of ISIS 326358, which is a perfect match to the Cynomolgous monkey apolipoprotein B sequence to which ISIS 301012 hybridizes. The compounds were analyzed for their effect on Cynomolgous monkey apolipoprotein B mRNA levels in primary Cynomolgous monkey hepatocytes purchased from In Vitro Technologies (Gaithersburg, MD). Pre-plated primary Cynomolgous monkey hepatocytes were purchased from Invitro Technologies (Baltimore, MD). Cells were cultured in high-glucose DMEM (Invitrogen Corporation, Carlsbad, CA) supplemented with 10% fetal bovine serum (Invitrogen Corporation, Carlsbad, CA), 100 units/mL and 100 µg/mL streptomycin (Invitrogen Corporation, Carlsbad, CA).

[0460] Primary Cynomolgous monkey hepatocytes were treated with 10, 50, 150 or 300 nM of antisense oligonucleotides for 48 hours. ISIS 113529 (SEQ ID NO: 859) was used as a control oligonucleotide. Untreated cells also served as a control. Cynomolgous monkey apolipoprotein B mRNA levels were quantitated by real-time PCR using the human apolipoprotein B and GAPDH primers and probes described by other examples herein. The results, shown in Table 44, are the average of 6 experiments and are expressed as percent inhibition of apolipoprotein B mRNA normalized to untreated control cells.

Table 44 Inhibition of Cynomolgous monkey apolipoprotein B mRNA by ISIS 301012 and ISIS 326358

		% Inhibition of apolipoprotein B mRNA		
		ISIS #		
Dose of oligonucleotide	Time of treatment (hours)	326358 treatment	301012	113529
10 nM	24	35	24	N.D.
	48	85	76	N.D.
50 nM	24	66	60	N.D.
	48	88	77	N.D.
150 nM	24	61	56	5
	48	82	88	42
300 nM	24	64	61	19
	48	87	86	13

[0461] These data demonstrate that both ISIS 326359 and ISIS 301012 (despite two mismatches with the Cynomolgous monkey apolipoprotein B sequence) can inhibit the expression of apolipoprotein B mRNA in cynomolgous monkey primary hepatocytes, in a dose- and time-dependent manner.

[0462] Apolipoprotein B protein secreted from primary Cynomolgous hepatocytes treated with 150 and 300 nM of oligonucleotide was measured by ELISA using an apolipoprotein B protein specific kit (ALerCHEK Inc., Portland, ME). Each result represents the average of 3 experiments. The data are normalized to untreated control cells and are shown in Table 45.

Table 45 Reduction in apolipoprotein B protein secreted from Cynomolgous monkey hepatocytes following antisense oligonucleotide treatment

		% Reduction in secreted apolipoprotein B protein		
		ISIS #		
Dose of oligonucleotide	Time of treatment (hours)	326358	301012	113529
150 nM	24	21	31	11
	48	29	25	18
300 nM	24	17	10	12
	48	35	17	8

[0463] These results demonstrate that antisense inhibition by ISIS 301012 and ISIS 326358 leads to a decrease in the secretion of apolipoprotein B protein from cultured primary Cynomolgous hepatocytes.

[0464] Additionally, protein was isolated from oligonucleotide-treated primary Cynomolgous monkey hepatocytes and

**EP 2 336 318 A1**

subjected to immunoblot analysis to further assess apolipoprotein B protein expression. Immunoblotting was performed as described herein, using an antibody to human apolipoprotein B protein (US Biologicals, Swampscott, MA). Immunoblot analysis of apolipoprotein B expression following antisense oligonucleotide treatment with ISIS 326358 and ISIS 301012 reveals a substantial reduction in apolipoprotein B expression.

**[0465]** In a further embodiment of the invention, antisense inhibition by ISIS 281625 was compared to that by ISIS 315089, which is a perfect match to the Cynomolgous monkey apolipoprotein B sequence to which ISIS 281625 hybridizes. Primary Cynomolgous monkey hepatocytes, cultured as described herein, were treated with 10, 50, 150 or 300 nM of ISIS 315089 or ISIS 281625 for 24 hours. Cells were treated with the control oligonucleotide ISIS 13650 (SEQ ID NO: 806) at 150 and 300 nM or ISIS 113529 (SEQ ID NO: 859) at 300 nM. Untreated cells also served as a control. Cynomolgous monkey apolipoprotein B mRNA levels in primary Cynomolgous monkey hepatocytes was quantitated using real-time PCR with human primers and probe as described by other examples herein. The results, shown in Table 46, are the average of 3 experiments and are expressed as percent inhibition of apolipoprotein B mRNA normalized to untreated control cells. Where present, a "+" preceding the value indicates that mRNA expression was increased.

Table 46 Antisense inhibition of apolipoprotein B mRNA expression in Cynomolgous monkey hepatocytes

Dose of oligonucleotide	% Inhibition of apolipoprotein B mRNA			
	ISIS #			
	315089	281625	13650	113529
10 nM	70	+5	N.D.	N.D.
50 nM	83	41	N.D.	N.D.
150 nM	81	35	+50	N.D.
300 nM	82	69	33	28

**[0466]** These data demonstrate that both ISIS 315089 and ISIS 281625 can inhibit the expression of apolipoprotein B mRNA in Cynomolgous monkey primary hepatocytes, in a dose-dependent manner.

**[0467]** Apolipoprotein B protein secreted primary Cynomolgous hepatocytes treated with 50 and 150 nM of ISIS 315089 and ISIS 281625 was measured by ELISA using an apolipoprotein B protein specific kit (ALerCHEK Inc., Portland, ME). Each result represents the average of 3 experiments. The data are normalized to untreated control cells and are shown in Table 47.

Table 47 Reduction in apolipoprotein B protein secreted from Cynomolgous monkey hepatocytes following antisense oligonucleotide treatment

Dose of oligonucleotide	% Reduction of monkey apolipoprotein B protein secretion			
	ISIS #			
	315089	281625	13650	113529
50 nM	11	6	16	N.D.
150 nM	25	13	13	12

**[0468]** These results demonstrate that antisense inhibition by 150 nM of ISIS 315089 leads to a decrease in the secretion of apolipoprotein B protein from cultured primary Cynomolgous hepatocytes.

**[0469]** ISIS 271009 (SEQ ID NO: 319) and ISIS 301027 (SEQ ID NO: 262) were also tested for their effects on apolipoprotein B mRNA and protein expression in Cynomolgous primary hepatocytes. Cells, cultured as described herein, were treated with 10, 50 and 150 nM of ISIS 271009 or ISIS 301027 for 24 hours. Cells were treated with the control oligonucleotide ISIS 113529 (SEQ ID NO: 859) at 150 nM. Untreated cells also served as a control. Cynomolgous monkey apolipoprotein B mRNA levels in primary Cynomolgous monkey hepatocytes was quantitated using real-time PCR with human primers and probe as described by other examples herein. The results, shown in Table 48, are the average of 2 experiments and are expressed as percent inhibition of apolipoprotein B mRNA normalized to untreated control cells.

## EP 2 336 318 A1

Table 48 Antisense inhibition of apolipoprotein B mRNA expression in Cynomolgous monkey hepatocytes

	% Inhibition of apolipoprotein B mRNA		
	ISIS #		
Dose of oligonucleotide	271009	301027	113529
10 nM	42	40	N.D.
50 nM	66	54	N.D.
150 nM	69	67	11

[0470] These data demonstrate that both ISIS 271009 and ISIS 301027 can inhibit the expression of apolipoprotein B mRNA in Cynomolgous monkey primary hepatocytes, in a dose-dependent manner.

[0471] Apolipoprotein B protein secreted from primary Cynomolgous hepatocytes treated with 50 and 150 nM of ISIS 271009 and ISIS 301027 was measured by ELISA using an apolipoprotein B protein specific kit (ALerCHEK Inc., Portland, ME). Each result represents the average of 3 experiments. The data are shown as percent reduction in secreted protein, normalized to untreated control cells, and are shown in Table 49. Where present, a "+" indicates that protein secretion was increased.

Table 49 Reduction in apolipoprotein B protein secreted from Cynomolgous monkey hepatocytes following antisense oligonucleotide treatment

	% Reduction of monkey apolipoprotein B protein secretion			
	ISIS #			
Dose of oligonucleotide	271009	301027	13650	113529
50 nM	+30	25	N.D.	N.D.
150 nM	26	31	+1	15

[0472] These results demonstrate that antisense inhibition by ISIS 315089 and ISIS 281625 leads to a decrease in the secretion of apolipoprotein B protein from cultured primary Cynomolgous hepatocytes.

### Example 62

#### Methods for evaluating hepatic steatosis

[0473] Hepatic steatosis refers to the accumulation of lipids in the liver, or "fatty liver", which is frequently caused by alcohol consumption, diabetes and hyperlipidemia. Livers of animals treated with antisense oligonucleotides targeted to apolipoprotein B were evaluated for the presence of steatosis. Steatosis is assessed by histological analysis of liver tissue and measurement of liver triglyceride levels.

[0474] Tissue resected from liver is immediately immersed in Tissue Tek OCT embedding compound (Ted Pella, Inc., Redding, CA) and frozen in a 2-methyl-butane dry ice slurry. Tissue sections are cut at a thickness of 4-5  $\mu$ m and then fixed in 5% neutral-buffered formalin. Tissue sections are stained with hematoxylin and eosin following standard histological procedures to visualize nuclei and cytoplasm, respectively, and oil red O according to the manufacturer's instructions (Newcomers Supply, Middleton, WI) to visualize lipids.

[0475] Alternatively, tissues are fixed in 10% neutral-buffered formalin, embedded in paraffin, sectioned at a thickness of 4-5  $\mu$ m, deparaffinized and stained with hematoxylin and eosin, all according to standard histological procedures.

[0476] Quantitation of liver triglyceride content is also used to assess steatosis. Tissue triglyceride levels are measured using a Triglyceride GPO Assay (Sigma-Aldrich, St. Louis, MO).

### Example 63

#### Effects of antisense inhibition by ISIS 301012 in lean mice: long-term study

[0477] In accordance with the present invention, the toxicity of ISIS 301012 (SEQ ID NO: 247) is investigated in a long-term, 3 month study in mice. Two-month old male and female CD-1 mice (Charles River Laboratories, Wilmington,

MA) are dosed with 2, 5, 12.5, 25 or 50 mg/kg of ISIS 301012 twice per week for first week, and every 4 days thereafter. The mice are maintained on a standard rodent diet. Saline and control oligonucleotide animals serve as controls and are injected on the same schedule. Each treatment group contains 6 to 10 mice of each sex, and each treatment group is duplicated, one group for a 1 month study termination, the other for a 3 month study termination. After the 1 or 3 month treatment periods, the mice are sacrificed and evaluated for target expression in liver, lipid levels in serum and indicators of toxicity. Liver samples are procured, RNA is isolated and apolipoprotein B mRNA expression is measured by real-time PCR as described in other examples herein. Serum lipids, including total cholesterol, LDL-cholesterol, HDL-cholesterol and triglycerides, are evaluated by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY). Ratios of LDL-cholesterol to HDL-cholesterol and total cholesterol to HDL-cholesterol are also calculated. Analyses of serum ALT and AST, inflammatory infiltrates in tissue and basophilic granules in tissue provide an assessment of toxicities related to the treatment. Hepatic steatosis, or accumulation of lipids in the liver, is assessed by routine histological analysis with oil red O stain and measurement of liver tissue triglycerides using a Triglyceride GPO Assay (Sigma-Aldrich, St. Louis, MO).

**[0478]** The toxicity study also includes groups of animals allowed to recover following cessation of oligonucleotide treatment. Both male and female CD-1 mice (Charles River Laboratories, Wilmington, MA) are treated with 5, 10, 50 mg/kg of ISIS 301012 twice per week for the first week and every 4 days thereafter. Saline and control oligonucleotide injected animals serve as controls. Each treatment group includes 6 animals per sex. After 3 months of treatment, animals remain untreated for an additional 3 months, after which they are sacrificed. The same parameters are evaluated as in the mice sacrificed immediately after 3 months of treatment.

**[0479]** After one month of treatment, real-time PCR quantitation reveals that mouse apolipoprotein B mRNA levels in liver are reduced by 53%. Additionally, the expected dose-response toxicities were observed. ALT and AST levels, measured by routine clinical procedures on an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY), are increased in mice treated with 25 or 50 mg/kg of ISIS 301012. Tissues were prepared for analysis by routine histological procedures. Basophilic granules in liver and kidney tissue were observed at doses of ISIS 301012 above 12.5 mg/kg. Mild lymphohistiocytic infiltrates were observed in various tissues at doses greater than 12.5 mg/kg of ISIS 301012. Staining of tissue sections with oil red O reveals no steatosis present following the oligonucleotide treatments.

#### **Example 64**

##### **Effects of antisense inhibition by ISIS 301012 in lean Cynomolgous monkeys: long-term study**

**[0480]** As discussed in Example 45, Cynomolgous monkeys (male or female) are used to evaluate antisense oligonucleotides for their potential to lower apolipoprotein B mRNA or protein levels, as well as phenotypic endpoints associated with apolipoprotein B including, but not limited to cardiovascular indicators, atherosclerosis, lipid diseases, obesity, and plaque formation. Accordingly, in a further embodiment of the invention, ISIS 301012 (SEQ ID NO: 247) is investigated in a long-term study for its effects on apolipoprotein B expression and serum lipids in Cynomolgous monkeys. Such a long-term study is also used to evaluate the toxicity of antisense compounds.

**[0481]** Male and female Cynomolgous monkeys are treated with 2, 4 or 12 mg/kg of ISIS 301012 intravenously or 2 or 20 mg/kg subcutaneously at a frequency of every two days for the first week, and every 4 days thereafter, for 1 and 3 month treatment periods. Saline-treated animals serve as controls. Each treatment group includes 2 to 3 animals of each sex.

**[0482]** At a one month interval and at the 3 month study termination, the animals are sacrificed and evaluated for target expression in liver, lipid levels in serum and indicators of toxicity. Liver samples are procured, RNA is isolated and apolipoprotein B mRNA expression is measured by real-time PCR as described in other examples herein. Serum lipids, including total cholesterol, LDL-cholesterol, HDL-cholesterol and triglycerides, are evaluated by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY). Ratios of LDL-cholesterol to HDL-cholesterol and total cholesterol to HDL-cholesterol are also calculated. Analyses of serum ALT and AST, inflammatory infiltrates in tissue and basophilic granules in tissue provide an assessment of toxicities related to the treatment. Hepatic steatosis, or accumulation of lipids in the liver, is assessed by routine histological analysis with oil red O stain and measurement of liver tissue triglycerides using a Triglyceride GPO Assay (Sigma-Aldrich, St. Louis, MO).

**[0483]** Additional treatment groups consisting of 2 animals per sex are treated with saline (0 mg/kg), 12 or 20 mg/kg ISIS 301012 at a frequency of every two days for the first week, and every 4 days thereafter, for a 3 month period. Following the treatment period, the animals receive no treatment for an additional three months. These treatment groups are for the purpose of studying the effects of apolipoprotein B inhibition 3 months after cessation of treatment. At the end of the 3 month recovery period, animals are sacrificed and evaluated for the same parameters as the animals sacrificed immediately after 1 and 3 months of treatment.

**[0484]** The results from the one month interval of the long term treatment are shown in Table 50 and are normalized to saline-treated animals for mRNA and to untreated baseline values for lipid levels. Total cholesterol, LDL-cholesterol,

## EP 2 336 318 A1

HDL-cholesterol, LDL particle concentration and triglyceride levels in serum were measured by nuclear magnetic resonance spectroscopy by Liposcience (Raleigh, NC). Additionally, the concentration of intact oligonucleotide in liver was measured by capillary gel electrophoresis and is presented as micrograms of oligonucleotide per gram of liver tissue. Each result represents the average of data from 4 animals (2 males and 2 females).

Table 50 Effects of antisense inhibition by ISIS 301012 in lean Cynomolgous monkeys

	Intravenous delivery			Subcutaneous injection		
	2 mg/kg	4 mg/kg	12 mg/kg	3.5 mg/kg	20 mg/kg	
<b>apolipoprotein B expression % change normalized to saline</b>	-45	-76	-96	N.D.	-94	
<b>antisense oligonucleotide concentration <math>\mu\text{g/g}</math></b>	92	179	550	N.D.	855	
<b>Lipid parameters, % change normalized to untreated baseline value</b>	<b>saline</b>	<b>2 mg/kg</b>	<b>4 mg/kg</b>	<b>12 mg/kg</b>	<b>3.5 mg/kg</b>	<b>20 mg/kg</b>
<b>Total cholesterol</b>	+1	-6	-2	-2	+5	-5
<b>LDL-cholesterol</b>	+17	+15	+9	+3	-4	-16
<b>HDL-cholesterol</b>	-11	-23	-15	-8	+13	+5
<b>LDL/HDL</b>	+62	+94	+38	+44	-15	-19
<b>Total cholesterol/HDL</b>	+30	+44	+22	+21	-7	-10
<b>Triglyceride</b>	+37	+26	+32	+15	+1	-3
<b>LDL Particle concentration</b>	+15	+8	+8	-11	-14	-21

[0485] These data show that ISIS 301012 inhibits apolipoprotein B expression in a dose-dependent manner in a primate species and concomitantly lowers lipid levels at higher doses of ISIS 301012. Furthermore, these results demonstrate that antisense oligonucleotide accumulates in the liver in a dose-dependent manner.

[0486] Hepatic steatosis, or accumulation of lipids in the liver, was not observed following 4 weeks of treatment with the doses indicated. Expected dose-related toxicities were observed at the higher doses of 12 and 20 mg/kg, including a transient 1.2-1.3 fold increase in activated partial thromboplastin time (APTT) during the first 4 hours and basophilic granules in the liver and kidney (as assessed by routine histological examination of tissue samples). No functional changes in kidney were observed.

[0487] In a similar experiment, male and female Cynomolgous monkeys received an intravenous dose of ISIS 301012 at 4 mg/kg, every two days for the first week and every 4 days thereafter. Groups of animals were sacrificed after the first dose and the fourth dose, as well as 11, 15 and 23 days following the fourth and final dose. Liver RNA was isolated and apolipoprotein B mRNA levels were evaluated by real-time PCR as described herein. The results of this experiment demonstrate a 40% reduction in apolipoprotein B mRNA expression after a single intravenous dose of 4 mg/kg ISIS 301012. Furthermore, after 4 doses of ISIS 301012 at 4 mg/kg, target mRNA was reduced by approximately 85% and a 50% reduction in target mRNA was sustained for up to 16 days following the cessation of antisense oligonucleotide treatment.

### Example 65

#### Microarray analysis: gene expression patterns in lean versus high-fat fed mice

[0488] Male C57B1/6 mice were divided into the following groups, consisting of 5 animals each: (1) mice on a lean diet, injected with saline (lean control); (2) mice on a high fat diet; (3) mice on a high fat diet injected with 50 mg/kg of the control oligonucleotide 141923 (SEQ ID NO: 858); (4) mice on a high fat diet given 20 mg/kg atorvastatin calcium (Lipitor®, Pfizer Inc.); (5) mice on a high fat diet injected with 10, 25 or 50 mg/kg ISIS 147764 (SEQ ID NO: 109). Saline and oligonucleotide treatments were administered intraperitoneally twice weekly for 6 weeks. Atorvastatin was administered daily for 6 weeks. At study termination, liver samples were isolated from each animal and RNA was isolated for

Northern blot qualitative assessment, DNA microarray and quantitative real-time PCR. Northern blot assessment and quantitative real-time PCR were performed as described by other examples herein.

**[0489]** For DNA microarray analysis, hybridization samples were prepared from 10 µg of total RNA isolated from each mouse liver according to the Affymetrix Expression Analysis Technical Manual (Affymetrix, Inc., Santa Clara, CA). Samples were hybridized to a mouse gene chip containing approximately 22,000 genes, which was subsequently washed and double-stained using the Fluidics Station 400 (Affymetrix, Inc., Santa Clara, CA) as defined by the manufacturer's protocol. Stained gene chips were scanned for probe cell intensity with the GeneArray scanner (Affymetrix, Inc., Santa Clara, CA). Signal values for each probe set were calculated using the Affymetrix Microarray Suite v5.0 software (Affymetrix, Inc., Santa Clara, CA). Each condition was profiled from 5 biological samples per group, one chip per sample. Fold change in expression was computed using the geometric mean of signal values as generated by Microarray Suite v5.0. Statistical analysis utilized one-way ANOVA followed by 9 pair-wise comparisons. All groups were compared to the high fat group to determine gene expression changes resulting from ISIS 147764 treatment. Microarray data was interpreted using hierarchical clustering to visualize global gene expression patterns.

**[0490]** The results of the microarray analysis reveal that treatment with ISIS 147764 drives the gene expression profile in high fat fed mice to the profile observed in lean mice. Real-time PCR analysis confirmed the reduction in mRNA expression for the following genes involved in the lipid metabolism: hepatic lipase, fatty acid synthase ATP-binding cassette, sub-family D (ALD) member 2, intestinal fatty acid binding protein 2, stearol CoA desaturase-1 and HMG CoA reductase.

**[0491]** Mouse apolipoprotein B mRNA and serum cholesterol levels, measured as described herein, were evaluated to confirm antisense inhibition by ISIS 147764 and ISIS 147483. Both mRNA and cholesterol levels were lowered in a dose-dependent manner following treatment with ISIS 147764 or ISIS 147483, as demonstrated in other examples herein. The 50 mg/kg dose of ISIS 147483 increased ALT and AST levels. The 10, 25 and 50 mg/kg doses of ISIS 147764 and the 10 and 25 mg/kg doses of ISIS 147483 did not significantly elevate ALT or AST levels.

## Example 66

### Evaluation of hepatic steatosis in animals treated with apolipoprotein B antisense oligonucleotides

**[0492]** Livers of animals treated with antisense oligonucleotides targeted to apolipoprotein B were evaluated for the presence of steatosis. Steatosis is assessed by histological analysis of liver tissue and measurement of liver triglyceride levels.

### Evaluation of steatosis in high fat fed animals treated with ISIS 147764 for 6 weeks

**[0493]** Liver tissue from ISIS 147764 (SEQ ID NO: 109) and control-treated animals described in Example 21 was evaluated for steatosis at study termination following 6 weeks of treatment. Tissue sections were stained with oil red O and hematoxylin to visualize lipids and nuclei, respectively. Tissue sections were also stained with hematoxylin and eosin to visualize nuclei and cytoplasm, respectively. Histological analysis of tissue sections stained by either method reveal no difference in steatosis between saline treated and ISIS 147764 treated animals, demonstrating that a 6 week treatment with ISIS 147764 does not lead to accumulation of lipids in the liver.

### Evaluation of steatosis following long-term treatment with apolipoprotein B inhibitor in high-fat fed animals

**[0494]** Male C57B1/6 mice were treated with twice weekly intraperitoneal injections of 25 mg/kg ISIS 147764 (SEQ ID NO: 109) or 25 mg/kg ISIS 141923 (SEQ ID NO: 858) for 6, 12 and 20 weeks. Saline treated animals served as controls. Each treatment group contained 4 animals. Animals were sacrificed at 6, 12 and 20 weeks and liver tissue was procured for histological analysis and measurement of tissue triglyceride content. The results reveal no significant differences in liver tissue triglyceride content when ISIS 147764 treated animals are compared to saline treated animals. Furthermore, histological analysis of liver tissue section demonstrates that steatosis is reduced at 12 and 20 weeks following treatment of high fat fed mice with ISIS 147764, in comparison to saline control animals that received a high fat diet.

### Evaluation of steatosis in lean mice

**[0495]** The accumulation of lipids in liver tissue was also evaluated in lean mice. Male C67B1/6 mice (Charles River Laboratories (Wilmington, MA) at 6 to 7 weeks of age were maintained on a standard rodent diet and were treated twice weekly with intraperitoneal injections of 25 or 50 mg/kg 147764 (SEQ ID NO: 109) or 147483 (SEQ ID NO: 79) for 6 weeks. Saline treated animals served as controls. Each treatment group was comprised of 4 animals. Animals were



sacrificed after the 6 week treatment period, at which point liver tissue and serum were collected.

[0496] Apolipoprotein B mRNA levels were measured by real-time PCR as described by other examples herein. The data, shown in Table 51, represent the average of 4 animals and are presented as inhibition relative to saline treated controls. The results demonstrate that both ISIS 147483 and ISIS 147764 inhibit apolipoprotein B mRNA expression in lean mice in a dose-dependent manner.

Table 51 Antisense inhibition of apolipoprotein B mRNA in lean mice

	Treatment and dose			
	ISIS 147483		ISIS 147764	
	25 mg/kg	50 mg/kg	25 mg/kg	50 mg/kg
<b>% inhibition apolipoprotein B mRNA</b>	79	91	48	77

[0497] Total cholesterol, LDL-cholesterol, HDL-cholesterol and triglycerides in serum were measured by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY). The liver enzymes ALT and ALT in serum were also measured using the Olympus Clinical Analyzer. These results demonstrate that ISIS 147764 lowers serum lipids relative to saline-treated control animals. ALT and AST levels do not exceed the normal range for mice (300 IU/L), indicating a lack of treatment-associated toxicity. The results are the average of data from 4 animals and are shown in Table 52.

Table 52 Serum lipids and liver enzyme levels in lean mice treated with ISIS 147764 and ISIS 147483

	Treatment and dose				
	Saline	ISIS 147483		ISIS 147764	
		25 mg/kg	50 mg/kg	25 mg/kg	50 mg/kg
<b>Serum lipids</b>					
Total cholesterol mg/dL	164	153	183	114	57
LDL-cholesterol mg/dL	25	26	39	29	18
HDL-cholesterol mg/dL	127	117	131	79	38
Triglycerides mg/dL	121	138	127	80	30
<b>Liver enzymes</b>					
ALT IU/L	105	73	57	47	48
AST IU/L	109	78	72	81	101

[0498] Liver tissue was prepared by routine histological methods to evaluate steatosis, as described herein. Examination of tissue samples stained with oil red O or hematoxylin and eosin reveals that treatment of lean mice with apolipoprotein B antisense oligonucleotides does not result in steatosis.

#### Six month study to further evaluate steatosis in mice treated with apolipoprotein B antisense oligonucleotides

[0499] A long-term treatment of mice with antisense oligonucleotides targeted to apolipoprotein B is used to evaluate the toxicological and pharmacological effects of extended treatment with antisense compounds. Both male and female C57B1/6 mice at 2 months of age are treated with 2, 5, 25 or 50 mg/kg of apolipoprotein B antisense oligonucleotide. Treatments are administered intraperitoneally every 2 days for the first week and every 4 days thereafter. Mice treated with saline alone or control oligonucleotide serve as control groups. Each treatment group contains 25 to 30 mice. After 6 months of treatment, a subset of the mice in each treatment group is sacrificed. The remaining mice are allowed a 3 month recovery period without treatment, after which they are sacrificed. Apolipoprotein B mRNA expression in liver is measured by real-time PCR as described by other methods herein. Liver tissue is also prepared for measurement of triglyceride content using a Triglyceride GPO Assay (Sigma-Aldrich, St. Louis, MO). Serum is collected and evaluated for lipid content, including total cholesterol, LDL-cholesterol, HDL-cholesterol and triglyceride, using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY). The liver enzymes ALT and AST are also measured in serum, also using the clinical analyzer. Serum samples are subjected to immunoblot analysis using an antibody directed to apolipoprotein

B (Santa Cruz Biotechnology, Inc., Santa Cruz, CA). Liver, kidney and other tissues are prepared by routine procedures for histological analyses. Tissues are evaluated for the presence of basophilic granules and inflammatory infiltrates. Steatosis is evaluated by oil red O stain of liver tissue sections.

#### 5 **Example 67**

##### **A mouse model for atherosclerotic plaque formation: human apolipoprotein B transgenic mice lacking the LDL receptor gene**

10 **[0500]** The LDL receptor is responsible for clearing apolipoprotein B-containing LDL particles. Without the LDL receptor, animals cannot effectively clear apolipoprotein B-containing LDL particles from the plasma. Thus the serum levels of apolipoprotein B and LDL cholesterol are markedly elevated. Mice expressing the human apolipoprotein B transgene (TgN-hApoB +/+) and mice deficient for the LDL receptor (LDLr -/-) are both used as animal models of atherosclerotic plaque development. When the LDL receptor deficiency genotype is combined with a human apolipoprotein B transgenic genotype (TgN-hApoB +/+; LDLr -/-), atherosclerotic plaques develop rapidly. In accordance with the present invention, mice of this genetic background are used to investigate the ability of compounds to prevent atherosclerosis and plaque formation.

15 **[0501]** Male TgN-hApoB +/+;LDLr -/- mice are treated twice weekly with 10 or 20 mg/kg of human apolipoprotein B antisense oligonucleotides for 12 weeks. Control groups are treated with saline or control oligonucleotide. Serum total cholesterol, HDL-cholesterol, LDL-cholesterol and triglycerides are measured at 2, 4, 6, 8 and 12 weeks by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY). Serum human apolipoprotein B protein is measured at 2, 4, 6, 8 and 12 weeks using an ELISA kit (ALerCHEK Inc., Portland, ME). Human and mouse apolipoprotein mRNA in liver is measured at 12 weeks. The results of the 12 week study serve to evaluate the pharmacological behavior of ISIS 301012 in a doubly transgenic model.

20 **[0502]** Additionally, a four month study is performed in TgN-hApoB +/+;LDLr -/- mice, with treatment conditions used in the 12 week study. Mice are treated for 4 months with antisense oligonucleotides targeted to human apolipoprotein B to evaluate the ability of such compounds to prevent atherosclerotic plaque formation. At the end of the 4 month treatment period, mice are anesthetized and perfused with 10% formalin. The perfused arterial tree is isolated and examined for the presence of atherosclerotic plaques. Sections of the arterial tree are embedded in paraffin and prepared for histological analysis using routine methods. Serum total cholesterol, HDL-cholesterol, LDL-cholesterol and triglycerides are measured at 2, 4, 6, 8, 12 and 16 weeks by routine clinical analysis using an Olympus Clinical Analyzer (Olympus America Inc., Melville, NY). Serum human apolipoprotein B protein is measured at 2, 4, 6, 8, 12 and 16 weeks using an ELISA kit (ALerCHEK Inc., Portland, ME). Human and mouse apolipoprotein mRNA in liver at 16 weeks is measured by real-time PCR.

#### 25 **Example 68**

##### **Rabbit models for study of atherosclerotic plaque formation**

30 **[0503]** The Watanabe heritable hyperlipidemic (WHHL) strain of rabbit is used as a model for atherosclerotic plaque formation. New Zealand white rabbits on a high-fat diet are also used as a model of atherosclerotic plaque formation. Treatment of WHHL or high fat fed New Zealand white rabbits with apolipoprotein B antisense compounds is used to test their potential as therapeutic or prophylactic treatments for atherosclerotic plaque disease. Rabbits are injected with 5, 10, 25 or 50 mg/kg of antisense oligonucleotides targeted to apolipoprotein B. Animals treated with saline alone or a control oligonucleotide serve as controls. Throughout the treatment, serum samples are collected and evaluated for apolipoprotein B protein levels by ELISA (kit from ALerCHEK Inc., Portland, ME) and serum lipids (cholesterol, LDL-cholesterol, VLDL-cholesterol, HDL-cholesterol, triglycerides) by routine clinical analysis. Liver tissue triglyceride content is measured using a Triglyceride GPO Assay (Sigma-Aldrich, St. Louis, MO). Liver, kidney, heart, aorta and other tissues are procured and processed for histological analysis using routine procedures. Liver and kidney tissues are examined for evidence of basophilic granules and inflammatory infiltrates. Liver tissue is evaluated for steatosis using oil red O stain. Additionally, aortic sections stained with oil red O stain and hematoxylin are examined to evaluate the formation of atherosclerotic lesions.

#### 35 **Example 69**

##### **Oral delivery of apolipoprotein B inhibitors**

40 **[0504]** Oligonucleotides may be formulated for delivery *in vivo* in an acceptable dosage form, e.g. as parenteral or

non-parenteral formulations. Parenteral formulations include intravenous (IV), subcutaneous (SC), intraperitoneal (IP), intravitreal and intramuscular (IM) formulations, as well as formulations for delivery via pulmonary inhalation, intranasal administration, topical administration, etc. Non-parenteral formulations include formulations for delivery via the alimentary canal, e.g. oral administration, rectal administration, intrajejunal instillation, etc. Rectal administration includes administration as an enema or a suppository. Oral administration includes administration as a capsule, a gel capsule, a pill, an elixir, etc.

**[0505]** In some embodiments, an oligonucleotide may be administered to a subject via an oral route of administration. The subject may be an animal or a human (man). An animal subject may be a mammal, such as a mouse, rat, mouse, a rat, a dog, a guinea pig, a monkey, a non-human primate, a cat or a pig. Non-human primates include monkeys and chimpanzees. A suitable animal subject may be an experimental animal, such as a mouse, rat, mouse, a rat, a dog, a monkey, a non-human primate, a cat or a pig.

**[0506]** In some embodiments, the subject may be a human. In certain embodiments, the subject may be a human patient in need of therapeutic treatment as discussed in more detail herein. In certain embodiments, the subject may be in need of modulation of expression of one or more genes as discussed in more detail herein. In some particular embodiments, the subject may be in need of inhibition of expression of one or more genes as discussed in more detail herein. In particular embodiments, the subject may be in need of modulation, i.e. inhibition or enhancement, of apolipoprotein B in order to obtain therapeutic indications discussed in more detail herein.

**[0507]** In some embodiments, non-parenteral (e.g. oral) oligonucleotide formulations according to the present invention result in enhanced bioavailability of the oligonucleotide. In this context, the term "bioavailability" refers to a measurement of that portion of an administered drug which reaches the circulatory system (e.g. blood, especially blood plasma) when a particular mode of administration is used to deliver the drug. Enhanced bioavailability refers to a particular mode of administration's ability to deliver oligonucleotide to the peripheral blood plasma of a subject relative to another mode of administration. For example, when a non-parenteral mode of administration (e.g. an oral mode) is used to introduce the drug into a subject, the bioavailability for that mode of administration may be compared to a different mode of administration, e.g. an IV mode of administration. In some embodiments, the area under a compound's blood plasma concentration curve ( $AUC_0$ ) after non-parenteral (e.g. oral, rectal, intrajejunal) administration may be divided by the area under the drug's plasma concentration curve after intravenous (i.v.) administration ( $AUC_{iv}$ ) to provide a dimensionless quotient (relative bioavailability, RB) that represents fraction of compound absorbed via the non-parenteral route as compared to the IV route. A composition's bioavailability is said to be enhanced in comparison to another composition's bioavailability when the first composition's relative bioavailability ( $RB_1$ ) is greater than the second composition's relative bioavailability ( $RB_2$ ).

**[0508]** In general, bioavailability correlates with therapeutic efficacy when a compound's therapeutic efficacy is related to the blood concentration achieved, even if the drug's ultimate site of action is intracellular (van Berge-Henegouwen et al., *Gastroenterol.*, 1977, 73, 300). Bioavailability studies have been used to determine the degree of intestinal absorption of a drug by measuring the change in peripheral blood levels of the drug after an oral dose (DiSanto, Chapter 76 In: Remington's *Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 1451-1458).

**[0509]** In general, an oral composition's bioavailability is said to be "enhanced" when its relative bioavailability is greater than the bioavailability of a composition substantially consisting of pure oligonucleotide, i.e. oligonucleotide in the absence of a penetration enhancer.

**[0510]** Organ bioavailability refers to the concentration of compound in an organ. Organ bioavailability may be measured in test subjects by a number of means, such as by whole-body radiography. Organ bioavailability may be modified, e.g. enhanced, by one or more modifications to the oligonucleotide, by use of one or more carrier compounds or excipients, etc. as discussed in more detail herein. In general, an increase in bioavailability will result in an increase in organ bioavailability.

**[0511]** Oral oligonucleotide compositions according to the present invention may comprise one or more "mucosal penetration enhancers," also known as "absorption enhancers" or simply as "penetration enhancers." Accordingly, some embodiments of the invention comprise at least one oligonucleotide in combination with at least one penetration enhancer. In general, a penetration enhancer is a substance that facilitates the transport of a drug across mucous membrane(s) associated with the desired mode of administration, e.g. intestinal epithelial membranes. Accordingly it is desirable to select one or more penetration enhancers that facilitate the uptake of an oligonucleotide, without interfering with the activity of the oligonucleotide, and in a such a manner the oligonucleotide can be introduced into the body of an animal without unacceptable side-effects such as toxicity, irritation or allergic response.

**[0512]** Embodiments of the present invention provide compositions comprising one or more pharmaceutically acceptable penetration enhancers, and methods of using such compositions, which result in the improved bioavailability of oligonucleotides administered via non-parenteral modes of administration. Heretofore, certain penetration enhancers have been used to improve the bioavailability of certain drugs. See Muranishi, *Crit. Rev. Ther. Drug Carrier Systems*, 1990, 7, 1 and Lee et al., *Crit. Rev. Ther. Drug Carrier Systems*, 1991, 8, 91. It has been found that the uptake and

delivery of oligonucleotides, relatively complex molecules which are known to be difficult to administer to animals and man, can be greatly improved even when administered by non-parenteral means through the use of a number of different classes of penetration enhancers.

5 [0513] In some embodiments, compositions for non-parenteral administration include one or more modifications from naturally-occurring oligonucleotides (i.e. full-phosphodiester deoxyribosyl or full-phosphodiester ribosyl oligonucleotides). Such modifications may increase binding affinity, nuclease stability, cell or tissue permeability, tissue distribution, or other biological or pharmacokinetic property. Modifications may be made to the base, the linker, or the sugar, in general, as discussed in more detail herein with regards to oligonucleotide chemistry.

10 [0514] In some embodiments of the invention, compositions for administration to a subject, and in particular oral compositions for administration to an animal or human subject, will comprise modified oligonucleotides having one or more modifications for enhancing affinity, stability, tissue distribution, or other biological property.

15 [0515] Suitable modified linkers include phosphorothioate linkers. In some embodiments according to the invention, the oligonucleotide has at least one phosphorothioate linker. Phosphorothioate linkers provide nuclease stability as well as plasma protein binding characteristics to the oligonucleotide. Nuclease stability is useful for increasing the *in vivo* lifetime of oligonucleotides, while plasma protein binding decreases the rate of first pass clearance of oligonucleotide via renal excretion. In some embodiments according to the present invention, the oligonucleotide has at least two phosphorothioate linkers. In some embodiments, wherein the oligonucleotide has exactly  $n$  nucleosides, the oligonucleotide has from one to  $n-1$  phosphorothioate linkages. In some embodiments, wherein the oligonucleotide has exactly  $n$  nucleosides, the oligonucleotide has  $n-1$  phosphorothioate linkages. In some embodiments wherein the oligonucleotide has exactly  $n$  nucleosides, and  $n$  is even, the oligonucleotide has from  $n/2$  phosphorothioate linkages, or, when  $n$  is odd, from 1 to  $(n-1)/2$  phosphorothioate linkages. In some embodiments, the oligonucleotide has alternating phosphodiester (PO) and phosphorothioate (PS) linkages. In other embodiments, the oligonucleotide has at least one stretch of two or more consecutive PO linkages and at least one stretch of two or more PS linkages. In other embodiments, the oligonucleotide has at least two stretches of PO linkages interrupted by at least one PS linkage.

25 [0516] In some embodiments, at least one of the nucleosides is modified on the ribosyl sugar unit by a modification that imparts nuclease stability, binding affinity or some other beneficial biological property to the sugar. In some cases, the sugar modification includes a 2'-modification, e.g. the 2'-OH of the ribosyl sugar is replaced or substituted. Suitable replacements for 2'-OH include 2'-F and 2'-arabino-F. Suitable substitutions for OH include 2'-*O*-alkyl, e.g. 2'-*O*-methyl, and 2'-*O*-substituted alkyl, e.g. 2'-*O*-methoxyethyl, 2'-*O*-aminopropyl, etc. In some embodiments, the oligonucleotide contains at least one 2'-modification. In some embodiments, the oligonucleotide contains at least 2 2'-modifications. In some embodiments, the oligonucleotide has at least one 2'-modification at each of the termini (i.e. the 3'- and 5'-terminal nucleosides each have the same or different 2'-modifications). In some embodiments, the oligonucleotide has at least two sequential 2'-modifications at each end of the oligonucleotide. In some embodiments, oligonucleotides further comprise at least one deoxynucleoside. In particular embodiments, oligonucleotides comprise a stretch of deoxynucleosides such that the stretch is capable of activating RNase (e.g. RNase H) cleavage of an RNA to which the oligonucleotide is capable of hybridizing. In some embodiments, a stretch of deoxynucleosides capable of activating RNase-mediated cleavage of RNA comprises about 6 to about 16, e.g. about 8 to about 16 consecutive deoxynucleosides.

30 [0517] Oral compositions for administration of non-parenteral oligonucleotide compositions of the present invention may be formulated in various dosage forms such as, but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The term "alimentary delivery" encompasses e.g. oral, rectal, endoscopic and sublingual/buccal administration. A common requirement for these modes of administration is absorption over some portion or all of the alimentary tract and a need for efficient mucosal penetration of the nucleic acid(s) so administered.

35 [0518] Delivery of a drug via the oral mucosa, as in the case of buccal and sublingual administration, has several desirable features, including, in many instances, a more rapid rise in plasma concentration of the drug than via oral delivery (Harvey, Chapter 35 In: Remington's Pharmaceutical Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, page 711).

40 [0519] Endoscopy may be used for drug delivery directly to an interior portion of the alimentary tract. For example, endoscopic retrograde cystopancreatography (ERCP) takes advantage of extended gastroscopy and permits selective access to the biliary tract and the pancreatic duct (Hirahata et al., Gan To Kagaku Ryoho, 1992, 19(10 Suppl.), 1591). Pharmaceutical compositions, including liposomal formulations, can be delivered directly into portions of the alimentary canal, such as, e.g., the duodenum (Somogyi et al., Pharm. Res., 1995, 12, 149) or the gastric submucosa (Akamo et al., Japanese J. Cancer Res., 1994, 85, 652) via endoscopic means. Gastric lavage devices (Inoue et al., Artif. Organs, 1997, 21, 28) and percutaneous endoscopic feeding devices (Pennington et al., Ailment Pharmacol. Ther., 1995, 9, 471) can also be used for direct alimentary delivery of pharmaceutical compositions.

45 [0520] In some embodiments, oligonucleotide formulations may be administered through the anus into the rectum or lower intestine. Rectal suppositories, retention enemas or rectal catheters can be used for this purpose and may be preferred when patient compliance might otherwise be difficult to achieve (e.g., in pediatric and geriatric applications, or when the patient is vomiting or unconscious). Rectal administration can result in more prompt and higher blood levels

than the oral route. (Harvey, Chapter 35 In: Remington=s Pharmaceutical Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, page 711). Because about 50% of the drug that is absorbed from the rectum will bypass the liver, administration by this route significantly reduces the potential for first-pass metabolism (Benet et al., Chapter 1 In: Goodman & Gilman=s The Pharmacological Basis of Therapeutics, 9th Ed., Hardman et al., eds., McGraw-Hill, New York, NY, 1996).

**[0521]** One advantageous method of non-parenteral administration oligonucleotide compositions is oral delivery. Some embodiments employ various penetration enhancers in order to effect transport of oligonucleotides and other nucleic acids across mucosal and epithelial membranes. Penetration enhancers may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p. 92). Accordingly, some embodiments comprise oral oligonucleotide compositions comprising at least one member of the group consisting of surfactants, fatty acids, bile salts, chelating agents, and non-chelating surfactants. Further embodiments comprise oral oligonucleotide comprising at least one fatty acid, e.g. capric or lauric acid, or combinations or salts thereof. Other embodiments comprise methods of enhancing the oral bioavailability of an oligonucleotide, the method comprising co-administering the oligonucleotide and at least one penetration enhancer.

**[0522]** Other excipients that may be added to oral oligonucleotide compositions include surfactants (or "surface-active agents"), which are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the alimentary mucosa and other epithelial membranes is enhanced. In addition to bile salts and fatty acids, surfactants include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92); and perfluorohemical emulsions, such as FC-43 (Takahashi et al., J. Pharm. Pharmacol., 1988, 40, 252).

**[0523]** Fatty acids and their derivatives which act as penetration enhancers and may be used in compositions of the present invention include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines and mono- and di-glycerides thereof and/or physiologically acceptable salts thereof (*i.e.*, oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, *etc.*) (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1; El-Hariri et al., J. Pharm. Pharmacol., 1992, 44, 651).

**[0524]** In some embodiments, oligonucleotide compositions for oral delivery comprise at least two discrete phases, which phases may comprise particles, capsules, gel-capsules, microspheres, etc. Each phase may contain one or more oligonucleotides, penetration enhancers, surfactants, bioadhesives, effervescent agents, or other adjuvant, excipient or diluent. In some embodiments, one phase comprises at least one oligonucleotide and at least one penetration enhancer. In some embodiments, a first phase comprises at least one oligonucleotide and at least one penetration enhancer, while a second phase comprises at least one penetration enhancer. In some embodiments, a first phase comprises at least one oligonucleotide and at least one penetration enhancer, while a second phase comprises at least one penetration enhancer and substantially no oligonucleotide. In some embodiments, at least one phase is compounded with at least one degradation retardant, such as a coating or a matrix, which delays release of the contents of that phase. In some embodiments, at least one phase In some embodiments, a first phase comprises at least one oligonucleotide, at least one penetration enhancer, while a second phase comprises at least one penetration enhancer and a release-retardant. In particular embodiments, an oral oligonucleotide comprises a first phase comprising particles containing an oligonucleotide and a penetration enhancer, and a second phase comprising particles coated with a release-retarding agent and containing penetration enhancer.

**[0525]** A variety of bile salts also function as penetration enhancers to facilitate the uptake and bioavailability of drugs. The physiological roles of bile include the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 In: Goodman & Gilman=s The Pharmacological Basis of Therapeutics, 9th Ed., Hardman et al., eds., McGraw-Hill, New York, NY, 1996, pages 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus, the term "bile salt" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (CDCA, sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Swinyard, Chapter 39 In: Remington=s Pharmaceutical Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1; Yamamoto et al., J. Pharm. Exp. Ther., 1992, 263, 25; Yamashita et al., J. Pharm. Sci., 1990, 79, 579).

**[0526]** In some embodiments, penetration enhancers useful in some embodiments of present invention are mixtures of penetration enhancing compounds. One such penetration enhancer is a mixture of UDCA (and/or CDCA) with capric and/or lauric acids or salts thereof e.g. sodium. Such mixtures are useful for enhancing the delivery of biologically active substances across mucosal membranes, in particular intestinal mucosa. Other penetration enhancer mixtures comprise about 5-95% of bile acid or salt(s) UDCA and/or CDCA with 5-95% capric and/or lauric acid. Particular penetration enhancers are mixtures of the sodium salts of UDCA, capric acid and lauric acid in a ratio of about 1:2:2 respectively. Another such penetration enhancer is a mixture of capric and lauric acid (or salts thereof) in a 0.01:1 to 1:0.01 ratio (mole basis). In particular embodiments capric acid and lauric acid are present in molar ratios of e.g. about 0.1:1 to about 1:0.1, in particular about 0.5:1 to about 1:0.5.

**[0527]** Other excipients include chelating agents, i.e. compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the alimentary and other mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, J. Chromatogr., 1993, 618, 315). Chelating agents of the invention include, but are not limited to, disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), *N*-acyl derivatives of collagen, laurith-9 and *N*-amino acyl derivatives of beta-diketones (enamines)(Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1; Buur et al., J. Control Rel., 1990, 14, 43).

**[0528]** As used herein, non-chelating non-surfactant penetration enhancers may be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary and other mucosal membranes (Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1). This class of penetration enhancers includes, but is not limited to, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., J. Pharm. Pharmacol., 1987, 39, 621).

**[0529]** Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), can be used.

**[0530]** Some oral oligonucleotide compositions also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which may be inert (*i.e.*, does not possess biological activity *per se*) or may be necessary for transport, recognition or pathway activation or mediation, or is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., Antisense Res. Dev., 1995, 5, 115; Takakura et al., Antisense & Nucl. Acid Drug Dev., 1996, 6, 177).

**[0531]** A "pharmaceutical carrier" or "excipient" may be a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, *etc.*, when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, *etc.*); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, *etc.*); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, *etc.*); disintegrants (e.g., starch, sodium starch glycolate, EXPLOTAB); and wetting agents (e.g., sodium lauryl sulphate, *etc.*).

**[0532]** Oral oligonucleotide compositions may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the composition of present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. Further embodiments of the invention are described below:

## EP 2 336 318 A1

1. An antisense compound 8 to 50 nucleobases in length, wherein said compound specifically hybridizes with nucleotides 2920-3420 as set forth in SEQ ID NO:3 and inhibits expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.
- 5 2. The antisense compound of embodiment 1, wherein said compound specifically hybridizes with nucleotides 3230-3288 as set forth in SEQ ID NO:3 and inhibits expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.
- 10 3. The antisense compound of embodiment 2 that is an antisense oligonucleotide.
4. The antisense compound of embodiment 3, wherein the antisense oligonucleotide is an oligonucleotide mimetic compound.
- 15 5. The antisense compound of embodiment 2, twelve to thirty nucleobases in length.
6. The antisense compound of embodiment 5, fourteen to twenty nucleobases in length.
- 20 7. The antisense compound of embodiment 4, wherein the oligonucleotide mimetic compound comprises at least one phosphorothioate linkage.
8. The antisense compound of embodiment 4, wherein the oligonucleotide mimetic compound comprises at least one 2'-O-methoxyethyl sugar moiety.
- 25 9. The antisense compound of embodiment 4, wherein the oligonucleotide mimetic compound comprises at least one 5-methylcytosine.
10. The antisense compound of embodiment 2, wherein the antisense compound is a chimeric antisense compound.
- 30 11. The antisense compound of embodiment 10, wherein the chimeric antisense compound is a chimeric phosphorothioate antisense compound.
12. The antisense compound of embodiment 11, wherein the chimeric phosphorothioate antisense compound comprises 2'-methoxyethoxyl nucleotide wings and a 2'-deoxynucleotide gap.
- 35 13. The antisense compound of embodiment 12, wherein the chimeric phosphorothioate antisense compound comprises ten 2'-deoxynucleotides.
14. The antisense compound of any one of embodiments 1-13, wherein said antisense compound inhibits expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 50% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.
- 40 15. The antisense compound of any one of embodiments 1-13, wherein at least one nucleobase is covalently linked to a conjugate.
- 45 16. A composition comprising the antisense compound of any one of embodiments 1-13 and a pharmaceutically acceptable carrier or diluent.
17. The composition of embodiment 16 further comprising a colloidal dispersion system.
- 50 18. A composition comprising an antisense compound of any of embodiments 1-13 hybridized to a complementary strand.
19. The composition of embodiment 18, wherein the hybridization of the antisense compound to the complementary strand forms at least one blunt end.
- 55 20. The composition of embodiment 19, wherein the hybridization of the antisense compound to the complementary strand forms two blunt ends.

21. An antisense oligonucleotide compound 8 to 50 nucleobases in length comprising at least 8 contiguous nucleotides of SEQ ID NO:247.

5 22. The antisense oligonucleotide compound of embodiment 21, wherein the antisense oligonucleotide compound has a sequence comprising SEQ ID NO:247.

23. The antisense oligonucleotide compound of embodiment 22, twelve to thirty nucleobases in length.

10 24. The antisense oligonucleotide compound of embodiment 23, fourteen to twenty nucleobases in length.

25. The antisense oligonucleotide compound of embodiment 24, wherein the antisense oligonucleotide compound has a sequence consisting of SEQ ID NO:247.

15 26. The antisense oligonucleotide compound of embodiment 25, wherein the antisense oligonucleotide compound is an oligonucleotide mimetic compound.

27. The antisense oligonucleotide compound of embodiment 26, wherein the oligonucleotide mimetic compound is a chimeric phosphorothioate oligonucleotide compound.

20 28. The antisense oligonucleotide compound of embodiment 27, wherein the chimeric phosphorothioate oligonucleotide compound comprises 2'-methoxyethoxyl nucleotide wings and a 2'-deoxynucleotide gap.

25 29. The antisense oligonucleotide compound of embodiment 28, wherein the chimeric phosphorothioate oligonucleotide compound comprises ten 2'-deoxynucleotides.

30 30. The antisense oligonucleotide compound of any one of embodiments 21-29, wherein at least one oligonucleotide is covalently linked to a conjugate.

31. A composition comprising the antisense oligonucleotide compound of any of embodiments 21-29 and a pharmaceutically acceptable carrier or diluent.

32. The composition of embodiment 31 further comprising a colloidal dispersion system.

35 33. A composition comprising an oligonucleotide compound of any of embodiments 22-29 hybridized to a complementary strand.

34. The composition of embodiment 33, wherein the hybridization of the oligonucleotide compound to the complementary strand forms at least one blunt end.

40 35. The composition of embodiment 34, wherein the hybridization of the oligonucleotide compound to the complementary strand forms two blunt ends.

45 36. A method of inhibiting the expression of apolipoprotein B in cells or tissues comprising contacting said cells or tissues with a compound of embodiment 2 under conditions such that expression of apolipoprotein B is inhibited.

37. A method of inhibiting the expression of apolipoprotein B in cells or tissues comprising contacting said cells or tissues with a compound of embodiment 21 under conditions such that expression of apolipoprotein B is inhibited.

50 38. The method of embodiment 36 or embodiment 37, wherein the cells or tissues are contacted *in vivo*.

39. The method of embodiment 38, wherein said contacting comprises the step of administering the compound to an animal.

55 40. The method of embodiment 39, wherein the animal is a human.

41. The method of embodiment 40, wherein the human has a disease or condition associated with apolipoprotein B expression and a therapeutically or prophylactically effective amount of the compound is administered.



42. The method of embodiment 41, wherein the human has a condition associated with abnormal lipid metabolism.
43. The method of embodiment 41, wherein the human has a condition associated with abnormal cholesterol metabolism.
- 5 44. The method of embodiment 41, wherein the human has a cardiovascular disease.
45. The method of embodiment 44, wherein the cardiovascular disease is atherosclerosis.
- 10 46. The method of embodiment 41, wherein the human has an abnormal metabolic condition associated with apolipoprotein B expression.
47. The method of embodiment 46, wherein the abnormal metabolic condition is hyperlipidemia.
- 15 48. The method of embodiment 41, wherein the human has diabetes.
49. The method of embodiment 41, wherein the human is obese.
- 20 50. The method of embodiment 40, wherein an effective amount of the compound is administered to prevent a disease or condition associated with apolipoprotein B expression.
51. The method of embodiment 40, wherein an effective amount of the compound is administered to delay a disease or condition associated with apolipoprotein B expression.
- 25 52. A method of preventing or delaying the onset of an increase in glucose levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of embodiment 1.
53. A method of preventing or delaying the onset of an increase in glucose levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of embodiment 22.
- 30 54. The method of embodiment 52 or embodiment 53 wherein the animal is a human.
55. The method of embodiment 54 wherein the glucose levels are serum or plasma glucose levels.
- 35 56. A method of modulating serum cholesterol levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of embodiment 1 or 21.
57. The method of embodiment 56 wherein the animal is a human.
- 40 58. A method of modulating lipoprotein levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of embodiment 1.
59. A method of modulating lipoprotein levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of embodiment 22.
- 45 60. The method of embodiment 58 or embodiment 59 wherein the animal is a human.
61. The method of embodiment 60 wherein the lipoprotein is VLDL.
- 50 62. The method of embodiment 60 wherein the lipoprotein is HDL.
63. The method of embodiment 60 wherein the lipoprotein is LDL.
- 55 64. The method of any one of embodiments 39, 52, 53, 56, 58, and 59 wherein the compound is administered intravenously.
65. The method of any one of embodiments 39, 52, 53, 56, 58, and 59 wherein the compound is administered subcutaneously.

## EP 2 336 318 A1

5 66. An antisense oligonucleotide compound 20 nucleobases in length having a sequence of nucleobases as set forth in SEQ ID NO:247 and comprising 5-methylcytidine at nucleobases 2, 3, 5, 9, 12, 15, 17, 19, and 20, wherein every internucleoside linkage is a phosphothioate linkage, nucleobases 1-5 and 16-20 comprise a 2'-methoxyethoxyl modification, and nucleobases 6-15 are deoxynucleotides.

67. The antisense oligonucleotide compound of embodiment 66, wherein at least one oligonucleotide is covalently linked to a conjugate.

10 68. A composition comprising the antisense oligonucleotide compound of embodiment 66 and a pharmaceutically acceptable carrier or diluent.

69. The composition of embodiment 68 further comprising a colloidal dispersion system.

15 70. A composition comprising the antisense oligonucleotide compound of embodiment 66 hybridized to a complementary strand.

71. A method of inhibiting the expression of apolipoprotein B in cells or tissues comprising contacting said cells or tissues with a compound of embodiment 66 so that expression of apolipoprotein B is inhibited.

20 72. The method of embodiment 71, wherein the cells or tissues are contacted *in vivo*.

73. The method of embodiment 72, wherein said contacting comprises the step of administering the compound to an animal.

25 74. The method of embodiment 73, wherein the animal is a human.

75. The method of embodiment 74, wherein the human has a disease or condition associated with apolipoprotein B expression and a therapeutically or prophylactically effective amount of the compound is administered.

30 76. The method of embodiment 75, wherein the human has a condition associated with abnormal lipid metabolism.

77. The method of embodiment 75, wherein the human has a condition associated with abnormal cholesterol metabolism.

35 78. The method of embodiment 75, wherein the human has a cardiovascular disease.

79. The method of embodiment 78, wherein the cardiovascular disease is atherosclerosis.

40 80. The method of embodiment 75, wherein the human has an abnormal metabolic condition associated with apolipoprotein B expression.

81. The method of embodiment 80, wherein the abnormal metabolic condition is hyperlipidemia.

45 82. The method of embodiment 75, wherein the human has diabetes.

83. The method of embodiment 75, wherein the human is obese.

50 84. The method of embodiment 74, wherein an effective amount of the compound is administered to prevent a disease or condition associated with apolipoprotein B expression.

85. The method of embodiment 74, wherein an effective amount of the compound is administered to delay a disease or condition associated with apolipoprotein B expression.

55 86. A method of preventing or delaying the onset of an increase in glucose levels in a human comprising administering to said human a therapeutically or prophylactically effective amount of the compound of embodiment 66.

87. The method of embodiment 86 wherein the glucose levels are serum glucose levels.

88. The method of embodiment 86 wherein the glucose levels are plasma glucose levels.

89. A method of modulating serum cholesterol levels in a human comprising administering to said human a therapeutically or prophylactically effective amount of the compound of embodiment 66.

5

90. A method of modulating lipoprotein levels in a human comprising administering to said human a therapeutically or prophylactically effective amount of the compound of embodiment 66.

10

91. The method of embodiment 90 wherein the lipoprotein is VLDL.

92. The method of embodiment 90 wherein the lipoprotein is HDL.

93. The method of embodiment 90 wherein the lipoprotein is LDL.

15

94. The method of any one of embodiments 73-93 wherein the compound is administered intravenously.

95. The method of any one of embodiments 73-93 wherein the compound is administered subcutaneously.

20

96. The method of any one of embodiments 73-93 wherein the compound is administered orally.

97. A kit comprising a compound of any one of embodiments 1-15, 21-30, and 66-67.

25

98. A compound comprising a first nucleobase strand hybridized to a second nucleobase strand, each strand 8 to 50 nucleobases in length, said first nucleobase strand comprising a sequence of at least 8 contiguous nucleobases of the sequence set forth in SEQ ID NO:3, said second nucleobase strand comprising a sequence sufficiently complementary to said first strand so as to permit stable hybridization, said compound inhibiting expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 100 nM.

30

99. The compound of embodiment 98, wherein the first strand comprises a sequence of 12 to 30 contiguous nucleobases of the sequence set forth in SEQ ID NO:3.

100. The compound of embodiment 98, wherein the first strand comprises a sequence of 20 contiguous nucleobases of the sequence set forth in SEQ ID NO:3.

35

101. The compound of embodiments 98, 99, or 100, wherein the second strand comprises a sequence perfectly complementary to at least 8 contiguous nucleobases of the sequence set forth in SEQ ID NO:3.

40

102. The compound of embodiment 101, wherein the second strand comprises a sequence perfectly complementary to 12 to 30 nucleobases of the sequence set forth in SEQ ID NO:3.

103. The compound of embodiment 101, wherein the second strand comprises a sequence perfectly complementary to 20 nucleobases of the sequence set forth in SEQ ID NO:3.

45

104. The compound of any of embodiments 98-103, wherein at least one strand comprises RNA.

105. The compound of any of embodiments 98-104, wherein at least one strand comprises one or more deoxynucleosides.

50

106. The compound of any of embodiments 98-105, wherein the hybridized strands form at least one overhanging end.

107. The compound of embodiment 106, wherein the overhanging end comprises at least one modified base.

55

108. The compound of any of embodiments 98-107, wherein said compound inhibits expression of mRNA encoding human apolipoprotein B after 16 to 24 hours by at least 50% in 80% confluent HepG2 cells in culture at a concentration of 100 nM.

## EP 2 336 318 A1

109. A vesicle comprising a compound any of embodiments 98-108.

110. The vesicle of embodiment 109, wherein the vesicle is a liposome.

5 111. A composition comprising the compound of any one of embodiments 98-108 and a pharmaceutically acceptable carrier or diluent.

112. The composition of embodiment 111 further comprising a colloidal dispersion system.

10 113. A method of inhibiting the expression of apolipoprotein B in cells or tissues comprising contacting said cells or tissues with the compound of any one of embodiments 98-108 under conditions such that expression of apolipoprotein B is inhibited.

15 114. The method of embodiment 113, wherein the cells or tissues are contacted *in vivo*.

115. The method of embodiment 114, wherein said contacting comprises the step of administering the compound to an animal.

20 116. The method of embodiment 115, wherein the animal is a human.

117. The method of embodiment 116, wherein the human has a condition associated with apolipoprotein B expression and a therapeutically or prophylactically effective amount of the compound is administered.

25 118. The method of embodiment 117, wherein said condition is associated with abnormal lipid metabolism.

119. The method of embodiment 117, wherein said condition is associated with abnormal cholesterol metabolism.

120. The method of embodiment 117, wherein said condition is cardiovascular disease.

30 121. The method of embodiment 120, wherein the cardiovascular disease is atherosclerosis.

122. The method of embodiment 117, wherein said condition is an abnormal metabolic condition associated with apolipoprotein B expression.

35 123. The method of embodiment 122, wherein the abnormal metabolic condition associated with apolipoprotein B expression is hyperlipidemia.

124. The method of embodiment 117, wherein the condition is diabetes.

40 125. The method of embodiment 117, wherein the condition is obesity.

126. The method of embodiment 116, wherein an effective amount of the compound is administered to prevent a condition associated with apolipoprotein B expression.

45 127. The method of embodiment 126, wherein an effective amount of the compound is administered to delay a condition associated with apolipoprotein B expression.

128. A method of reducing lipoprotein(a) secretion by hepatocytes comprising:

50 (a) contacting hepatocytes with an amount of a composition comprising a non-catalytic compound 8 to 50 nucleobases in length that specifically hybridizes with mRNA encoding human apolipoprotein B and inhibits expression of the mRNA after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM, wherein said amount is effective to inhibit expression of apolipoprotein B in the hepatocytes; and

55 (b) measuring lipoprotein(a) secretion by the hepatocytes.

129. The method of embodiment 128, wherein the non-catalytic compound specifically hybridizes with nucleotides

## EP 2 336 318 A1

3230-3288 as set forth in SEQ ID NO:3.

130. The method of embodiment 129, wherein the non-catalytic compound comprises a sequence of nucleobases as set forth in SEQ ID NO:247.

5

131. The method of any of embodiments 128-130, wherein the non-catalytic compound is an antisense oligonucleotide mimetic.

10

132. A method of treating a condition associated with apolipoprotein B expression in a primate comprising administering to the primate a therapeutically or prophylactically effective amount of a non-catalytic compound 8 to 50 nucleobases in length that specifically hybridizes with mRNA encoding human apolipoprotein B and inhibits expression of the mRNA after 16 to 24 hours by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.

15

133. The method of embodiment 132, wherein the primate is a human.

134. The method of embodiment 133 wherein the condition is selected from the group consisting of abnormal lipid metabolism, abnormal cholesterol metabolism, cardiovascular disease, hyperlipidemia, diabetes, and obesity.

20

135. A method of reducing apolipoprotein B expression in the liver of an animal, comprising administering to the animal between 2 mg/kg and 20 mg/kg of a non-catalytic compound 8 to 50 nucleobases in length that specifically hybridizes with mRNA encoding human apolipoprotein B by at least 30% in 80% confluent HepG2 cells in culture at a concentration of 150 nM.

25

136. The method of embodiment 135, wherein the compound is administered subcutaneously.

137. The method of embodiment 135, wherein the compound is administered intravenously.

30

138. The method of embodiment 135, wherein the compound is administered orally.

139. The method of any one of embodiments 135-138, wherein administration to the animal is repeated.

140. The method of any one of embodiments 135-139, wherein the animal is a human.

35

141. Use of a compound of any one of embodiments 1-15, 21-30, 66-67, and 98-108 in the production of a medicament.

142. Use of a compound of any one of embodiments 1-15, 21-30, 66-67, and 98-108 in a medicament for altering lipid metabolism.

40

143. Use of a compound of any one of embodiments 1-15, 21-30, 66-67, and 98-108 in a medicament for a disease or condition associated with apolipoprotein B expression.

45

144. A method of making a compound of any one of embodiments 98-108 comprising specifically hybridizing *in vitro* a first nucleobase strand comprising a sequence of at least 8 contiguous nucleobases of the sequence set forth in SEQ ID NO:3 to a second nucleobase strand comprising a sequence sufficiently complementary to said first strand so as to permit stable hybridization.

50

55

EP 2 336 318 A1

SEQUENCE LISTING

<110> Genzyme Corporation

5 <120> ANTISENSE MODULATION OF APOLIPOPROTEIN B EXPRESSION

<130> P055617EP

<150> US 60/426,234

<151> 2002-11-13

10 <150> PCT/US03/015493

<151> 2003-05-15

<150> PCT/US03/036411

<151> 2003-11-13

15 <150> EP 03789763.4

<151> 2003-11-13

<160> 892

<210> 1

<211> 20

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctccctcaggg 20

25 <210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

30 <400> 2

atgcattctg cccccaagga 20

<210> 3

<211> 14121

<212> DNA

35 <213> Homo sapiens

<220>

<221> CDS

<222> (129)..(13820)

<400> 3

40 atccccaccg ggacctgcgg ggctgagtgc ctttctcggg tgctgccgct gaggagcccg 60

cccagccagc cagggccggg aggccgaggc cagggccgag cccaggagcc gccccaaccg 120

agctggcg atg gac ccg ccg agg ccc gcg ctg ctg gcg ctg ctg gcg ctg 170

Met Asp Pro Pro Arg Pro Ala Leu Leu Ala Leu Leu Ala Leu

1 5 10

45 cct gcg ctg ctg ctg ctg ctg gcg ggc gcc agg gcc gaa gag gaa 218

Pro Ala Leu Leu Leu Leu Leu Ala Gly Ala Arg Ala Glu Glu Glu

15 20 25 30

atg ctg gaa aat gtc agc ctg gtc tct cca aaa gat gcg acc cga ttc 266

Met Leu Glu Asn Val Ser Leu Val Cys Pro Lys Asp Ala Thr Arg Pro

35 40 45

50 aag cac ctg cgg aag cac aca cac aac tal gag gct gag agt tcc agc 314

Lys His Leu Arg Lys Tyr Thr Tyr Asn Tyr Glu Ala Glu Ser Ser Ser

EP 2 336 318 A1

			50					55				60		
			gga gtc cct ggg act gct gal lca aga agl gcc acc agg atc aac tgc											362
			Gly Val Pro Gly Thr Ala Asp Ser Arg Ser Ala Thr Arg Ile Asn Cys											
5			65					70				75		
			aaq qtt gag ctg gaq gtt ccc caq ctc tgc agc ttc atc ctg aag acc											410
			Lys Val Glu Leu Glu Val Pro Gln Leu Cys Ser Phe Ile Leu Lys Thr											
			80					85				90		
10			agc cag tgc acc ctg aaa gag gtg tat ggc ttc aac cct gag ggc aaa											458
			Ser Gln Cys Thr Leu Lys Glu Val Tyr Gly Phe Asn Pro Glu Gly Lys											
			95					100				105		110
			gcc ttg ctg aag aaa acc aag aac tat gag gag ttt gct gca gcc atg											506
			Ala Leu Leu Lys Lys Thr Lys Asn Ser Glu Glu Phe Ala Ala Ala Met											
			115					120				125		
15			lcc agg lal gag ctg aag ctg gcc al_ cca gaa ggg aag cag gll llc											554
			Ser Arg Tyr Glu Leu Lys Leu Ala Ile Pro Glu Gly Lys Gln Val Phe											
			130					135				140		
			ctt tac cag gag aaa gat gaa cct act tac atc ctg aac atc aag agg											602
			Leu Tyr Pro Glu Lys Asp Glu Pro Thr Tyr Ile Leu Asn Ile Lys Arg											
			145					150				155		
20			ggc atc att tct gcc ctg gtt ccc cca gag aca gaa gaa gcc aag											650
			Gly Ile Ile Ser Ala Leu Leu Val Pro Pro Glu Thr Glu Glu Ala Lys											
			160					165				170		
			caa gtg ttg ttt ctg gat acc gtg tat gga aac tgc tcc act cac ttt											698
			Gln Val Leu Phe Leu Asp Thr Val Tyr Gly Asn Cys Ser Thr His Phe											
			175					180				185		190
25			acc gtc aag acg agg aag ggc aat gtg gca aca gaa ata tcc act gaa											746
			Thr Val Lys Thr Arg Lys Gly Asn Val Ala Thr Glu Ile Ser Thr Glu											
			195					200				205		
			aga gac ctg ggg cag tgt gat cgc ttc aag ccc atc cgc aca ggc atc											794
			Arg Asp Leu Gly Gln Cys Asp Arg Phe Lys Pro Ile Arg Thr Gly Ile											
			210					215				220		
30			agc cca ctt gct ctg atc aaa ggc atg acc cgc ccc ttg tca act ctg											842
			Ser Pro Leu Ala Leu Ile Lys Gly Met Thr Arg Pro Leu Ser Thr Leu											
			225					230				235		
			atc agc agc agc cag tcc tgt cag tac aca ctg gac gct aag agg aag											890
			Ile Ser Ser Ser Gln Ser Cys Gln Tyr Thr Leu Asp Ala Lys Arg Lys											
			240					245				250		
35			cat gtg gca gaa gcc atc tgc aag gag caa cac ctg ttc ctg cct ttc											938
			His Val Ala Glu Ala Ile Cys Lys Glu Gln His Leu Phe Leu Pro Phe											
			255					260				265		270
			tcc tac aac aat aag tat ggg atg gta gca caa gtg aca cag act ttg											986
			Ser Tyr Asn Asn Lys Tyr Gly Met Val Ala Gln Val Thr Gln Thr Leu											
			275					280				285		
			aaa ctt gaa gac aca cca aag atc aac agc cgc ttc ttt ggt gaa ggt											1034
			Lys Leu Glu Asp Thr Pro Lys Ile Asn Ser Arg Phe Phe Gly Glu Gly											
			290					295				300		
45			act aag aag atg ggc ctg gca ttt gag agc acc aaa tcc aca tca cct											1082
			Thr Lys Lys Met Gly Leu Ala Phe Glu Ser Thr Lys Ser Thr Ser Pro											
			305					310				315		
			cca aaq caq gcc gaa gct gtt ttg aaq act ctg caq gaa ctg aaa asa											1130
			Pro Lys Gln Ala Glu Ala Val Leu Lys Thr Leu Gln Glu Leu Lys Lys											
			320					325				330		
50			cta acc atc tct gag caa aat atc cag aga gct aat ctg ttc aat aag											1178
			Leu Thr Ile Ser Glu Gln Asn Ile Gln Arg Ala Asn Leu Phe Asn Lys											
			335					340				345		350
55														

EP 2 336 318 A1

	ctg gtt act gag ctg aga ggc ctc agt gat gaa gca gtc aca tct ctc	1226
	Leu Val Thr Glu Leu Arg Gly Leu Ser Asp Glu Ala Val Thr Ser Leu	
	355 360 365	
5	ttg cca cag ctg att gag gtg tcc agc ccc atc act tta caa gcc ttg	1274
	Leu Pro Gln Leu Ile Glu Val Ser Ser Pro Ile Thr Leu Gln Ala Leu	
	370 375 380	
	ggt cag tgt gga cag cct cag tgc tcc act cac atc ctc cag tgg ctg	1322
	Val Gln Cys Gly Gln Pro Gln Cys Ser Thr His Ile Leu Gln Trp Leu	
	385 390 395	
10	aaa cgt qtg cat gcc aac ccc ctt ctg ata gat qtg gtc acc tac ctg	1370
	Lys Arg Val His Ala Asn Pro Leu Leu Ile Asp Val Val Thr Tyr Leu	
	400 405 410	
	gtg gcc ctg atc ccc gag ccc tca gca cag cag ctg cga gag atc ttc	1418
	Val Ala Leu Ile Pro Glu Pro Ser Ala Gln Gln Leu Arg Glu Ile Phe	
	415 420 425 430	
15	aac atg gcg agg gat cag cgc agc cga gcc acc ttg tat gcg ctg agc	1466
	Asn Met Ala Arg Asp Gln Arg Ser Arg Ala Thr Leu Tyr Ala Leu Ser	
	435 440 445	
	cac gcg glc aac aac tal cal aag aca aac ccl aca ggg acc cag gag	1514
	His Ala Val Asn Asn Tyr His Lys Trp Asn Pro Thr Gly Thr Gln Glu	
	450 455 460	
20	ctg ctg gac att gct aat tac ctg atg gaa cag att caa gat gac tgc	1562
	Leu Leu Asp Ile Ala Asn Tyr Leu Met Glu Gln Ile Gln Asp Asp Cys	
	465 470 475	
25	act ggg gat gaa gat tac acc tat ttg att ctg cgg gtc att gga aat	1610
	Thr Gly Asp Glu Asp Tyr Thr Tyr Leu Ile Leu Arg Val Ile Gly Asn	
	480 485 490	
	atg gcc csa acc atg gag cag tta act cca gas ctc aag tct tca atc	1658
	Met Gly Gln Thr Met Glu Gln Leu Thr Pro Glu Leu Lys Ser Ser Ile	
	495 500 505 510	
30	ctc aaa tgt gtc caa agt aca aag cca tca ctg atg atc cag aaa gct	1706
	Leu Lys Cys Val Gln Ser Thr Lys Pro Ser Leu Met Ile Gln Lys Ala	
	515 520 525	
	gcc atc cag gct ctg cgg aaa atg gag cct aaa gac aag gac cag gag	1754
	Ala Ile Gln Ala Leu Arg Lys Met Glu Pro Lys Asp Lys Asp Gln Glu	
	530 535 540	
35	ggt ctt ctt cag act ttc ctt gat gat gct tct ccg gga gat aag cga	1802
	Val Leu Leu Gln Thr Phe Leu Asp Asp Ala Ser Pro Gly Asp Lys Arg	
	545 550 555	
	ctg gct gcc tat ctt atg ttg atg agg agt cct tca cag gca gat att	1850
	Leu Ala Ala Tyr Leu Met Leu Met Arg Ser Pro Ser Gln Ala Asp Ile	
	560 565 570	
40	aac aaa att gtc caa att cta cca tgg gaa cag aat gag caa gtg aag	1898
	Asn Lys Ile Val Gln Ile Leu Pro Trp Glu Gln Asn Glu Gln Val Lys	
	575 580 585 590	
	aac ttt gtg gct tcc cat att gcc aat atc ttg aac tca gaa gaa ttg	1946
	Asn Phe Val Ala Ser His Ile Ala Asn Ile Leu Asn Ser Glu Glu Leu	
	595 600 605	
	gat atc caa gat ctg aaa aag tta gtg aaa gaa gct ctg aaa gaa tcc	1994
	Asp Ile Gln Asp Leu Lys Lys Leu Val Lys Glu Ala Leu Lys Glu Ser	
	610 615 620	
50	caa ctt cca act gtc atg gac ttc aga aaa ttc tct cgg aac tat caa	2042
	Gln Leu Pro Thr Val Met Asp Phe Arg Lys Phe Ser Arg Asn Tyr Gln	
	625 630 635	

55



EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

ctc tac aaa tct gtt tct ctt cca tca ctt gac cca gcc tca gcc aaa 2090  
 Leu Tyr Lys Ser Val Ser Leu Pro Ser Leu Asp Pro Ala Ser Ala Lys  
 640 645 650

ata gaa ggg aat ctt ata ttt gat cca aat aac tac ctt cct aaa gaa 2138  
 Ile Glu Gly Asn Leu Ile Phe Asp Pro Asn Asn Tyr Leu Pro Lys Glu  
 655 660 665 670

agc atg ctg aaa act acc ctc act gcc ttt gga ttt gct tca gct gac 2186  
 Ser Met Leu Lys Thr Thr Leu Thr Ala Phe Gly Phe Ala Ser Ala Asp  
 675 680 685

ctc atc gag att ggc ctg gaa gga aaa ggc ttt gag cca aca ttg gaa 2234  
 Leu Ile Glu Ile Gly Leu Glu Gly Lys Gly Phe Glu Pro Thr Leu Glu  
 690 695 700

gct ctt ttt ggg aag caa gga ttt ttc cca gac agt gtc aac aaa gct 2282  
 Ala Leu Phe Gly Lys Gln Gly Phe Phe Pro Asp Ser Val Asn Lys Ala  
 705 710 715

ttg tac tgg gtt aat ggt caa gtt cct gat ggt gtc tct aag gtc tta 2330  
 Leu Tyr Trp Val Asn Gly Gln Val Pro Asp Gly Val Ser Lys Val Leu  
 720 725 730

ccg gac cac lll ggc cal acc aaa gal gal aaa cal gag cag gal atg 2378  
 Val Asp His Phe Gly Tyr Thr Lys Asp Asp Lys His Glu Gln Asp Met  
 735 740 745 750

gta aat gga ata atg ctc agt gtt gag aag ctg att aaa gat ttg aaa 2426  
 Val Asn Gly Ile Met Leu Ser Val Glu Lys Leu Ile Lys Asp Leu Lys  
 755 760 765

tcc aaa gaa gtc ccg gaa gcc aga gcc tac ctc cgc atc ttg gga gag 2474  
 Ser Lys Glu Val Pro Glu Ala Arg Ala Tyr Leu Arg Ile Leu Gly Glu  
 770 775 780

gag ctt ggt ttt gcc agt ctc cat gac ctc cag ctc ctg gga aag ctg 2522  
 Glu Leu Gly Phe Ala Ser Leu His Asp Leu Gln Leu Leu Gly Lys Leu  
 785 790 795

ctt ctg atg ggt gcc cgc act ctg cag ggg atc ccc cag atg att gga 2570  
 Leu Leu Met Gly Ala Arg Thr Leu Gln Gly Ile Pro Gln Met Ile Gly  
 800 805 810

gag gtc atc agg aag ggc tca aag aat gac ttt ttt ctt cac tac atc 2618  
 Glu Val Ile Arg Lys Gly Ser Lys Asn Asp Phe Phe Leu His Tyr Ile  
 815 820 825 830

ttc atg gag aat gcc ttt gaa ctc ccc act gga gct gga tta cag ttg 2666  
 Phe Met Glu Asn Ala Phe Glu Leu Pro Thr Gly Ala Gly Leu Gln Leu  
 835 840 845

caa ata tct tca tct gga gtc att gct ccc gga gcc aag gct gga gta 2714  
 Gln Ile Ser Ser Ser Gly Val Ile Ala Pro Gly Ala Lys Ala Gly Val  
 850 855 860

aaa ctg gaa gta gcc aac atg cag gct gaa ctg gtg gca aaa ccc tcc 2762  
 Lys Leu Glu Val Ala Asn Met Gln Ala Glu Leu Val Ala Lys Pro Ser  
 865 870 875

cgg tct gtg gag ttt gtg aca cat atg ggc atc atc att ccg gac ttc 2810  
 Val Ser Val Glu Phe Val Thr Asn Met Gly Ile Ile Ile Pro Asp Phe  
 880 885 890

gct agg agt ggg gtc cag atg aac acc aac ttc ttc cac gag tgg ggt 2858  
 Ala Arg Ser Gly Val Gln Met Asn Thr Asn Phe Phe His Glu Ser Gly  
 895 900 905 910

ctg gag gct cat gtt gcc cta aaa gct ggg aag ctg aag ttt atc att 2906  
 Leu Glu Ala His Val Ala Leu Lys Ala Gly Lys Leu Lys Phe Ile Ile  
 915 920 925

EP 2 336 318 A1

cct tcc cca aag aga cca gtc aag ctg ctc agt gga ggc aac aca tta 2954  
 Pro Ser Pro Lys Arg Pro Val Lys Leu Leu Ser Gly Gly Asn Thr Leu  
 930 935 940

5 cat ttg gtc tct acc acc aaa acg gag gtg atc cca cct ctc att gag 3002  
 His Leu Val Ser Thr Thr Lys Thr Glu Val Ile Pro Pro Leu Ile Glu  
 945 950 955

aac agg cag tcc tgg tca gtt tgc aag caa gtc ttt cct ggc ctg aat 3050  
 Asn Arg Gln Ser Trp Ser Val Cys Lys Gln Val Phe Pro Gly Leu Asn  
 960 965 970

10 tac tgc acc tca ggc gct tac tcc aac gcc agc tcc aca gac tcc gcc 3098  
 Tyr Cys Thr Ser Gly Ala Tyr Ser Asn Ala Ser Ser Thr Asp Ser Ala  
 975 980 985 990

tcc tac tat ccg ctg acc ggg gac acc aga tta gag ctg gaa ctg agg 3146  
 Ser Tyr Tyr Pro Leu Thr Gly Asp Thr Arg Leu Glu Leu Clu Leu Arg  
 995 1000 1005

15 cct aca gga gac att caq caq tat tct gtc aqc gca acc tat caq ctc 3194  
 Pro Thr Gly Glu Ile Glu Gln Tyr Ser Val Ser Ala Thr Tyr Glu Leu  
 1010 1015 1020

cag aga gag gac aga gcc ttg gtg gat acc ctg aag ttt gta act caa 3242  
 Gln Arg Glu Asp Arg Ala Leu Val Asp Thr Leu Lys Phe Val Thr Gln  
 1025 1030 1035

20 gca gaa ggt gcg aag cag act gag gct acc atg aca ttc aaa tat aat 3290  
 Ala Glu Gly Ala Lys Gln Thr Glu Ala Thr Met Thr Phe Lys Tyr Asn  
 1040 1045 1050

cgg cag agt atg acc ctg tcc agt gaa gtc caa att ccg gat ttt gat 3338  
 Arg Gln Ser Met Thr Leu Ser Ser Glu Val Gln Ile Pro Asp Phe Asp  
 1055 1060 1065 1070

ctt gac ctc gga aca atc ctc aga gtc aat gat gaa tct act caq gcc 3386  
 Val Asp Leu Gly Thr Ile Leu Arg Val Asn Asp Glu Ser Thr Glu Gly  
 1075 1080 1085

30 aaa acg tct tac aga ctc acc ctg gac att cag aac aag aaa att act 3434  
 Lys Thr Ser Tyr Arg Leu Thr Leu Asp Ile Gln Asn Lys Lys Ile Thr  
 1090 1095 1100

gag gtc gcc ctc atg ggc cac cta agt tgt gac aca aag gaa gaa aga 3482  
 Glu Val Ala Leu Met Gly His Leu Ser Cys Asp Thr Lys Clu Glu Arg  
 1105 1110 1115

aaa atc aag ggt gtt att tcc ata ccc cgt ttg caa gca gaa gcc aga 3530  
 Lys Ile Lys Gly Val Ile Ser Ile Pro Arg Leu Gln Ala Glu Ala Arg  
 1120 1125 1130

40 agt gag atc ctc gcc cac tgg tgg cct gcc aaa ctg ctt ctc caa atg 3578  
 Ser Glu Ile Leu Ala His Trp Ser Pro Ala Lys Leu Leu Leu Gln Met  
 1135 1140 1145 1150

gac tca tct gct aca gct tat ggc tcc aca gtt tcc aag agg ctg gca 3626  
 Asp Ser Ser Ala Thr Ala Tyr Gly Ser Thr Val Ser Lys Arg Val Ala  
 1155 1160 1165

45 tgg cat tat gat gaa gag aag att gaa ttt gaa tgg aac aca ggc acc 3674  
 Trp His Tyr Asp Glu Glu Lys Ile Glu Phe Glu Trp Asn Thr Gly Thr  
 1170 1175 1180

aat gta gat acc aaa aaa atg act tcc aat ttc cct ctg gat ctc tcc 3722  
 Asn Val Asp Thr Lys Lys Met Thr Ser Asn Phe Pro Val Asp Leu Ser  
 1185 1190 1195

50 gat tat cct aag agc ctg cat atg tat gct aat aga ctc ctg gat cac 3770  
 Asp Tyr Pro Lys Ser Leu His Met Tyr Ala Asn Arg Leu Leu Asp His  
 1200 1205 1210

55

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

aga gtc cct gaa aca gac atg act ttc cgg cac gtg ggt tcc aaa tta 3818  
 Arg Val Pro Glu Thr Asp Met Thr Phe Arg His Val Gly Ser Lys Leu  
 1215 1220 1225 1230  
 ata gtt gca atg agc tca tgg ctt cag aag gca tct ggg agt ctt cct 3866  
 Ile Val Ala Met Ser Ser Trp Leu Gln Lys Ala Ser Gly Ser Leu Pro  
 1235 1240 1245  
 tat acc cag act ttg caa gac cac ctg aat agc ctg aag gag ttc aac 3914  
 Tyr Thr Gln Thr Leu Gln Asp His Leu Asn Ser Leu Lys Glu Phe Asn  
 1250 1255 1260  
 ctg caq aac atg gga ttg cca gac ttc cac atc cca gaa aac ctg ttc 3962  
 Leu Gln Asn Met Gly Leu Pro Asp Phe His Ile Pro Glu Asn Leu Phe  
 1265 1270 1275  
 tta aaa agc gat ggc cgg gtc aaa tat acc ttg aac aag aac agt ttg 4010  
 Leu Lys Ser Asp Gly Arg Val Lys Tyr Thr Leu Asn Lys Asn Ser Leu  
 1280 1285 1290  
 aaa att gag att cct ttg cct ttt ggt ggc aaa tcc tcc aga gat cta 4058  
 Lys Ile Glu Ile Pro Leu Pro Phe Gly Gly Lys Ser Ser Arg Asp Leu  
 1295 1300 1305 1310  
 aag atg tta gag act gll agg aca cca gcc ctg cac ttc aag tct glg 4106  
 Lys Met Leu Glu Thr Val Arg Thr Pro Ala Leu His Phe Lys Ser Val  
 1315 1320 1325  
 gga ttc cat ctg cca tct cga gag ttc caa gtc cct act ttt acc att 4154  
 Gly Phe His Leu Pro Ser Arg Glu Phe Gln Val Pro Thr Phe Thr Ile  
 1330 1335 1340  
 ccc aag ttg tat caa ctg caa gtg cct ctg ctg ggt gtt cta gac ctg 4202  
 Pro Lys Leu Tyr Gln Leu Gln Val Pro Leu Leu Gly Val Leu Asp Leu  
 1345 1350 1355  
 tcc acg aat gtc tac agc aac ttg tac aac tgg tcc gcc tcc tac agt 4250  
 Ser Thr Asn Val Tyr Ser Asn Leu Tyr Asn Trp Ser Ala Ser Tyr Ser  
 1360 1365 1370  
 ggt ggc aac acc agc aca gac cat ttc agc ctt cgg gct cgt tac cac 4298  
 Gly Gly Asn Thr Ser Thr Asp His Phe Ser Leu Arg Ala Arg Tyr His  
 1375 1380 1385 1390  
 atg aag gct gac tct gtg gtt gac ctg ctt tcc tac aat gtg caa gga 4346  
 Met Lys Ala Asp Ser Val Val Asp Leu Leu Ser Tyr Asn Val Gln Gly  
 1395 1400 1405  
 tct gga gaa aca aca tat gac cac aag aat acg ttc aca cta tca tgt 4394  
 Ser Gly Glu Thr Thr Tyr Asp His Lys Asn Thr Phe Thr Leu Ser Cys  
 1410 1415 1420  
 gat ggg tct cta cgc cac aaa ttt cta gat tgg aat atc aaa ttc agt 4442  
 Asp Gly Ser Leu Arg His Lys Phe Leu Asp Ser Asn Ile Lys Phe Ser  
 1425 1430 1435  
 cat gta gaa aaa ctt gga aac aac cca gtc tca aaa ggt tta cta ata 4490  
 His Val Glu Lys Leu Gly Asn Asn Pro Val Ser Lys Gly Leu Leu Ile  
 1440 1445 1450  
 ttc gat gca tct agt tcc tgg gga cca cag atg tct gct tca gtt cat 4538  
 Phe Asp Ala Ser Ser Ser Trp Gly Pro Gln Met Ser Ala Ser Val His  
 1455 1460 1465 1470  
 ttg gac tcc aaa aag aaa cag cat ttg ttt gtc aaa gaa gtc aag att 4586  
 Leu Asp Ser Lys Lys Lys Gln His Leu Phe Val Lys Glu Val Lys Ile  
 1475 1480 1485  
 gat ggg cag ttc aga gtc tct tgg ttc tat gct aaa ggc aca tat ggc 4634  
 Asp Gly Gln Phe Arg Val Ser Ser Phe Tyr Ala Lys Gly Thr Tyr Gly  
 1490 1495 1500

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

ctg tct tgt cag agg gat cct aac acg ggc cgg ctc aat gga gag tcc 4682  
 Leu Ser Cys Gln Arg Asp Pro Asn Thr Gly Arg Leu Asn Cys Glu Ser  
 1505 1510 1515

aac ctg agg ttt aac tcc tcc tac ctc caa ggc acc aac cag ata aca 4730  
 Asn Leu Arg Phe Asn Ser Ser Tyr Leu Gln Gly Thr Asn Gln Ile Thr  
 1520 1525 1530

gga aga tat gaa gat gga acc ctc tcc ctc acc tcc acc tct gat ctg 4778  
 Gly Arg Tyr Glu Asp Gly Thr Leu Ser Leu Thr Ser Thr Ser Asp Leu  
 1535 1540 1545 1550

caa agt ggc atc att aaa aat act gct tcc ctc aag tat gag aac tac 4826  
 Gln Ser Gly Ile Ile Lys Asn Thr Ala Ser Leu Lys Tyr Glu Asn Tyr  
 1555 1560 1565

gag ctg acg lla aaa cct gac acc aac ggg aag lal aag aac lll gcc 4874  
 Glu Leu Thr Leu Lys Ser Asp Thr Asn Gly Lys Tyr Lys Asn Phe Ala  
 1570 1575 1580

act tct aac aag atg gat atg acc ttc tct aag caa aat gca ctg ctg 4922  
 Thr Ser Asn Lys Met Asp Met Thr Phe Ser Lys Gln Asn Ala Leu Leu  
 1585 1590 1595

cgt tct gaa tat cag gct gat tac gag tca ttg agg ttc ttc agc ctg 4970  
 Arg Ser Glu Tyr Gln Ala Asp Tyr Glu Ser Leu Arg Phe Phe Ser Leu  
 1600 1605 1610

ctt tct gga tca cta aat tcc cat ggt ctt gag tta aat gct gac atc 5018  
 Leu Ser Gly Ser Leu Asn Ser His Gly Leu Glu Leu Asn Ala Asp Ile  
 1615 1620 1625 1630

tta ggc act gac aaa att aat agt ggt gct cac aag gcg aca cta agg 5066  
 Leu Gly Thr Asp Gly Ile Asn Ser Gly Ala His Lys Ala Thr Leu Arg  
 1635 1640 1645

att ggc caa gat gga ata tct acc agt gca acg acc aac ttg aag tct 5114  
 Ile Gly Gln Asp Gly Ile Ser Thr Ser Ala Thr Thr Asn Leu Lys Cys  
 1650 1655 1660

agt ctc ctg gtg ctg gag aat gag ctg aat gca gag ctt ggc ctc tct 5162  
 Ser Leu Leu Val Leu Glu Asn Glu Leu Asn Ala Glu Leu Gly Leu Ser  
 1665 1670 1675

ggg gca tct atg aaa tta aca aca aat ggc cgc ttc agg gaa cac aat 5210  
 Gly Ala Ser Met Lys Leu Thr Thr Asn Gly Arg Phe Arg Glu His Asn  
 1680 1685 1690

gca aaa ttc agt ctg gat ggg aaa gcc gcc ctc aca gag cta tca ctg 5258  
 Ala Lys Phe Ser Leu Asp Gly Lys Ala Ala Leu Thr Glu Leu Ser Leu  
 1695 1700 1705 1710

gga agt gct tat cag gcc atg att ctg ggt gtc gac agc aaa aac att 5306  
 Gly Ser Ala Tyr Gln Ala Met Ile Leu Gly Val Asp Ser Lys Asn Ile  
 1715 1720 1725

ttc aac ttc aag gtc agt caa gaa gga ctt aag ctc tca aat gac atg 5354  
 Phe Asn Phe Lys Val Ser Gln Glu Gly Leu Lys Leu Ser Asn Asp Met  
 1730 1735 1740

atg ggc tca tat gct gaa atg aaa ttc gac cac aca aac agt ctg aac 5402  
 Met Gly Ser Tyr Ala Glu Met Lys Phe Asp His Thr Asn Ser Leu Asn  
 1745 1750 1755

att gca ggc tta tca ctg gac ttc tct tca aaa ctt gac aac att tac 5450  
 Ile Ala Gly Leu Ser Leu Asp Phe Ser Ser Lys Leu Asp Asn Ile Tyr  
 1760 1765 1770

agc tct gac aag lll lal aag caa acg gll aal lla cag cta cag ccc 5498  
 Ser Ser Asp Lys Phe Tyr Lys Gln Thr Val Asn Leu Gln Leu Gln Pro

EP 2 336 318 A1

	1775		1780		1785		1790	
	Lal Lcl clg gla act act lla aac agl gac clg aaa Lac aal gcl clg							5546
5	Tyr Ser Leu Val Thr Thr Leu Asn Ser Asp Leu Lys Tyr Asn Ala Leu		1795	1800			1805	
	cat ctc acc aac aat ggg aaa cta ccg cta gaa ccc ctg aag ctg cat							5594
	Asp Leu Thr Asn Asn Gly Lys Leu Arg Leu Glu Pro Leu Lys Leu His	1810		1815			1820	
10	gng gct ggt aac cta aaa gga gcc tac caa aat aat gaa ata aaa cac							5642
	Val Ala Gly Asn Leu Lys Gly Ala Tyr Gln Asn Asn Glu Ile Lys His	1825		1830			1835	
	atc tat gcc atc tct tct gct gcc tta tca gca agc tat aaa gca gac							5690
	Ile Tyr Ala Ile Ser Ser Ala Ala Leu Ser Ala Ser Tyr Lys Ala Asp	1840		1845			1850	
15	act gll gcl aag gll cag ggl glg gag lll agc cal cgg clc aac aca							5738
	Thr Val Ala Lys Val Gln Gly Val Glu Phe Ser His Arg Leu Asn Thr	1855		1860			1865	1870
	gac atc gct ggg ctg gct tca gcc att gac atg agc aca aac tat aat							5786
20	Asp Ile Ala Gly Leu Ala Ser Ala Ile Asp Met Ser Thr Asn Tyr Asn	1875		1880			1885	
	tca gac tca ctg cat ttc agc aat gtc ttc cgt tct gta atg gcc ccg							5834
	Ser Asp Ser Leu His Phe Ser Asn Val Phe Arg Ser Val Met Ala Pro	1890		1895			1900	
	ttt acc atg acc atc gat gca cat aca aat ggc aat ggg aaa ctc gct							5882
25	Phe Thr Met Thr Ile Asp Ala His Thr Asn Gly Asn Gly Lys Leu Ala	1905		1910			1915	
	ctc tgg gga gaa cat act ggg cag ctg tat agc aaa ttc ctg ttg aaa							5930
	Leu Thr Gly Glu His Thr Gly Gln Leu Tyr Ser Lys Phe Leu Leu Lys	1920		1925			1930	
	gca gaa cct ctg gca ttt act ttc tct cat gat tac aaa ggc tcc aca							5978
30	Ala Glu Pro Leu Ala Phe Thr Phe Ser His Asp Tyr Lys Gly Ser Thr	1935		1940			1945	1950
	agt cat cat ctc gtg tct agg aaa agc atc agt gca gct ctt gaa cac							6026
	Ser His His Leu Val Ser Arg Lys Ser Ile Ser Ala Ala Leu Glu His	1955		1960			1965	
	aaa gtc agt gcc ctg ctt act cca gct gag cag aca ggc acc tgg aaa							6074
35	Lys Val Ser Ala Leu Leu Thr Pro Ala Glu Gln Thr Gly Thr Trp Lys	1970		1975			1980	
	ctc aag acc caa ttt aac aac aat gaa tac agc cag gac ttg gat gct							6122
	Leu Lys Thr Gln Phe Asn Asn Asn Glu Tyr Ser Gln Asp Leu Asp Ala	1985		1990			1995	
	tac aac act aaa gat aaa att ggc gtg gag ctt act gga cga act ctg							6170
40	Tyr Asn Thr Lys Asp Lys Ile Gly Val Glu Leu Thr Gly Arg Thr Leu	2000		2005			2010	
	gct gac cta act cta cta gac tcc cca att aaa gtg cca ctt tta ctc							6218
	Ala Asp Leu Thr Leu Leu Asp Ser Pro Ile Lys Val Pro Leu Leu Leu	2015		2020			2025	2030
45	agt gag ccc atc aat atc att gat gct tta gag atg aga gat gcc gtt							6266
	Ser Glu Pro Ile Asn Ile Ile Asp Ala Leu Glu Met Arg Asp Ala Val	2035		2040			2045	
	cag aag ccc caa gaa ttt aca att gtc gct ttt gta aag tat gat aaa							6314
50	Glu Lys Pro Gln Glu Phe Thr Ile Val Ala Phe Val Lys Tyr Asp Lys	2050		2055			2060	
	aac caa gat gtt cac tcc att aac ctc cca ttt ttt gag acc ttg caa							6362
	Asn Gln Asp Val His Ser Ile Asn Leu Pro Phe Phe Glu Thr Leu Gln	2065		2070			2075	

55

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

gaa tat ttt gag agg aat cga caa acc att ata gtt gta gtg gaa aac 6410  
 Glu Tyr Phe Glu Arg Asn Arg Gln Thr Ile Ile Val Val Val Glu Asn  
 2380 2085 2090

gta cag aga aac ctg aag cac atc aat att gat caa ttt gta aga aaa 6458  
 Val Gln Arg Asn Leu Lys His Ile Asn Ile Asp Gln Phe Val Arg Lys  
 2095 2100 2105 2110

tac aga gca gcc ctg gga aaa stc cca cag caa gct aat gat tat ctg 6506  
 Tyr Arg Ala Ala Leu Gly Lys Leu Pro Gln Gln Ala Asn Asp Tyr Leu  
 2115 2120 2125

aat tca ttc aat tgg gag aga caa gtc tca cat gcc aag gag aaa ctg 6554  
 Asn Ser Phe Asn Trp Glu Arg Gln Val Ser His Ala Lys Glu Tyr Leu  
 2130 2135 2140

act gct ctc aca aaa aag tat aga atc aca gaa aat gat ata caa att 6602  
 Thr Ala Leu Thr Lys Lys Tyr Arg Ile Thr Glu Asn Asp Ile Gln Ile  
 2145 2150 2155

gca tta gat gat gcc aaa atc aac ttc aat gaa aaa cta tct caa ctg 6650  
 Ala Leu Asp Asp Ala Lys Ile Asn Phe Asn Glu Lys Leu Ser Gln Leu  
 2160 2165 2170

cag aca tal atg ala caa tll gal cag tal all aaa gal agt tal gal 6698  
 Gln Thr Tyr Met Ile Gln Phe Asp Gln Tyr Ile Lys Asp Ser Tyr Asp  
 2175 2180 2185 2190

tta cat gat ttg aaa ata gct att gct aat att att gat gaa atc att 6746  
 Leu His Asp Leu Tyr Phe Ala Ile Ala Asn Ile Ile Asp Glu Ile Ile  
 2195 2200 2205

gaa aaa tta aaa agt ctt gat gag cac tat cat atc cgt gta aat tta 6794  
 Glu Lys Leu Lys Ser Leu Asp Glu His Tyr His Ile Arg Val Asn Leu  
 2210 2215 2220

gta aaa aca atc cat gat cta cat ttg ttt att gaa aat att gat ttt 6842  
 Val Lys Ser Thr Ile His Asp Leu His Leu Phe Ile Glu Asn Ile Asp Phe  
 2225 2230 2235

aac aaa agt gga agt agt act gca tcc tgg att caa aat gtg gat act 6890  
 Asn Lys Ser Gly Ser Ser Thr Ala Ser Trp Ile Gln Asn Val Asp Thr  
 2240 2245 2250

aag tac caa atc aga atc cag ata caa gaa aaa ctg cag cag ctt aag 6938  
 Lys Tyr Gln Ile Arg Ile Gln Ile Gln Glu Lys Leu Gln Gln Leu Lys  
 2255 2260 2265 2270

aga cac ata cag aat ata gac atc cag cac cta gct gga aag tta aaa 6986  
 Arg His Ile Gln Asn Ile Asp Ile Gln His Leu Ala Gly Lys Leu Lys  
 2275 2280 2285

caa cac att gag gct att gat gtt aga gtg ctt tta gat caa ttg gga 7034  
 Gln His Ile Glu Ala Ile Asp Val Arg Val Leu Leu Asp Gln Leu Gly  
 2290 2295 2300

act aca att tca ttt gaa aga ata aat gat gtt ctt gag cat gtc aaa 7082  
 Thr Thr Ile Ser Phe Glu Arg Ile Asn Asp Val Leu Glu His Val Lys  
 2305 2310 2315

cac ttt gtt ata aat ctt att ggg gat ttt gaa gta gct gag aaa atc 7130  
 His Phe Val Ile Asn Leu Ile Gly Asp Phe Glu Val Ala Glu Lys Ile  
 2320 2325 2330

aat gcc ttc aga gcc aaa gtc cat gag tta atc gag agg tat gaa gta 7178  
 Asn Ala Phe Arg Ala Lys Val His Glu Leu Ile Glu Arg Tyr Glu Val  
 2335 2340 2345 2350

gac caa caa atc cag gtt tta atg gat aaa tta gta gag ttg acc cac 7226  
 Asp Gln Gln Ile Gln Val Leu Met Asp Lys Leu Val Glu Leu Thr His  
 2355 2360 2365

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

caa tac aag ttg aag gag act att cag aag cta agc aat gtc cta caa 7274  
 Gln Tyr Lys Leu Lys Glu Thr Ile Cln Lys Leu Ser Asn Val Leu Gln  
 2370 2375 2380

caa gtt aag ata aaa gat tac ttt gag aaa ttg gtt gga ttt att gat 7322  
 Gln Val Lys Ile Lys Asp Tyr Phe Glu Lys Leu Val Gly Phe Ile Asp  
 2385 2390 2395

gat gct gtg aag aag ctt aat gaa tta tct ttt aaa aca ttc att gaa 7370  
 Asp Ala Val Lys Lys Leu Asn Glu Leu Ser Phe Lys Thr Phe Ile Glu  
 2400 2405 2410

gat gtt aac aaa ttc ctt gac atg ttg ata aag aaa tta aag tca ttt 7418  
 Asp Val Asn Lys Phe Leu Asp Met Leu Ile Lys Lys Leu Lys Ser Phe  
 2415 2420 2425 2430

gal lac cac cag lll gla gal gaa acc aal gac aaa atc cgl gag glg 7466  
 Asp Tyr His Gln Phe Val Asp Glu Thr Asn Asp Lys Ile Arg Glu Val  
 2435 2440 2445

act cag aga ctc aat ggt gaa att cag gct ctg gaa cta cca caa aaa 7514  
 Thr Gln Arg Leu Gly Glu Ile Cln Ala Leu Glu Leu Pro Gln Lys  
 2450 2455 2460

gct gaa gca tta aaa ctg ttt tta gag gaa acc aag gcc aca gtt gca 7562  
 Ala Glu Ala Leu Lys Leu Phe Leu Glu Glu Thr Lys Ala Thr Val Ala  
 2465 2470 2475

gtg tat ctg gaa agc cta cag gac acc aaa ata acc tta atc atc aat 7610  
 Val Tyr Leu Glu Ser Leu Cln Asp Thr Lys Ile Thr Leu Ile Ile Asn  
 2480 2485 2490

tgg tta cag gag gct tta agt tca gca tct ttg gct cac atg aag gcc 7658  
 Trp Leu Gln Glu Ala Leu Ser Ser Ala Ser Leu Ala His Met Lys Ala  
 2495 2500 2505 2510

aaa ttc cga gag act cta gaa gat aca cga gac cga atg tat caa atg 7706  
 Lys Phe Arg Glu Thr Leu Glu Asp Thr Arg Asp Arg Met Tyr Gln Met  
 2515 2520 2525

gac att cag cag gaa ctt caa cga tac ctg tct ctg gta gcc cag gtt 7754  
 Asp Ile Gln Gln Glu Leu Cln Arg Tyr Leu Ser Leu Val Gly Gln Val  
 2530 2535 2540

tat agc aca ctt gtc acc tac att tct gat tgg tgg act ctt gct gct 7802  
 Tyr Ser Thr Leu Val Thr Tyr Ile Ser Asp Trp Trp Thr Leu Ala Ala  
 2545 2550 2555

aag aac ctt act gac ttt gca gag caa tat tct atc caa gat tgg gct 7850  
 Lys Asn Leu Thr Asp Phe Ala Glu Gln Tyr Ser Ile Gln Asp Trp Ala  
 2560 2565 2570

aaa cgt atg aaa gca ttg gta gag caa ggg ttc act gtt cct gaa atc 7898  
 Lys Arg Met Lys Ala Leu Val Glu Gln Gly Phe Thr Val Pro Glu Ile  
 2575 2580 2585 2590

aag acc atc ctt ggg acc atg cct gcc ttt gaa gtc agt ctt cag gct 7946  
 Lys Thr Ile Leu Gly Thr Met Pro Ala Phe Glu Val Ser Leu Gln Ala  
 2595 2600 2605

ctt cag aaa gct acc ttc cag aca cct gat ttt ata gtc ccc cta aca 7994  
 Leu Gln Lys Ala Thr Phe Cln Thr Pro Asp Phe Ile Val Pro Leu Thr  
 2610 2615 2620

cat ttg agg att cca tca gtt caa ata aac ttc aaa gac tta aaa aat 8042  
 Asp Leu Arg Ile Pro Ser Val Gln Ile Asn Phe Lys Asp Leu Lys Asn  
 2625 2630 2635

ata aaa atc cca tcc agg ttt tcc aca cca gaa ttt acc atc ctt aac 8090  
 Ile Lys Ile Pro Ser Arg Phe Ser Thr Pro Glu Phe Thr Ile Leu Asn  
 2640 2645 2650

EP 2 336 318 A1

acc ttc ccc att cct tcc ttt cca att gac ttt gtc gaa atg aaa gta 8138  
 Thr Phe His Ile Pro Ser Phe Thr Ile Asp Phe Val Glu Met Lys Val  
 2655 2660 2665 2670  
 5  
 aag atc atc aga acc att gac cag atg cag aac agt gag ctg cag tgg 8186  
 Lys Ile Ile Arg Thr Ile Asp Gln Met Gln Asn Ser Glu Leu Gln Trp  
 2675 2680 2685  
 ccc gtt cca gat ata tat ctc agg gat ctg aag gtg gag gac att cct 8234  
 Pro Val Pro Asp Ile Tyr Leu Arg Asp Leu Lys Val Glu Asp Ile Pro  
 2690 2695 2700  
 10  
 cta gcg aga atc acc ctg cca gac ttc cgt tta cca gaa atc cca att 8282  
 Ile Ala Arg Ile Thr Leu Pro Asp Phe Arg Ile Pro Glu Ile Ala Ile  
 2705 2710 2715  
 cca gaa ttc ata atc cca act ctc aac ctt aat gat ttt caa gtt cct 8330  
 Pro Glu Phe Ile Ile Pro Thr Leu Asn Leu Asn Asp Phe Gln Val Pro  
 2720 2725 2730  
 15  
 gac ctt cac ata cca gaa ttc cag ctt ccc cac atc tca cac aca att 8378  
 Asp Leu His Ile Pro Glu Phe Gln Leu Pro His Ile Ser His Thr Ile  
 2735 2740 2745 2750  
 20  
 gaa gla cct act lll ggc aag cta lac agl all ctg aaa atc caa lcl 8426  
 Glu Val Pro Thr Phe Gly Lys Leu Tyr Ser Ile Leu Lys Ile Gln Ser  
 2755 2760 2765  
 cct ctt ttc aca tta gat gca aat gct gac ata ggg aat gga acc acc 8474  
 Pro Leu Phe Thr Leu Asp Ala Asn Ala Asp Ile Gly Asn Gly Thr Thr  
 2770 2775 2780  
 25  
 tca gca aac gaa gca ggt atc gca gct tcc atc act gcc aaa gga gag 8522  
 Ser Ala Asn Glu Ala Gly Ile Ala Ala Ser Ile Thr Ala Lys Gly Glu  
 2785 2790 2795  
 tcc aaa tta gaa gtt ctc aat ttt gat ttt caa gca aat gca caa ctc 8570  
 Ser Lys Leu Glu Val Leu Asn Phe Asp Phe Gln Ala Asn Ala Gln Leu  
 2800 2805 2810  
 30  
 tca aac cct aac att aat ccg ctg gct ctg aag gag tca gtg aag ttc 8618  
 Ser Asn Pro Lys Ile Asn Pro Leu Ala Leu Lys Glu Ser Val Lys Phe  
 2815 2820 2825 2830  
 35  
 tcc agc acg tac ctg aga acg gag cat ggg agt gaa atg ctg ttt ttt 8666  
 Ser Ser Lys Tyr Leu Arg Thr Glu His Gly Ser Glu Met Leu Phe Phe  
 2835 2840 2845  
 gga aat gct att gag gga aaa tca aac aca gtg gca agt tta cac aca 8714  
 Gly Asn Ala Ile Glu Gly Lys Ser Asn Thr Val Ala Ser Leu His Thr  
 2850 2855 2860  
 40  
 gaa aaa aal aca ctg gag cll agl aal gga gly all gtc aag ala aac 8762  
 Glu Lys Asn Thr Leu Glu Leu Ser Asn Gly Val Ile Val Lys Ile Asn  
 2865 2870 2875  
 aat cag ctt acc ctg gat agc aac act aaa tac ttc cac aaa ttg aac 8810  
 Asn Gln Leu Thr Leu Asp Ser Asn Thr Lys Tyr Phe His Lys Leu Asn  
 2880 2885 2890  
 45  
 atc ccc aca ctg gac ttc tct agt cag gct gac ctg cgc aac gag atc 8858  
 Ile Pro Lys Leu Asp Phe Ser Ser Gln Ala Asp Leu Arg Asn Glu Ile  
 2895 2900 2905 2910  
 aag aca ctg ttg aaa gct ggc cac ata gca tgg act tct tct gga aaa 8906  
 Lys Thr Leu Leu Lys Ala Gly His Ile Ala Trp Thr Ser Ser Gly Lys  
 2915 2920 2925  
 50  
 ggg tca tgg aaa tgg gcc tgc ccc aga ttc tca gal gag gga aca cal 8954  
 Gly Ser Trp Lys Trp Ala Cys Pro Arg Phe Ser Asp Glu Gly Thr His  
 2930 2935 2940 2945  
 55



EP 2 336 318 A1

		2930		2935		2940		
		gaa lea caa all agl llc acc ala gaa gga ccc ctc acf lcc lll gga	9002					
5		Glu Ser Gln Ile Ser Phe Thr Ile Glu Gly Pro Leu Thr Ser Phe Gly						
		2945		2950		2955		
		ctg tcc aat aac atc aat agc aaa cac cta aga gta aac caa aac ttg	9050					
		Leu Ser Asn Lys Ile Asn Ser Lys His Leu Arg Val Asn Gln Asn Leu						
		2960		2965		2970		
10		ggt tat gaa tct ggc tcc ctc aac ttt tct aaa ctt gaa att caa tca	9098					
		Val Tyr Glu Ser Gly Ser Leu Asn Phe Ser Lys Leu Glu Ile Gln Ser						
		2975		2980		2985		2990
		caa gtc gat tcc cag cat gtg ggc cac agt gtt cta act gct aaa ggc	9146					
		Gln Val Asp Ser Gln His Val Gly His Ser Val Leu Thr Ala Lys Gly						
		2995		3000		3005		
15		a-g gca clg lll gga gaa ggg aag gca gag lll acf ggg agg cal gal	9194					
		Met Ala Leu Phe Gly Glu Gly Lys Ala Glu Phe Thr Gly Arg His Asp						
		3010		3015		3020		
		gct cat tta aat gga aag gtt att gga act ttg aaa aat tct ctt ttc	9242					
		Ala His Leu Asn Gly Lys Val Ile Gly Thr Leu Lys Asn Ser Leu Phe						
		3025		3030		3035		
20		ttt tca gcc cag cca ttt gag atc acg gca tcc aca aac aat gaa ggg	9290					
		Phe Ser Ala Gln Pro Phe Glu Ile Thr Ala Ser Thr Asn Asn Glu Gly						
		3040		3045		3050		
		aat ttg aaa gtt cgt ttt cca tta agg tta aca ggg aag ata gac ttc	9338					
		Asn Leu Lys Val Arg Phe Pro Leu Arg Leu Thr Gly Lys Ile Asp Phe						
25		3055		3060		3065		3070
		ctg aat aac tat gca ctg ttt ctg agt ccc agt gcc cag caa gca agt	9386					
		Leu Asn Asn Tyr Ala Leu Phe Leu Ser Pro Ser Ala Gln Gln Ala Ser						
		3075		3080		3085		
		tgg caa gta agt gct agg ttc aat cag tat aag tac aac caa aat ttc	9434					
		Trp Gln Val Ser Ala Arg Phe Asn Gln Tyr Lys Tyr Asn Gln Asn Phe						
30		3090		3095		3100		
		tct gct gga aac aac gag aac att atg gag gcc cat gta gga ata aat	9482					
		Ser Ala Gly Asn Asn Glu Asn Ile Met Glu Ala His Val Gly Ile Asn						
		3105		3110		3115		
		gga gaa gca aat ctg gat ttc tta aac att cct tta aca att cct gaa	9530					
		Gly Glu Ala Asn Leu Asp Phe Leu Asn Ile Pro Leu Thr Ile Pro Glu						
35		3120		3125		3130		
		atg cgt cta cct tac aca ata atc aca act cct cca ctg aaa gat ttc	9578					
		Met Arg Leu Pro Tyr Thr Ile Ile Thr Thr Pro Pro Leu Lys Asp Phe						
		3135		3140		3145		3150
		tct cta tgg gaa aaa aca ggc ttg aag gaa ttc ttg aaa acg aca aag	9626					
		Ser Leu Trp Glu Lys Thr Gly Leu Lys Glu Phe Leu Lys Thr Thr Lys						
		3155		3160		3165		
		caa tca ttt gat tta agt gta aaa gcc cag tat aag aaa aac aaa cac	9674					
		Gln Ser Phe Asp Leu Ser Val Lys Ala Gln Tyr Lys Lys Asn Lys His						
		3170		3175		3180		
45		agg cat tcc atc aca aat cct ttg gcc gtg ctt tgt gag ttt aac agt	9722					
		Arg His Ser Ile Thr Asn Pro Leu Ala Val Leu Cys Glu Phe Ile Ser						
		3185		3190		3195		
		cag agc atc aaa tcc ttt gac agg cat ttt gaa aaa aac aga aac aat	9770					
		Gln Ser Ile Lys Ser Phe Asp Arg His Phe Glu Lys Asn Arg Asn Asn						
		3200		3205		3210		
50		gca tta gat ttt gtc acc aaa tcc tat aat gac aca aaa att aag ttt	9818					
		Ala Leu Asp Phe Val Thr Lys Ser Tyr Asn Glu Thr Lys Ile Lys Phe						
		3215		3220		3225		3230

55

EP 2 336 318 A1

gat aag tac aaa gct gaa aaa tct cac gac gag ctc ccc agg acc ttt 9866  
 Asp Lys Tyr Lys Ala Glu Lys Ser His Asp Glu Leu Pro Arg Thr Phe  
 3235 3240 3245

5  
 caa att cct gga tac act gtt cca gtt gtc aat gtt gaa gtg tct cca 9914  
 Gln Ile Pro Gly Tyr Thr Val Pro Val Val Asn Val Glu Val Ser Pro  
 3250 3255 3260

10  
 ttc acc ata gag atg tgg gca ttc gcc tat gtg ttc cca aaa gca gtc 9962  
 Phe Thr Ile Glu Met Ser Ala Phe Gly Tyr Val Phe Pro Lys Ala Val  
 3265 3270 3275

15  
 agc atg cct agt ttc tcc atc cta ggt tct gac gtc cgt gtg cct tca 10010  
 Ser Met Pro Ser Phe Ser Ile Leu Gly Ser Asp Val Arg Val Pro Ser  
 3280 3285 3290

20  
 tac aca tta atc ctg cca tca tta gag ctg cca gtc ctt cat gtc cct 10058  
 Tyr Thr Leu Ile Leu Pro Ser Leu Glu Leu Pro Val Leu His Val Pro  
 3295 3300 3305 3310

25  
 aga aat ctc aag ctt tct ctt cca cat ttc aag gaa ttg tgt acc ata 10106  
 Arg Asn Leu Lys Leu Ser Leu Pro His Phe Lys Glu Leu Cys Thr Ile  
 3315 3320 3325

30  
 agc cat att ttt att cct gcc atg gcc aat att acc tat gat ttc tcc 10154  
 Ser His Ile Phe Ile Pro Ala Met Gly Asn Ile Thr Tyr Asp Phe Ser  
 3330 3335 3340

35  
 ttt aaa tca agt gtc atc aca ctg aat acc aat got gaa ctt ttt aac 10202  
 Phe Lys Ser Ser Val Ile Thr Leu Asn Thr Asn Ala Glu Leu Phe Asn  
 3345 3350 3355

40  
 cag tca gat att gtt gct cat ctc ctt tct tca tct tca tct gtc att 10250  
 Gln Ser Asp Ile Val Ala His Leu Ser Ser Ser Ser Ser Val Ile  
 3360 3365 3370

45  
 gat gca ctg cag tac aaa tta gag gcc acc aca aga ttg aca aga aaa 10298  
 Asp Ala Leu Gln Tyr Lys Leu Glu Gly Thr Thr Arg Leu Thr Arg Lys  
 3375 3380 3385 3390

50  
 aag gga ttg aac tta gcc aca gct ctg tct ctg aac aac aaa ttt gtg 10346  
 Arg Gly Leu Lys Leu Ala Thr Ala Leu Ser Leu Ser Asn Lys Phe Val  
 3395 3400 3405

55  
 gag ggt agt cat aac agt act gtg agc tta acc acg aaa aat atg gaa 10394  
 Glu Gly Ser His Asn Ser Thr Val Ser Leu Thr Thr Lys Asn Met Glu  
 3410 3415 3420

60  
 gtg tca gtg gca aaa acc aca aaa gcc gaa att cca att ttg aga atg 10442  
 Val Ser Val Ala Lys Thr Thr Lys Ala Glu Ile Pro Ile Leu Arg Met  
 3425 3430 3435

65  
 aat ttc aag caa gaa ctt aal gga aal acc aag tca aaa cct act gtc 10490  
 Asn Phe Lys Gln Glu Leu Asn Gly Asn Thr Lys Ser Lys Pro Thr Val  
 3440 3445 3450

70  
 tct tcc tcc atg gaa ttt aag tat gat ttc aat tct tca atg ctg tac 10538  
 Ser Ser Ser Met Glu Phe Lys Tyr Asp Phe Asn Ser Ser Met Leu Tyr  
 3455 3460 3465 3470

75  
 tct acc gct aaa gga gca gtt gac cac aag ctt agc ttg gaa agc ctc 10586  
 Ser Thr Ala Lys Gly Ala Val Asp His Lys Leu Ser Leu Glu Ser Leu  
 3475 3480 3485

80  
 acc tct tac ttt tcc att gag tca tct acc aae gga gat gtc aag ggt 10634  
 Thr Ser Tyr Phe Ser Ile Glu Ser Ser Thr Lys Gly Asp Val Lys Gly  
 3490 3495 3500

85  
 tgg gtt ctt tct cgg gaa tal tca gga act att gct agt gag gcc aac 10682  
 Ser Val Leu Ser Arg Glu Tyr Ser Gly Thr Ile Ala Ser Glu Ala Asn

EP 2 336 318 A1

	3505		3510		3515		
5	act lac llg aal lcc aag agc aca cgg lcl lca glg aag clg cag gcc					10730	
	Thr Tyr Leu Asn Ser Lys Ser Thr Arg Ser Ser Val Lys Leu Gln Gly						
	3520		3525		3530		
	act tcc aaa att gat gat atc tgg aac ctt gaa gta aaa gaa aat ttt					10778	
	Thr Ser Lys Ile Asp Asp Ile Trp Asn Leu Glu Val Lys Glu Asn Phe						
	3535		3540		3545		3550
10	gct gga gaa gcc aca ctc caa cgc ata tat tcc ctc tgg gag cac agt					10826	
	Ala Gly Glu Ala Thr Leu Gln Arg Ile Tyr Ser Leu Trp Glu His Ser						
		3555		3560		3565	
	acg aaa aac cac tta cag cta gag gcc ctc ttt ttc acc aac gga gaa					10874	
	Thr Lys Asn His Leu Gln Leu Glu Gly Leu Phe Phe Thr Asn Gly Glu						
		3570		3575		3580	
15	cal aca agc aaa gcc acc clg gaa ctc lcl cca lgg caa alg lca gcl					10922	
	His Thr Ser Lys Ala Thr Leu Glu Leu Ser Pro Trp Gln Met Ser Ala						
		3585		3590		3595	
	ctt gtt cag gtc cat gca agt cag ccc agt tcc ttc cat gat ttc cot					10970	
	Leu Val Gln Val His Ala Ser Gln Pro Ser Ser Phe His Asp Phe Pro						
		3600		3605		3610	
20	gac ctt gcc cag gaa gtg gcc ctg aat gct aac act aag aac cag aag					11018	
	Asp Leu Gly Gln Glu Val Ala Leu Asn Ala Asn Thr Lys Asn Gln Lys						
		3615		3620		3625	3630
	atc aga tgg aaa aat gaa gtc cgg att cat tct ggg tct ttc cag agc					11066	
	Ile Arg Trp Lys Asn Glu Val Arg Ile His Ser Gly Ser Phe Gln Ser						
		3635		3640		3645	
25	cag gtc gag ctt tcc aat gac caa gaa aag gca cac ctt gac att gca					11114	
	Gln Val Glu Leu Ser Asn Asp Gln Glu Lys Ala His Leu Asp Ile Ala						
		3650		3655		3660	
	gga tcc tta gaa gga cac cta agg ttc ctc aaa aat atc atc cta cca					11162	
	Gly Ser Leu Glu Gly His Leu Arg Phe Leu Lys Asn Ile Ile Leu Pro						
		3665		3670		3675	
30	gtc tat gac aag agc tta tgg gat ttc cta aag ctg gat gta acc acc					11210	
	Val Tyr Asp Lys Ser Leu Trp Asp Phe Leu Lys Leu Asp Val Thr Thr						
		3680		3685		3690	
35	agc all ggl agg aga cag cal cll cgl gll lca acl gcc lll glg lac					11258	
	Ser Ile Gly Arg Arg Gln His Leu Arg Val Ser Thr Ala Phe Val Tyr						
		3695		3700		3705	3710
	acc aaa aac ccc aat gcc tat tca ttc tcc atc cct gta aaa gtt ttg					11306	
	Thr Lys Asn Pro Asn Gly Tyr Ser Phe Ser Ile Pro Val Lys Val Leu						
		3715		3720		3725	
40	gct gat aaa ttc att act cct ggg ctg aaa cta aat gat cta aat tca					11354	
	Ala Asp Lys Phe Ile Thr Pro Gly Leu Lys Leu Asn Asp Leu Asn Ser						
		3730		3735		3740	
	gct ctt gtc atg cct acg ttc cat gtc cca ttt aca gat ctt cag gtt					11402	
	Val Leu Val Met Pro Thr Phe His Val Pro Phe Thr Asp Leu Gln Val						
		3745		3750		3755	
45	cca tcc tgc aaa ctt gac ttc aga gaa ata caa atc cat aag aag ctg					11450	
	Pro Ser Cys Lys Leu Asp Phe Arg Glu Ile Gln Ile Tyr Lys Lys Leu						
		3760		3765		3770	
50	aga act tca tca ttt gcc ctc aac cta cca aca ctc ccc gag gta aaa					11498	
	Arg Thr Ser Ser Phe Ala Leu Asn Leu Pro Thr Leu Pro Glu Val Lys						
		3775		3780		3785	3790
55							

EP 2 336 318 A1

ttc cct gaa gtt gat gtg tta aca aaa tat tct caa coa gaa gac tcc 11546  
 Phe Pro Glu Val Asp Val Leu Thr Lys Tyr Ser Gln Pro Glu Asp Ser  
 3795 3800 3805  
 5  
 ttg att ccc ttt ttt gag ata acc gtg cct gaa tct cag tta act gtg 11594  
 Leu Ile Pro Phe Phe Glu Ile Thr Val Pro Glu Ser Gln Leu Thr Val  
 3810 3815 3820  
 tcc cag ttc acg ctt cca aaa agt gtc tca gat ggc att gct gct ttg 11642  
 Ser Gln Phe Thr Leu Pro Lys Ser Val Ser Asp Gly Ile Ala Ala Leu  
 3825 3830 3835  
 10  
 cat cta aat gca gta gcc aac aag atc gca gac ttt gag ttg ccc acc 11690  
 Asp Leu Asn Ala Val Ala Asn Lys Ile Ala Asp Phe Glu Leu Pro Thr  
 3840 3845 3850  
 atc atc gtg cct gag cag acc att gag att ccc tcc att aag ttc tct 11738  
 Ile Ile Val Pro Glu Gln Thr Ile Glu Ile Pro Ser Ile Lys Phe Ser  
 3855 3860 3865 3870  
 15  
 gta cct gct gga att gtc att cct tcc ttt caa gca ctg act gca cgc 11786  
 Val Pro Ala Gly Ile Val Ile Pro Ser Phe Gln Ala Leu Thr Ala Arg  
 3875 3880 3885  
 lll gag gla gac tcl ccc glg tal aal gcc act lgg agl gcc agl llg 11834  
 Phe Glu Val Asp Ser Pro Val Tyr Asn Ala Thr Trp Ser Ala Ser Leu  
 3890 3895 3900  
 aaa aac aaa gca gat tat gtt gaa aca gtc ctg gat tcc aca tgc acc 11882  
 Lys Asn Lys Ala Asp Tyr Val Glu Thr Val Leu Asp Ser Thr Cys Ser  
 3905 3910 3915  
 25  
 tca acc gta cag ttc cta gaa tat gaa cta aat gtt ttg gga aca cac 11930  
 Ser Thr Val Gln Phe Leu Glu Tyr Glu Leu Asn Val Leu Gly Thr His  
 3920 3925 3930  
 aaa atc gaa gat ggt acg tta gcc tct aag act aaa gga aca ctt gca 11978  
 Lys Ile Glu Asp Gly Thr Leu Ala Ser Lys Thr Lys Gly Thr Leu Ala  
 3935 3940 3945 3950  
 30  
 cac cgt gac ttc agt gca gaa tat gaa gaa gat ggc aaa ttt gaa gca 12026  
 His Arg Asp Phe Ser Ala Glu Tyr Glu Glu Asp Gly Lys Phe Glu Gly  
 3955 3960 3965  
 ctt cag gaa tgg gaa gga aaa gcg cac ctg aat atc aaa agc cca gcg 12074  
 Leu Gln Glu Trp Glu Gly Lys Ala His Leu Asn Ile Lys Ser Pro Ala  
 3970 3975 3980  
 35  
 ttc acc gat ctg cat ctg cgc tac cag aaa gac aag aaa gcg atc tcc 12122  
 Phe Thr Asp Leu His Leu Arg Tyr Gln Lys Asp Lys Lys Gly Ile Ser  
 3985 3990 3995  
 acc tca gca gcc tcc cca gcc gta gcc acc gtg gcc atg gat atg gat 12170  
 Thr Ser Ala Ala Ser Pro Ala Val Gly Thr Val Gly Met Asp Met Asp  
 4000 4005 4010  
 gaa gat gac gac ttt tct aaa tgg aac ttc tac tac agc cct cag tcc 12218  
 Glu Asp Asp Asp Phe Ser Lys Trp Asn Phe Tyr Tyr Ser Pro Gln Ser  
 4015 4020 4025 4030  
 45  
 tct cca gat aaa aaa ctg acc ata ttc aaa act gag ttg agg gtc cgg 12266  
 Ser Pro Asp Lys Lys Leu Thr Ile Phe Lys Thr Glu Leu Arg Val Arg  
 4035 4040 4045  
 caa tct gat gac gaa act cag atc aaa gtt aat tgg gaa gaa cag gca 12314  
 Glu Ser Asp Glu Glu Thr Gln Ile Lys Val Asn Trp Glu Glu Glu Ala  
 4050 4055 4060  
 50  
 gct tct gcc ttg cta acc tct ctg aaa gac aac gtg ccc aag gcc aca 12362  
 Ala Ser Gly Leu Leu Thr Ser Leu Lys Asp Asn Val Pro Lys Ala Thr  
 4065 4070 4075  
 55

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

ggg gtc ott tat gat tat gtc aac aag tac cac tgg gaa cac aca ggg 12410  
 Gly Val Leu Tyr Asp Tyr Val Asn Lys Tyr His Trp Glu His Thr Gly  
 4380 4085 409C

ctc acc ctg aga gaa gtg tct tca aag ctg aga aga aat ctg cag aac 12458  
 Leu Thr Leu Arg Glu Val Ser Ser Lys Leu Arg Arg Asn Leu Gln Asn  
 4095 4100 4105 4110

aat gct gag tgg gtt tat caa ggg gcc att agg caa att gat gat atc 12506  
 Asn Ala Glu Trp Val Tyr Gln Gly Ala Ile Arg Gln Ile Asp Asp Ile  
 4115 4120 4125

cac gtg aag ttc cag aaa gca gcc acg gcc acc act ggg acc tac caa 12554  
 Asp Val Arg Phe Gln Lys Ala Ala Ser Gly Thr Thr Gly Thr Tyr Gln  
 4130 4135 4140

gag tgg aag gac aag gcc cag aat ctg tac cag gaa ctg ttg act cag 12602  
 Glu Trp Lys Asp Lys Ala Gln Asn Leu Tyr Gln Glu Leu Leu Thr Gln  
 4145 4150 4155

gaa gcc caa gcc agt ttc cag gga ctc aag gat aac gtg tct gat gcc 12650  
 Glu Gly Gln Ala Ser Phe Gln Gly Leu Lys Asp Asn Val Phe Asp Gly  
 4160 4165 417C

tlg gla cga gll acf caa aaa ttc cal atg aaa glc aag cal ctg all 12698  
 Leu Val Arg Val Thr Gln Lys Phe His Met Lys Val Lys His Leu Ile  
 4175 4180 4185 4190

cac tca ctc att gat ttt ctg aac ttc ccc aga ttc cag ttt ccg ggg 12746  
 Asp Ser Leu Ile Asp Phe Leu Asn Phe Pro Arg Phe Gln Phe Pro Gly  
 4195 4200 4205

aaa cct ggg ata tac act agg gag gaa ctt tgc act atg ttc sta agg 12794  
 Lys Pro Gly Ile Tyr Thr Arg Glu Glu Leu Cys Thr Met Phe Ile Arg  
 4210 4215 4220

gag gta ggg acg gta ctg tcc cag gta tat tcg aaa gtc cat aat ggt 12842  
 Glu Val Gly Thr Val Leu Ser Gln Val Tyr Ser Lys Val His Asn Gly  
 4225 4230 4235

tca gaa ata ctg ttt tcc tat ttc caa gac cta gtg att aca ctt cct 12890  
 Ser Glu Ile Leu Phe Ser Tyr Phe Gln Asp Leu Val Ile Thr Leu Pro  
 4240 4245 425C

ttc gag tta agg aaa cat aaa cta ata gat gta atc tcg atg tat agg 12938  
 Phe Glu Leu Arg Lys His Lys Leu Ile Asp Val Ile Ser Met Tyr Arg  
 4255 4260 4265 4270

gaa ctg ttg aaa gat tta tca aaa gaa gcc caa gag gta ttt aaa gcc 12986  
 Glu Leu Leu Lys Asp Leu Ser Lys Glu Ala Gln Glu Val Phe Lys Ala  
 4275 4280 4285

aat cag tct ctc aag acc aca gag gtg cta cgt aat ctt cag gac ctt 13034  
 Ile Gln Ser Leu Lys Thr Thr Glu Val Leu Arg Asn Leu Gln Asp Leu  
 4290 4295 4300

tta caa ttc att ttc caa cta ata gaa gat aac att aaa cag ctg aaa 13082  
 Leu Gln Phe Ile Phe Gln Leu Ile Glu Asp Asn Ile Lys Gln Leu Lys  
 4305 4310 4315

gag atg asa ttt act tat ctt att aac tat atc caa gat gag atc aac 13130  
 Glu Met Lys Phe Thr Tyr Leu Ile Asn Tyr Ile Gln Asp Glu Ile Asn  
 4320 4325 433C

aca atc ttc aat gat tat atc cca tat gtt ttt aaa ttg ttg aca gaa 13178  
 Thr Ile Phe Asn Asp Tyr Ile Pro Tyr Val Phe Lys Leu Leu Lys Glu  
 4335 4340 4345 4350

aac cta tgc cll aal cll cal aag ttc aal gaa lll all caa aac gag 13226  
 Asn Leu Cys Leu Asn Leu His Lys Phe Asn Glu Phe Ile Gln Asn Glu

EP 2 336 318 A1

		4355		4360		4365		
	5	ccl cag gaa gcl lcl caa gag lla cag cag atc cal caa lac all alg					13274	
		Leu Gln Glu Ala Ser Gln Glu Leu Gln Gln Ile His Gln Tyr Ile Met	4370	4375		4380		
		ccc ctt cgt gaa gaa tat ttt gat cca agt ata gtt gcc tgg aca gtg					13322	
		Ala Leu Arg Glu Glu Tyr Phe Asp Pro Ser Ile Val Gly Trp Thr Val	4385	4390		4395		
	10	aaa tat tat gaa ctt gaa gaa aag ata gtc agt ctg atc aag aac ctg					13370	
		Lys Tyr Tyr Glu Leu Glu Glu Lys Ile Val Ser Leu Ile Lys Asn Leu	4400	4405		4410		
		tta gtt gct ctt aag gac ttc cat tcc gaa tat att gtc agt gcc tcc					13418	
		Leu Val Ala Leu Lys Asp Phe His Ser Glu Tyr Ile Val Ser Ala Ser	4415	4420		4425	4430	
	15	aac lll acf lcc caa clc lca agl caa gll gag caa lll clg cac aga					13466	
		Asn Phe Thr Ser Gln Leu Ser Ser Gln Val Glu Gln Phe Leu His Arg	4435	4440		4445		
		aat att cag gaa tat ctt agc atc ccc acc gat cca gat gga aaa ggg					13514	
		Asn Ile Gln Glu Tyr Leu Ser Ile Leu Thr Asp Pro Asp Gly Lys Gly	4450	4455		4460		
	20	aaa gag aag att gca gag ctt tct gcc act gct cag gaa ata att aaa					13562	
		Lys Glu Lys Ile Ala Glu Leu Ser Ala Thr Ala Gln Glu Ile Ile Lys	4465	4470		4475		
		agc cag gcc att gcg acg aag aaa ata att tct gat tac cac cag cag					13610	
		Ser Gln Ala Ile Ala Thr Lys Lys Ile Ile Ser Asp Tyr His Gln Gln	4480	4485		4490		
	25	ttt aga tat aaa ctg caa gat ttt tca gac caa ctc tct gat tac tat					13658	
		Phe Arg Tyr Lys Leu Gln Asp Phe Ser Asp Gln Leu Ser Asp Tyr Tyr	4495	4500		4505	4510	
		gaa aaa ttt att gct gaa tcc aaa aga ttg att gac ctg tcc att caa					13706	
		Glu Lys Phe Ile Ala Glu Ser Lys Arg Leu Ile Asp Leu Ser Ile Gln	4515	4520		4525		
	30	aac tac cac aca ttt ctg ata tac atc acg gag tta ctg aaa aag ctg					13754	
		Asn Tyr His Thr Phe Leu Ile Tyr Ile Thr Glu Leu Leu Lys Lys Leu	4530	4535		4540		
		caa tca acc aca gtc atg aac ccc tac atg aag ctt gct cca gga gaa					13802	
		Gln Ser Thr Thr Val Met Asn Pro Tyr Met Lys Leu Ala Pro Gly Glu	4545	4550		4555		
	35	ctt act atc atc ctc taa ttttttaaaa gaaatcttca tttattcttc					13850	
		Leu Thr Ile Ile Leu *	4560					
	40	ttttccaatt gaactttcac atagcacaga aaaaattcaa actgcctata ttgataaac					13910	
		catacagtga gccagccttg cagtaggcag tagactataa gcagaagc acatgaactg					13970	
		gacctgcacc aaagctggca ccagggtctg gaaggctctt gaactcagaa gcatggcatt					14030	
		ttttgcaagt taagaaaaa caggatctga gtatttttc taaacttggg gcaggagcaa					14090	
	45	caaataaatg gactctttat tgtgtatcat a					14121	
		<210> 4						
		<211> 21						
		<212> DNA						
		<213> Artificial Sequence						
	50	<220>						
		<223> PCR Primer						
		<400> 4						
	55							

EP 2 336 318 A1

tgctaaaggc acatattggc t 21

5 <210> 5  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

10 <400> 5  
 ctcagggttg actccatt gag 23

<210> 6  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence

15 <220>  
 <223> PCR Probe

<400> 6  
 ctgtcagag ggatcctaac actggcgg 28

20 <210> 7  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

25 <400> 7  
 gaaggagaag gtcggagtc 19

<210> 8  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> PCR Primer

<400> 8  
 gaagatggtg atgggatttc 20

35 <210> 9  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Probe

40 <400> 9  
 caagcttccc gllccagcc 20

<210> 10  
 <211> 2354  
 <212> DNA  
 <213> Mus musculus

45 <400> 10  
 gaattccaac ttccaccct ctacataca atgaaatac ctgcttttgg caaactgcac 60  
 agcatcctta agatccaatc tctctctttt atattgatg ctaatgcaa cacacagaat 120  
 gtaacaactt caggaacaa agcagagatt gggcttctg tcaactgctaa aggagagacc 180

50 caatttgaag ctctcaattt tgattttcaa gcacaagctc aattcctgga gtaaatcct 240  
 calcclocag lccggaaga alccalgaac lcclocagla agcalglgag aaggagcal 300

55

EP 2 336 318 A1

gagggtaga tagtatttga tggaaaggcc atgagggga aatcagacac agtcgcaagt 360  
 Llacacacag agaaaaalga aglagaglll aalaalggla lgacglcaa aglaaacaaL 420  
 5 cagctcacc0 ttgacagtea cacaaagtac tccacaagt tgagtgttcc taggctegac 480  
 tctccaagta aggcctctcc taataatgaa atcaagacac tattagaagc tggacatgtg 540  
 gcattgacat ctccagggac agggctcatgg aactgggcoct gtoccaaactt ctgggatgaa 600  
 10 ggcatacatt cgtcccaaaL tagctttact ggggatggc0 ccatcgcttL tgttggacta 660  
 tccaataaca taaatggcaa acacttacgg gtcatacaaa aactgactta tgaatctggc 720  
 tccctcaact attc-aagtl- tgaagttgag tcaaaagttg aatctcagca cgtgggcacc 780  
 agcattctaa cagccaatgg tggggcactg cccaaggacg caaaggcaga aatgactggc 840  
 15 gagcacaalg ccaacllaaa lggaaaagll aLlggaacll lgaaaaallc tcccllcLL 900  
 tcaqcacaac catttgaat tactgcctcc acaataatg aaggaaattt gaaagtccgt 960  
 tttccactaa agctgactgg gaaaatagac tccctgaata actatgcatt gttctcagc 1020  
 ccccgtagcc aacaagcaag ctggcaagcg agtaccagat tcaatcagta caaatacaat 1080  
 20 caaaactttt ctgcataaaa caatgaccac aacatagaag ccagtatagg aatgaatgga 1140  
 gatgcaacc tggatttctL aaacatacct taaacaattc ctgaaattaa ctgacctac 1200  
 acggagttca aaacccctL actgaaggat tctccatat gggagaagac aggcttgaaa 1260  
 25 gaalLLLlgc agacaacaaa gcaalcaLLL galllgaglg laaaggclca alalaaaaag 1320  
 aacagtgaca agcattccat tgttgcctcc ctgggtatgt tttatgaatt tactctcaac 1380  
 aatgtcaatt cgtcggacag asaatttgag aaagtcagaa acaatgcttt acattttctt 1440  
 accacctcct ataaagaagc aaaaattaag gttgataagt acaaaactga aaattccctt 1500  
 30 aatcagccct ctgggaactL tcaaaatcat ggctacactc tccagttgt caacattgaa 1560  
 gtatctecat ttgcgtaga gacactggct tccaggcatg tgatcccac agcaataagc 1620  
 accccaagtg tcaaatccc tggctcaac atcatgggc ctccatacaa gttagtctg 1680  
 ccaccctgg agttgccagL tttccatggc cctgggaatc tattcaagtt ttcctccca 1740  
 35 galllcaagg gallcaacac tallgacaal aLLlalallc cagccalggg caaclllacc 1800  
 tatgactttt ctttaaatc aagtgtcatc aactgaata ccaatgctgg actttataac 1860  
 caatcagata tcttgccca tttcctttct tctcttcat ttgtactga cgcctccag 1920  
 40 tacaattag agggaacatc acgtctgatg cgaaaaaggg gattgaaact agccacagct 1980  
 gctctctaa ctacaatL tgtaaaggc agtcatgaca gcaccattag ttaaccaag 2040  
 aaaaacatg aagcatcagL gagaacaact gccaacctcc atgcctccat attctcaatg 2100  
 aacttcaagc aggaacttae tggaaatacc aagtcaaac ccaactgttc atcatccatt 2160  
 45 gaactaaact atgacttcaa ttcctcaaag ctgcactcta ctgcaacagg aggcattgat 2220  
 cacaagttca gottagaag tctcaactcc tacttttcca ttgagtcatt caccaaagga 2280  
 aatatcaaga gttccttccL tctcaggaa tattcaggaa gtgttgccaa tcaagccaat 2340  
 50 gTatatctga attc 2354  
 <210> 11  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

55



EP 2 336 318 A1

	<220>		
	<223> PCR Primer		
5	<400> 11	ogtgggctcc agcattcta	19
	<210> 12		
	<211> 21		
	<212> DNA		
10	<213> Artificial Sequence		
	<220>		
	<223> PCR Primer		
	<400> 12	agtcatttct gcctttgcct c	21
15	<210> 13		
	<211> 22		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> PCR Probe		
20	<400> 13	ccaatggctg ggcactgctc aa	22
	<210> 14		
	<211> 20		
	<212> DNA		
25	<213> Artificial Sequence		
	<220>		
	<223> PCR Primer		
	<400> 14	ggcaaatcca acggcacagt	20
30	<210> 15		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> PCR Primer		
35	<400> 15	gggtctcgtc cctggaaga	20
	<210> 16		
	<211> 27		
	<212> DNA		
40	<213> Artificial Sequence		
	<220>		
	<223> PCR Probe		
	<400> 16	aaggccgaga atgggaagct tgcata	27
45	<210> 17		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
50	<400> 17	ccgcaggctc cgtgggaa	20
55			

EP 2 336 318 A1

5  
 <210> 18  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 18  
 accgagaagg gcactcagcc 20

10  
 <210> 19  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 19  
 ccctcqqcct cqqcqcctq 20

20  
 <210> 20  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 20  
 tccatcgcca gctcgcgtcg 20

25  
 <210> 21  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 21  
 cagcgccagc agcggcagca 20

30  
 <210> 22  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 22  
 gcccgccagc agcagcagca 20

35  
 <210> 23  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 23  
 ctgaaatcag cagtcccagg 20

40  
 <210> 24  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 24  
 ctgaaatcag cagtcccagg 20

45  
 <210> 25  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 25  
 ctgaaatcag cagtcccagg 20

50  
 <210> 26  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 26  
 ctgaaatcag cagtcccagg 20

55  
 <210> 27  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 27  
 ctgaaatcag cagtcccagg 20

EP 2 336 318 A1

	<400> 24	
	c...cagcaag gcttggccc	20
5	<210> 25	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
10	<400> 25	
	t...ctgttgc cacattgcc	20
	<210> 26	
	<211> 20	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 26	
	ggaagagggtg ttgcacctg	20
20	<210> 27	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
25	<400> 27	
	tgtgtacca tccatactc	20
	<210> 28	
	<211> 20	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 28	
	tcaaatgcga ggccatctc	20
35	<210> 29	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
40	<400> 29	
	ggacaacctca atcagctgtg	20
	<210> 30	
	<211> 20	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 30	
	tcagggccac caggtaggtg	20
50	<210> 31	
	<211> 20	
	<212> DNA	
55		

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
 <400> 31  
 ctaatattca tccccagtcc 20  
 <210> 32  
 <211> 20  
 <212> DNA  
 10  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 32  
 tgctccatgg ttggccca- 20  
 15  
 <210> 33  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 20  
 <400> 33  
 gcagccagtc gtttatctcc 20  
 <210> 34  
 <211> 20  
 <212> DNA  
 25  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 34  
 gatatgcca agtggccac 20  
 30  
 <210> 35  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 35  
 <400> 35  
 cccaggagct ggaggtcatg 20  
 <210> 36  
 <211> 20  
 <212> DNA  
 40  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 36  
 t-gagccctt cctgatgacc 20  
 45  
 <210> 37  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 50  
 <400> 37  
 a-ctggacc cactcctagc 20  
 55

5  
 <210> 38  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 38  
 cagaccogac togtggaaga 20

10  
 <210> 39  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 39  
 gccctcagta gattcatca 20

15  
 <210> 40  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 40  
 gccalgccac cctc\_lggaa 20

20  
 <210> 41  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 41  
 aacccacgtg ccgcaaatc 20

25  
 <210> 42  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 42  
 actcccagat gcctctgaa 20

30  
 <210> 43  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 43  
 atgtgtaac gagccgaag 20

35  
 <210> 44  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

40  
 <210> 45  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

45  
 <210> 46  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

50  
 <210> 47  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

55

EP 2 336 318 A1

5 <223> Antisense Oligonucleotide  
 <400> 44  
 ggcgtagaga cccaacacac 20  
 <210> 45  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 45  
 g-gtttaggat ccctctgaca 20  
 <210> 46  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 46  
 ccacaglgata gclc-uglgag 20  
 <210> 47  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 47  
 a-tttcaqcat atgaqccac 20  
 <210> 48  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 48  
 ccctgaacct tagcaacagc 20  
 <210> 49  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 49  
 gctgaagcca gccacagca 20  
 <210> 50  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 50  
 acagctgcc agta-gttcc 20  
 <210> 51  
 <211> 20

55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 51  
 cccaataaga tttataacaa 20  
  
 <210> 52  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 52  
 lggcclacca gagacaggla 20  
 15  
  
 <210> 53  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 20  
  
 <400> 53  
 tcatacgttt agccaatc 20  
  
 <210> 54  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 54  
 gcatggtccc aagcatggtc 20  
 30  
  
 <210> 55  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 35  
  
 <400> 55  
 agtgatggaa gctgcatac 20  
  
 <210> 56  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 56  
 atgagcatca tgcctcccag 20  
 45  
  
 <210> 57  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 50  
  
 <400> 57  
  
 55

EP 2 336 318 A1

	gaacacatag ccgaatgccg	20
5	<210> 58 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 58 g-gggtgccct ctaa-attgta	20
	<210> 59 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 59 cccgagaaaag aaccgaacc	20
20	<210> 60 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 60 tgccctgcag cttcactgaa	20
	<210> 61 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 61 gaaatcccat aagc-cttgc-	20
35	<210> 62 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 62 agaagctgcc cLlLcLlLcc	20
	<210> 63 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 63 tcagggtgag ccctgtgtc-	20
50	<210> 64 <211> 20 <212> DNA <213> Artificial Sequence	
55		



EP 2 336 318 A1

	<220>	
	<223> Antisense Oligonucleotide	
5	<400> 64	
	c=aatggccc cttgataaac	20
	<210> 65	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Antisense Oligonucleotide	
	<400> 65	
	acgttatcct tgaçtcocctg	20
	<210> 66	
15	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
20	<400> 66	
	tatatcccag gtttccccg	20
	<210> 67	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Antisense Oligonucleotide	
	<400> 67	
	acctgggaca gtaccgtccc	20
	<210> 68	
30	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
35	<400> 68	
	c=ggcctactg caagcctggc	20
	<210> 69	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> Antisense Oligonucleotide	
	<400> 69	
	agaçacottc cgaçccocctg	20
	<210> 70	
45	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
50	<400> 70	
	a=gatacaca ataaagactc	20
	<210> 71	

55

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 71  
 atgtatgtg agagtgagg 20  
 <210> 72  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 72  
 gaggagattg gatcctaagg 20  
 <210> 73  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 73  
 ctcaaattg ggactctcc 20  
 <210> 74  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 74  
 tccaggaatt gagctgtgc 20  
 <210> 75  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 75  
 ttcaggactg gaggatgag 20  
 <210> 76  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 76  
 tctcacctc atgcaccat 20  
 <210> 77  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 77  
 tctcacctc atgcaccat 20  
 <210> 78  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 78  
 tctcacctc atgcaccat 20  
 <210> 79  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 79  
 tctcacctc atgcaccat 20  
 <210> 80  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 55  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 80  
 tctcacctc atgcaccat 20

EP 2 336 318 A1

5  
 <400> 77  
 tgactgtcaa gggtagctg 20  
 <210> 78  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 10  
 <400> 78  
 gccagccta ggaacctca 20  
 <210> 79  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 79  
 atgtcaatgc cacatgtcca 20  
 <210> 80  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 25  
 <400> 80  
 tccatccgag aagtgggac 20  
 <210> 81  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 81  
 attgggacg aatgatgcc 20  
 <210> 82  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 82  
 agttgaggaa gccagattca 20  
 <210> 83  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 83  
 tcccagtcg gcttagtgg 20  
 <210> 84  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 84  
 tcccagtcg gcttagtgg 20  
 <210> 85  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 85  
 tcccagtcg gcttagtgg 20  
 <210> 86  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 55  
 <220>  
 <223> Antisense Oligonucleotide

EP 2 336 318 A1

<220>  
 <223> Antisense Oligonucleotide  
 5 <400> 84  
 agcttgcttg ttgggcacgg 20  
 <210> 85  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 85  
 cctatactgg ctccatgtc 20  
 <210> 86  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 86  
 tgaactccgt gtaaggcaag 20  
 <210> 87  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 87  
 gagaaatcct tcagtaaggg 20  
 <210> 88  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 88  
 caatggaatg ctgcactg 20  
 <210> 89  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 89  
 gcttcattat aggaggtgg 20  
 <210> 90  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 90  
 acaactggga lagtglagcc 20  
 40  
 45  
 50  
 55

EP 2 336 318 A1

5 <210> 91  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 91  
 gtaggacca gggattgtga 20

10 <210> 92  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 92  
 accatqaaa actqcaact 20

15 <210> 93  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 93  
 tgggaggaa aactgaata 20

20 <210> 94  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 94  
 tgggcaacga taccgattg 20

25 <210> 95  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 95  
 ctagcagggcg tagtgacaa 20

30 <210> 96  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 96  
 gcatcagacg tgatgttccc 20

35 <210> 97  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

40  
 45  
 50  
 55

EP 2 336 318 A1

5 <400> 97  
c.LggLLaaa claaLggLgc 20

<210> 98  
<211> 20  
<212> DNA  
<213> Artificial Sequence

10 <220>  
<223> Antisense Oligonucleotide

<400> 98  
aTqqaqcat ggaqttgc 20

<210> 99  
<211> 20  
<212> DNA  
<213> Artificial Sequence

15 <220>  
<223> Antisense Oligonucleotide

<400> 99  
aatggatgat gaaacagtgc 20

20 <210> 100  
<211> 20  
<212> DNA  
<213> Artificial Sequence

25 <220>  
<223> Antisense Oligonucleotide

<400> 100  
aTcaatgcct cctgtgcag 20

30 <210> 101  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

35 <400> 101  
ggaagtgaga ctttctaagc 20

<210> 102  
<211> 20  
<212> DNA  
<213> Artificial Sequence

40 <220>  
<223> Antisense Oligonucleotide

<400> 102  
aggaaggaa ccttgatgc 20

<210> 103  
<211> 20  
<212> DNA  
<213> Artificial Sequence

45 <220>  
<223> Antisense Oligonucleotide

<400> 103  
aTtgcttca ttggcaaac 20

50 <210> 104  
<211> 20  
<212> DNA

55

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
 <400> 104  
 aggtqagga gttcqaattc 20  
 <210> 105  
 <211> 20  
 <212> DNA  
 10 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 105  
 t-gttccctg aagt-gttac 20  
 15 <210> 106  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 20 <223> Antisense Oligonucleotide  
 <400> 106  
 g-tcatggat t-cct-cagga 20  
 <210> 107  
 <211> 20  
 <212> DNA  
 25 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 107  
 a-gctccatt ctcacatgc 20  
 30 <210> 108  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 35 <223> Antisense Oligonucleotide  
 <400> 108  
 tgcgactgtg tctgatttc 20  
 <210> 109  
 <211> 20  
 <212> DNA  
 40 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 109  
 g-ccctgaag atgtcaatgc 20  
 45 <210> 110  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 50 <223> Antisense Oligonucleotide  
 <400> 110  
 aggcccagtt ccatgacct 20  
 55

EP 2 336 318 A1

5  
 <210> 111  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 111  
 ggagcccaag tgctgagatt 20

10  
 <210> 112  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 112  
 cgtccttgag cagtgccga 20

15  
 <210> 113  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 113  
 cccatagga gaaa\_ccttc 20

20  
 <210> 114  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 114  
 catgcctgga agccaqtgc 20

25  
 <210> 115  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 115  
 g-gttgaatc ccttgaatc 20

30  
 <210> 116  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 116  
 ggtaaagttg ccca-ggctg 20

35  
 <210> 117  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

40  
 <210> 118  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

45  
 <210> 119  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

50  
 <210> 120  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

55



EP 2 336 318 A1

5 <223> Antisense Oligonucleotide  
 <400> 117  
 gtataaaagt ccagcattgg 20  
 <210> 118  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 118  
 catcagacgt gatg-tcccc- 20  
 <210> 119  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 119  
 tggctaglll caatcccc- 20  
 <210> 120  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 120  
 c-qtcatgac tqccctttac 20  
 <210> 121  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 121  
 gottgaagtt cattgagaa- 20  
 <210> 122  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 122  
 t-cctgagaa aggaaggaac 20  
 <210> 123  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 123  
 tcagatatac attggettea 20  
 <210> 124  
 <211> 20

55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 124  
 ttcctcttgc gccctggcgc 20  
  
 <210> 125  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 125  
 cccacatgga acctcagcc 20  
  
 <210> 126  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 126  
 cctccagctc aacctgcag 20  
  
 <210> 127  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 127  
 gggttgaagc catacacctc 20  
  
 <210> 128  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 128  
 ccagcttgag ctcatacctg 20  
  
 <210> 129  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 129  
 ccctcttgat gttcaggatg 20  
  
 <210> 130  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 130  
  
 40  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 130  
  
 45  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 130  
  
 50  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 130  
  
 55

EP 2 336 318 A1

	gagcagtttc catacacgg-	20
5	<210> 131 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 131 cccttctctcg ttttgacgg-	20
	<210> 132 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 132 t-gaagcgat cacactgccc	20
20	<210> 133 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 133 gcctttgatg agagcaagtg	20
	<210> 134 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 134 tctctttagc gtccagtgtg	20
35	<210> 135 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 135 ccctctcagcl caglaaccag	20
	<210> 136 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 136 ccactgaggc tgtccacac-	20
50	<210> 137 <211> 20 <212> DNA <213> Artificial Sequence	
55		

EP 2 336 318 A1

	<220>		
	<223> Antisense Oligonucleotide		
5	<400> 137	cgctgatccc tggccatgtc	20
	<210> 138		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
10	<220>		
	<223> Antisense Oligonucleotide		
	<400> 138	gttgaccgog tggctcagcg	20
	<210> 139		
15	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
20	<400> 139	gcagctcctg ggtccctgta	20
	<210> 140		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
25	<220>		
	<223> Antisense Oligonucleotide		
	<400> 140	cccattgtag aatttggaca	20
	<210> 141		
30	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
35	<400> 141	aatctcgalg aggtcagctg	20
	<210> 142		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Antisense Oligonucleotide		
	<400> 142	gacaccatca ggaacttqac	20
	<210> 143		
45	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
50	<400> 143	gctcctctcc caagatgagg	20
	<210> 144		
55			

## EP 2 336 318 A1

<211> 20  
<212> DNA  
<213> Artificial Sequence

5 <220>  
<223> Antisense Oligonucleotide

<400> 144  
ggcaccatc agaagcagc 20

10 <210> 145  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

15 <400> 145  
agtccggaat gatgatgcc 20

<210> 146  
<211> 20  
<212> DNA  
<213> Artificial Sequence

20 <220>  
<223> Antisense Oligonucleotide

<400> 146  
ctgagcagct tgactggtc 20

25 <210> 147  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

30 <400> 147  
cccggtcagc ggatagtagg 20

<210> 148  
<211> 20  
<212> DNA  
<213> Artificial Sequence

35 <220>  
<223> Antisense Oligonucleotide

<400> 148  
tgtcacaact taggtggccc 20

40 <210> 149  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

45 <400> 149  
gctggcaat cccatgttc 20

<210> 150  
<211> 20  
<212> DNA  
<213> Artificial Sequence

50 <220>  
<223> Antisense Oligonucleotide

55

EP 2 336 318 A1

5  
 <400> 150  
 cccacagact tgaagtggag 20  
 <210> 151  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 10  
 <400> 151  
 gaactgcccc tcaatcttga 20  
 <210> 152  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 152  
 cccagagagg ccaagctctg 20  
 <210> 153  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 25  
 <400> 153  
 tgtgttcctt gaacggcca 20  
 <210> 154  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 154  
 acccagaatc atggcctgat 20  
 <210> 155  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 40  
 <400> 155  
 ggtgcctgtc tgctcagctg 20  
 <210> 156  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 156  
 atgtgaaact tgctctccc 20  
 50  
 <210> 157  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 55

EP 2 336 318 A1

<220>  
 <223> Antisense Oligonucleotide  
 5 <400> 157  
 tatgtctgca gttgagatag 20  
 <210> 158  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 158  
 t-gaatccag gatgcagtac 20  
 <210> 159  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 159  
 gagtctctga gtcacctcac 20  
 <210> 160  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 160  
 gatagaatat tgctctgcaa 20  
 <210> 161  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 161  
 cccttgctct accaatgct- 20  
 <210> 162  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 162  
 tccattccct atgtcagca- 20  
 <210> 163  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 163  
 gactccttca gagccagcgg 20  
 40  
 45  
 50  
 55

EP 2 336 318 A1

5 <210> 164  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 164  
 cccatgctcc gttctcaggc 20

10 <210> 165  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 165  
 ccqaqtcag cctcactaca 20

15 <210> 166  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 166  
 cagttagaac actctggccc 20

20 <210> 167  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 167  
 cagtgtgatg acacttgatt 20

25 <210> 168  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 168  
 ctgtggctaa ctccaatccc 20

30 <210> 169  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 169  
 cagtactggt atgactaccc 20

35 <210> 170  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

40  
 45  
 50  
 55



EP 2 336 318 A1

	<400> 170	
	cactgaagac cglgagctc	20
5	<210> 171	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
10	<400> 171	
	tcqtactgtg ctcccagaag	20
	<210> 172	
	<211> 20	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 172	
	aagaggccct cttagctgtaa	20
20	<210> 173	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
25	<400> 173	
	aagaccaga atgaatccgg	20
	<210> 174	
	<211> 20	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 174	
	gctactcca aagcgtgcag	20
35	<210> 175	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
40	<400> 175	
	tagaggctaa cgtaccatc	20
	<210> 176	
	<211> 20	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 176	
	ccatatccat gccacgggtg	20
50	<210> 177	
	<211> 20	
	<212> DNA	
55		

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
 <400> 177  
 agtttctca tcaattccc 20  
 <210> 178  
 <211> 20  
 <212> DNA  
 10 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 178  
 cccagtggta cttgtgaca 20  
 15 <210> 179  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 20 <223> Antisense Oligonucleotide  
 <400> 179  
 cccagtggta ccaatggctg 20  
 <210> 180  
 <211> 20  
 <212> DNA  
 25 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 180  
 gccaacagtt cctggtacag 20  
 30 <210> 181  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 35 <223> Antisense Oligonucleotide  
 <400> 181  
 ccctagtgtatatcccagg 20  
 <210> 182  
 <211> 20  
 <212> DNA  
 40 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 182  
 cgaagatta cgtagcacc 20  
 45 <210> 183  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 50 <223> Antisense Oligonucleotide  
 <400> 183  
 gccagccaa ctatacttg 20  
 55

EP 2 336 318 A1

5  
 <210> 184  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 184  
 cctggagcaa gtttcattga 20

10  
 <210> 185  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 185  
 tggacagacc aggcagaca 20

15  
 <210> 186  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 186  
 a\_gtgtactt ccggaggtc 20

20  
 <210> 187  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 187  
 tcttcaagat gaacctcag 20

25  
 <210> 188  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 188  
 tcagcaaggc ttgccctca 20

30  
 <210> 189  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 189  
 ctgcttcctt tctggaatg 20

35  
 <210> 190  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

40  
 <210> 190  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

45  
 <210> 190  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

50  
 <210> 190  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

55

EP 2 336 318 A1

5 <223> Antisense Oligonucleotide  
 <400> 190  
 tgccacattg ccctccctcg 20  
 <210> 191  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 191  
 gctgatcaga gttgacaagg 20  
 <210> 192  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 192  
 laclgacagy aclygclgc 20  
 <210> 193  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 193  
 catggtttct gccacatgc 20  
 <210> 194  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 194  
 gatgtggatt tgggtctctc 20  
 <210> 195  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 195  
 tgactgcttc atcactgagg 20  
 <210> 196  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 196  
 ggtaggtgac cacactctatc 20  
 <210> 197  
 <211> 20

55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 197  
 tcgcagctgc tgtcctgagg 20  
  
 <210> 198  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 198  
 Lccaalgc ccgcagaac 20  
  
 <210> 199  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 20  
  
 <400> 199  
 gatcatcagt gatcgtttg 20  
  
 <210> 200  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 200  
 agcctggatg gcagctttc 20  
  
 <210> 201  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 35  
  
 <400> 201  
 gctgaagaa gaacctcctg 20  
  
 <210> 202  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 202  
 tatctgctg tgaaggactc 20  
  
 <210> 203  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 50  
  
 <400> 203  
  
 55

EP 2 336 318 A1

	ctgagttcas gata-tggca	20
5	<210> 204 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 204 ctccaagcc aatc-cgatg	20
	<210> 205 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 205 tgcaactgta atccagctcc	20
20	<210> 206 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 206 ccagttcagc ctgcatgttg	20
	<210> 207 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 207 ctagagaccs aatg-taatg-	20
35	<210> 208 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 208 cgllggagla agcgcclgag	20
	<210> 209 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 209 caqctotaat ctgc-tgtccc	20
50	<210> 210 <211> 20 <212> DNA <213> Artificial Sequence	
55		

EP 2 336 318 A1

	<220>	
	<223> Antisense Oligonucleotide	
5	<400> 210 c-gtcctctc tctcgagctc	20
	<210> 211 <211> 20 <212> DNA <213> Artificial Sequence	
10	<220>	
	<223> Antisense Oligonucleotide	
	<400> 211 caaggtcata ctctgccgat	20
15	<210> 212 <211> 20 <212> DNA <213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
20	<400> 212 g-atggaaat aacacccttg	20
	<210> 213 <211> 20 <212> DNA <213> Artificial Sequence	
25	<220>	
	<223> Antisense Oligonucleotide	
	<400> 213 taagctgtag caga-gagtc	20
30	<210> 214 <211> 20 <212> DNA <213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
35	<400> 214 tagatcclcg gagcalttc	20
	<210> 215 <211> 20 <212> DNA <213> Artificial Sequence	
40	<220>	
	<223> Antisense Oligonucleotide	
	<400> 215 g-ctagaaca cccaggagag	20
45	<210> 216 <211> 20 <212> DNA <213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
50	<400> 216 accacagagt cagccttca-	20
	<210> 217	

55

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 217  
 aagcagacat ctgtggtccc 20  
 <210> 218  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 218  
 ctctccattg agccggccag 20  
 <210> 219  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 219  
 cctgatattc agaacgcagc 20  
 <210> 220  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 220  
 cagtgcctaa gatgtcagca 20  
 <210> 221  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 221  
 agcaccagga gactacactc 20  
 <210> 222  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 222  
 cccatccaga ctgaattttg 20  
 <210> 223  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 55



EP 2 336 318 A1

	<400> 223	
	ggttctagcc gtaçttccc	20
5	<210> 224	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
10	<400> 224	
	aggttaccag ccacatgcag	20
	<210> 225	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Antisense Oligonucleotide	
	<400> 225	
	atgtgcatcg atggcatgg	20
20	<210> 226	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
25	<400> 226	
	ccagagagcg agttcccat	20
	<210> 227	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Antisense Oligonucleotide	
	<400> 227	
	ctagacacga gatgatgac	20
35	<210> 228	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
40	<400> 228	
	tccaagtcct ggctgtattc	20
	<210> 229	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Antisense Oligonucleotide	
	<400> 229	
	cgtccagtaa gctccacgcc	20
50	<210> 230	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
55		

EP 2 336 318 A1

<220>  
 <223> Antisense Oligonucleotide  
 5 <400> 230  
 tcaacggcat ctctcatctc 20  
 <210> 231  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 231  
 tgatagtgt catcaagact 20  
 <210> 232  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 232  
 gattctgatt tggacttag 20  
 <210> 233  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 233  
 ctctcgatta actcatggac 20  
 <210> 234  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 234  
 atacactgca actgaggcc 20  
 <210> 235  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 235  
 gcaagagtcc accaatcaga 20  
 <210> 236  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 236  
 agagcctgaa gactgacttc 20  
 <210> 236  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40 <220>  
 <223> Antisense Oligonucleotide  
 <400> 236  
 agagcctgaa gactgacttc 20  
 <210> 236  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45 <220>  
 <223> Antisense Oligonucleotide  
 <400> 236  
 agagcctgaa gactgacttc 20  
 <210> 236  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50 <220>  
 <223> Antisense Oligonucleotide  
 <400> 236  
 agagcctgaa gactgacttc 20  
 <210> 236  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 55 <220>  
 <223> Antisense Oligonucleotide  
 <400> 236  
 agagcctgaa gactgacttc 20  
 <210> 236  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

EP 2 336 318 A1

5  
 <210> 237  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 237  
 tccctcatct gagaatctgg 20

10  
 <210> 238  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 238  
 caqtqcatca atgacagatg 20

15  
 <210> 239  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 239  
 ccgaaccctt gacatctcc 20

20  
 <210> 240  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 240  
 gcctcactag caatagtcc 20

25  
 <210> 241  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 241  
 gacattggcc atggagagag 20

30  
 <210> 242  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 242  
 cagtctccta ccaatgctgg 20

35  
 <210> 243  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

40  
 <210> 242  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 242  
 cagtctccta ccaatgctgg 20

45  
 <210> 243  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

50  
 <210> 243  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

55

EP 2 336 318 A1

5 <400> 243  
 lclgcactga aglccacggtg 20

<210> 244  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Antisense Oligonucleotide

<400> 244  
 tcccqgacct tcaactcaat 20

<210> 245  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

15 <220>  
 <223> Antisense Oligonucleotide

<400> 245  
 gcaggctccag ttcaatgtg 20

20 <210> 246  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

25 <400> 246  
 gccatccttc tgagtcaga 20

<210> 247  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> Antisense Oligonucleotide

<400> 247  
 gcctcagtct gcttcgcacc 20

35 <210> 248  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

40 <400> 248  
 ccccgcaggt cccggtggca 20

<210> 249  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

45 <220>  
 <223> Antisense Oligonucleotide

<400> 249  
 cagcccgca ggtcccggtg 20

50 <210> 250  
 <211> 20  
 <212> DNA

55

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
 <400> 250  
 caaccgagaa gggcaactcag 20  
 <210> 251  
 <211> 20  
 <212> DNA  
 10 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 251  
 cctcagcggc agcaaccgag 20  
 15 <210> 252  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 20 <223> Antisense Oligonucleotide  
 <400> 252  
 tctctcagcgg cagcaaccga 20  
 <210> 253  
 <211> 20  
 <212> DNA  
 25 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 253  
 ctcctcagcg gcagcaaccg 20  
 30 <210> 254  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 35 <223> Antisense Oligonucleotide  
 <400> 254  
 ggctcctcag cggcagcaac 20  
 <210> 255  
 <211> 20  
 <212> DNA  
 40 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 255  
 ggcgggctcc tcagcggcag 20  
 45 <210> 256  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 50 <223> Antisense Oligonucleotide  
 <400> 256  
 ggtccatcgc cagcgcggc 20  
 55

EP 2 336 318 A1

5  
 <210> 257  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

<400> 257  
 ggcggtcca tggcagctg 20

10  
 <210> 258  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

15  
 <400> 258  
 tagaggatga tagtaagttc 20

<210> 259  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

20  
 <220>  
 <223> Antisense Oligonucleotide

<400> 259  
 aaatgaagat llctllaaa 20

25  
 <210> 260  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30  
 <220>  
 <223> Antisense Oligonucleotide

<400> 260  
 tatgtgaaaq ttcaattga 20

35  
 <210> 261  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

40  
 <400> 261  
 aataggcag tttgaattt 20

<210> 262  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Antisense Oligonucleotide

<400> 262  
 gctcactgta tggttttatc 20

<210> 263  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

50  
 <220>

55

EP 2 336 318 A1

<223> Antisense Oligonucleotide  
 <400> 263  
 ggctcactgt atggctttat 20  
 5  
 <210> 264  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 10  
 <400> 264  
 ggctggctca ctgtatggt 20  
 <210> 265  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 265  
 aggctggctc actgcatggc 20  
 20  
 <210> 266  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 25  
 <400> 266  
 aaagctggct cactgtatcg 20  
 <210> 267  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 267  
 ctactgcaag gctggctcac 20  
 35  
 <210> 268  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 40  
 <400> 268  
 actgcctact gcaaggctgg 20  
 <210> 269  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 269  
 tgcttatagt ctactgcta 20  
 50  
 <210> 270  
 <211> 20

55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 270  
 tctgttat agtctactgc 20  
  
 <210> 271  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 271  
 LtggLgcag glccagLlca 20  
  
 <210> 272  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 272  
 caqctttggt gcaqgtccag 20  
  
 <210> 273  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 273  
 gccagctttg gtgcaggtcc 20  
  
 <210> 274  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 274  
 tgggtgccagc tttcgtgcag 20  
  
 <210> 275  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 275  
 gccctggtgc cagccttggc 20  
  
 <210> 276  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 276  
  
 40  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 276  
  
 45  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 276  
  
 50  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 276  
  
 55



EP 2 336 318 A1

	gagttcagag acctccgag	20
5	<210> 277 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 277 aaatgccatc ctccgagtc	20
	<210> 278 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 278 aaaaatgcc a tctctgag	20
20	<210> 279 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 279 aaaataactc agatcctgac	20
	<210> 280 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 280 agcaaaatas ctcagatccc	20
35	<210> 281 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 281 aglllagcas aalaaclcag	20
	<210> 282 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 282 tcccccaagt ttaccaaaaac	20
50	<210> 283 <211> 20 <212> DNA <213> Artificial Sequence	
55		

EP 2 336 318 A1

	<220> <223> Antisense Oligonucleotide	
5	<400> 283 tccctcctcc cccaagtta	20
	<210> 284 <211> 20 <212> DNA <213> Artificial Sequence	
10	<220> <223> Antisense Oligonucleotide	
	<400> 284 agactccatt tatttgttcc	20
15	<210> 285 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
20	<400> 285 ctctgcttg agttacaaac	20
	<210> 286 <211> 20 <212> DNA <213> Artificial Sequence	
25	<220> <223> Antisense Oligonucleotide	
	<400> 286 accttctgct tgaattacaa	20
30	<210> 287 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
35	<400> 287 gcacclclg clcgagllac	20
	<210> 288 <211> 20 <212> DNA <213> Artificial Sequence	
40	<220> <223> Antisense Oligonucleotide	
	<400> 288 tcgcacottc tgcttgaatt	20
45	<210> 289 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
50	<400> 289 cttcgcacct tctccttgag	20
	<210> 290	
55		

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 290  
 tgcttgcac ctccgcttg 20  
 <210> 291  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 291  
 tctgcttcgc acctctctgc 20  
 <210> 292  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 292  
 agtctgcttc gcacctctg 20  
 <210> 293  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 293  
 tcagtctgct tcgcaccttc 20  
 <210> 294  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 294  
 cctcagtctg cttcgcacct 20  
 <210> 295  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 295  
 agcctcagtc tgcttcgcac 20  
 <210> 296  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 55

EP 2 336 318 A1

	<400> 296	
	gtagcctcag tctcttcgc	20
5	<210> 297 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 297	
	tgtagcctc agtctgcttc	20
	<210> 298 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 298	
	catggtagcc tcagtctgc	20
20	<210> 299 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 299	
	gcatggtag cctcagtctg	20
	<210> 300 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 300	
	atgcatggt agcctcagtc	20
35	<210> 301 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 301	
	gaatgcatg gtagcctcag	20
	<210> 302 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 302	
	tgaatgtcc tgtagcctc	20
50	<210> 303 <211> 20 <212> DNA <213> Artificial Sequence	
55		

EP 2 336 318 A1

<220>  
 <223> Antisense Oligonucleotide  
 5 <400> 303  
 a-ttgaatgt catggtagcc 20  
 <210> 304  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 304  
 a-atttgaat gtca-ggtag 20  
 <210> 305  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 305  
 cagccacatg cagc-ttcagg 20  
 <210> 306  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 306  
 accagccaca tgcagc-ttca 20  
 <210> 307  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 307  
 t-accagcca catgcagc-t 20  
 <210> 308  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 308  
 ggttaccagc caca-gccagc 20  
 <210> 309  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 309  
 laggtlaccg gccacal-gca 20  
 <210> 309  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40 <220>  
 <223> Antisense Oligonucleotide  
 <400> 308  
 ggttaccagc caca-gccagc 20  
 <210> 309  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45 <220>  
 <223> Antisense Oligonucleotide  
 <400> 309  
 laggtlaccg gccacal-gca 20  
 <210> 309  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50 <220>  
 <223> Antisense Oligonucleotide  
 <400> 309  
 laggtlaccg gccacal-gca 20  
 <210> 309  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 55 <220>  
 <223> Antisense Oligonucleotide  
 <400> 309  
 laggtlaccg gccacal-gca 20  
 <210> 309  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

EP 2 336 318 A1

<210> 3'0  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 3'0  
 tttaggttac cagccacatg 20  
 10  
 <210> 3'1  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 15  
 <400> 3'1  
 cttttaqgtt accaqccaca 20  
 <210> 3'2  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 3'2  
 tccttttagg ttaccagcca 20  
 25  
 <210> 3'3  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 30  
 <400> 3'3  
 gtccttttta ggttaccagc 20  
 <210> 3'4  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 3'4  
 aggctccttt taggtacca 20  
 40  
 <210> 3'5  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 45  
 <400> 3'5  
 gtaggctcct ttaggttac 20  
 <210> 3'6  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 55

EP 2 336 318 A1

<400> 316  
 lgglaggcgc cllllaggll 20  
 5  
 <210> 317  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 10  
 <400> 317  
 tttgqtaggc tccttttaag 20  
 <210> 318  
 <211> 13993  
 <212> DNA  
 <213> H. sapiens  
 15  
 <220>  
 <221> CDS  
 <222> (1).. (13692)  
 <400> 318  
 20  
 a\_g gac cgg ccg agg ccc gcg ctg ctg gcg ctg ctg gcg ctg cct ggg 48  
 Met Asp Pro Pro Arg Pro Ala Leu Leu Ala Leu Leu Ala Leu Pro Ala  
 1 5 10 15  
 ctg ctg ctg ctg ctg ctg gcg gcc gcc agg gcc gaa gag gaa atg ctg 96  
 Leu Leu Leu Leu Leu Leu Ala Gly Ala Arg Ala Glu Gln Glu Met Leu  
 20 25 30  
 25  
 gaa aat gtc agc ctg gtc tgt cca aaa gat gcg acc cga ttc sag cac 144  
 Glu Asn Val Ser Leu Val Cys Pro Lys Asp Ala Thr Arg Phe Lys His  
 35 40 45  
 ctg cgg aag tac aca tac aac tat gag gct gag agt tcc agt gga gtc 192  
 Leu Arg Lys Tyr Thr Tyr Asn Tyr Glu Ala Glu Ser Ser Ser Gly Val  
 50 55 60  
 30  
 cct ggg act gct gat tca aga agt gcc acc agg atc aac tgc aag gtt 240  
 Pro Gly Thr Ala Asp Ser Arg Ser Ala Thr Arg Ile Asn Cys Lys Val  
 65 70 75 80  
 35  
 gag ctg gag gtt ccc cag ctc tgc agc ttc atc ctg aag acc agc cag 288  
 Glu Leu Glu Val Pro Gln Leu Cys Ser Phe Ile Leu Lys Thr Ser Gln  
 85 90 95  
 40  
 tgc acc ctg aaa gag gtg tat gcc ttc aac cct gag gcc aaa gcc ttg 336  
 Cys Thr Leu Lys Glu Val Tyr Gly Phe Asn Pro Glu Gly Lys Ala Leu  
 100 105 110  
 ctg aag aaa acc aag aac tct gag gag ttt gct gca gcc atg tcc agg 384  
 Leu Lys Lys Thr Lys Asn Ser Glu Glu Phe Ala Ala Ala Met Ser Arg  
 115 120 125  
 45  
 tat gag ctg aag ctg gcc att cca gaa ggg aag cag gtt ttc ctg tac 432  
 Tyr Glu Leu Lys Leu Ala Ile Pro Glu Gly Lys Gln Val Phe Leu Tyr  
 130 135 140  
 ccg gag asa gat gaa cct act tac atc ctg aac atc aag agg gcc atc 480  
 Pro Glu Lys Asp Glu Pro Thr Tyr Ile Leu Asn Ile Lys Arg Gly Ile  
 145 150 155 160  
 50  
 att tct gcc ctg ctg gtt ccc cca gag aca gaa gaa gcc aag caa gtg 528  
 Ile Ser Ala Leu Leu Val Pro Pro Glu Thr Glu Glu Ala Lys Gln Val  
 165 170 175  
 55  
 t\_g tll ctg gal acc glg tal gga aac tgc tcc act cac tll acc glc 576  
 Leu Phe Leu Asp Thr Val Tyr Gly Asn Cys Ser Thr His Phe Thr Val

EP 2 336 318 A1

	180	185	190	
5	aag acg agg aag ggc aal glg gca aca gaa ala lcc acl gaa aga gac Lys Trp Arg Lys Gly Asn Val Ala Trp Glu Ile Ser Thr Glu Arg Asp 195 200 205	624		
	ctg qgg cag tgt gat cgc ttc aag ccc atc cgc aca ggc atc agc coa Leu Gly Gln Cys Asp Arg Phe Lys Pro Ile Arg Thr Gly Ile Ser Pro 210 215 220	672		
10	ctt gct ctc atc aaa ggc atg acc cgc ccc ttg tca act ctg atc agc Leu Ala Leu Ile Lys Gly Met Thr Arg Pro Leu Ser Thr Leu Ile Ser 225 230 235 240	720		
	agc agc cag tcc tgt cag tac aca ctg gac gct aag agg aag cat gtg Ser Ser Gln Ser Cys Gln Tyr Thr Leu Asp Ala Lys Arg Lys His Val 245 250 255	768		
15	gca gaa gcc atc lgc aag gag caa cac ctc ltc ctg ccl ltc lcc lac Ala Glu Ala Ile Cys Lys Glu Gln His Leu Phe Leu Pro Phe Ser Tyr 260 265 270	816		
	aag aat aag tat ggg atg gta gca caa gtg aca cag act ttg aaa ctt Lys Asn Lys Tyr Gly Met Val Ala Gln Val Thr Gln Thr Leu Lys Leu 275 280 285	864		
20	gaa gac aca cca aag atc aac agc cgc ttc ttt ggt gaa ggt act aag Glu Asp Thr Pro Lys Ile Asn Ser Arg Phe Phe Gly Glu Gly Thr Lys 290 295 300	912		
	aag atg ggc ctc gca ttt gag agc acc aaa tcc aca tca cct cca aag Lys Met Gly Leu Ala Phe Clu Ser Thr Lys Ser Thr Ser Pro Pro Lys 305 310 315 320	960		
25	cag gcc gaa gct gtt ttg aag act ctc cag gaa ctg aaa aaa cta acc Gln Ala Glu Ala Val Leu Lys Thr Leu Gln Glu Leu Lys Lys Leu Thr 325 330 335	1008		
	atc tot gag caa aat atc cag aga gct aat ctc ttc aat aag ctg gtt Ile Ser Glu Gln Asn Ile Gln Arg Ala Asn Leu Phe Asn Lys Leu Val 340 345 350	1056		
30	act gag ctg aga ggc ctc agt gat gaa gca gtc aca tct ctc ttg cca Thr Glu Leu Arg Gly Leu Ser Asp Glu Ala Val Thr Ser Leu Leu Pro 355 360 365	1104		
	cag ctg att gag gtg tcc agc ccc atc act tta caa gcc ttg gtt cag Gln Leu Ile Glu Val Ser Ser Pro Ile Thr Leu Gln Ala Leu Val Gln 370 375 380	1152		
35	tgt gga cag cct cag tgc tcc act cac atc ctc cag tgg ctg aaa cgt Cys Gly Gln Pro Gln Cys Ser Thr His Ile Leu Gln Trp Leu Lys Arg 385 390 395 400	1200		
	gtg cat gcc aac ccc ctt ctg ata gat gtg gtc acc tac ctg gtg gcc Val His Ala Asn Pro Leu Leu Ile Asp Val Val Thr Tyr Leu Val Ala 405 410 415	1248		
40	ctg atc ccc gag ccc tca gca cag cag ctg cga gag atc ttc aac atg Leu Ile Pro Glu Pro Ser Ala Gln Gln Leu Arg Glu Ile Phe Asn Met 420 425 430	1296		
45	gcg agg gat cag cgc agc cga gcc acc ttg tat gcg ctg agc cac gcg Ala Arg Asp Gln Arg Ser Arg Ala Thr Leu Tyr Ala Leu Ser His Ala 435 440 445	1344		
	ctc aac aac tat cat aag aca aac cct aca ggg acc cag gag ctg ctg Val Asn Asn Tyr His Lys Thr Asn Pro Thr Gly Thr Gln Glu Leu Leu 450 455 460	1392		
50	gac att gct aat tac ctg atg gaa cag att caa gat gac tgc act gcg Asp Ile Ala Asn Tyr Leu Met Glu Gln Ile Gln Asp Asp Cys Thr Gly 465 470 475 480	1440		
55				



EP 2 336 318 A1

gat gaa gat tac acc tat ttg att ctg cgg gtc att gga aat atg gcc 1488  
 Asp Glu Asp Tyr Thr Tyr Leu Ile Leu Arg Val Ile Gly Asn Met Gly  
 485 490 495

5  
 caa acc atg gag cag tta act cca gaa ctc aag tct tca atc atg aaa 1536  
 Gln Thr Met Glu Gln Leu Thr Pro Glu Leu Lys Ser Ser Ile Leu Lys  
 500 505 510

tgt gtc caa agt aca aag cca tca ctg atg atc cag aaa gct gcc atc 1584  
 Cys Val Gln Ser Thr Lys Pro Ser Leu Met Ile Gln Lys Ala Ala Ile  
 515 520 525

10  
 caq gct ctg cgg aaa atg gaq cct aaa gac aag gac cag gag gtt ctt 1632  
 Gln Ala Leu Arg Lys Met Glu Pro Lys Asp Lys Asp Gln Glu Val Leu  
 530 535 540

ctt cag act ttc ctt gat gat gct tct cgg gga gat aag cga atg gct 1680  
 Leu Gln Thr Phe Leu Asp Asp Ala Ser Pro Gly Asp Lys Arg Leu Ala  
 545 550 555 560

15  
 gcc tat ctt atg ttg atg agg agt cct tca cag gca gat att aac aaa 1728  
 Ala Tyr Leu Met Leu Met Arg Ser Pro Ser Gln Ala Asp Ile Asn Lys  
 565 570 575

20  
 aul glc caa all cla cca lgg gaa cag aal gag caa glg aag aac lll 1776  
 Ile Val Gln Ile Leu Pro Trp Glu Gln Asn Glu Gln Val Lys Asn Phe  
 580 585 590

ctg gct tcc cat att gcc aat atc ttg aac tca gaa gaa ttg gat atc 1824  
 Val Ala Ser His Ile Ala Asn Ile Leu Asn Ser Glu Glu Leu Asp Ile  
 595 600 605

25  
 caa gat ctg aaa aag tta gtg aaa gaa gct ctg aaa gaa tct caa ctt 1872  
 Gln Asp Leu Lys Lys Leu Val Lys Glu Ala Leu Lys Glu Ser Gln Leu  
 610 615 620

cca act gtc atg gac ttc aga aaa ttc tct cgg aac tat caa ctc tac 1920  
 Pro Thr Val Met Asp Phe Arg Lys Phe Ser Arg Asn Tyr Gln Leu Tyr  
 625 630 635 640

30  
 aaa tct gtt tct ctt cca tca ctt gac cca gcc tca gcc aaa ata gaa 1968  
 Lys Ser Val Ser Leu Pro Ser Leu Asp Pro Ala Ser Ala Lys Ile Glu  
 645 650 655

ggg aat ctt ata ttt gat cca aat aac tac ctt cct aaa gaa agc atg 2016  
 Gly Asn Leu Ile Phe Asp Pro Asn Asn Tyr Leu Pro Lys Glu Ser Met  
 660 665 670

35  
 ctg aaa act acc ctc act gcc ttt gga ttt gct tca gct gac ctc atc 2064  
 Leu Lys Thr Thr Leu Thr Ala Phe Gly Phe Ala Ser Ala Asp Leu Ile  
 675 680 685

gag att ggc ttg gaa gga aaa ggc ttt gag cca aca ttg gag gct cct 2112  
 Glu Ile Gly Leu Glu Gly Lys Gly Phe Glu Pro Thr Leu Glu Ala Pro  
 690 695 700

40  
 ttt ggg aag caa gga ttt ttc cca gac agt gtc aac aaa gct ttg tac 2160  
 Phe Gly Lys Gln Gly Phe Phe Pro Asp Ser Val Asn Lys Ala Leu Tyr  
 705 710 715 720

45  
 tgg gtt ast ggt caa gtt cct gat ggt gtc tct aag gtc tta gtg gac 2208  
 Trp Val Asn Gly Gln Val Pro Asp Gly Val Ser Lys Val Leu Val Asp  
 725 730 735

cac ttt ggc tat acc aaa gat gat aaa cat gag cag gat atg gla aat 2256  
 His Phe Gly Tyr Thr Lys Asp Asp Lys His Glu Gln Asp Met Val Asn  
 740 745 750

50  
 gga ala atg ctc agt gll gag aag ctg all aaa gal llg aaa tcc aaa 2304  
 Gly Ile Met Leu Ser Val Glu Lys Leu Ile Lys Asp Leu Lys Ser Lys

55

EP 2 336 318 A1

	755	760	765	
5	gaa glc cgg gaa gcc aga gcc lac ctc cgc atc ttg gga gag gag clc Glu Val Pro Glu Ala Arg Ala Tyr Leu Arg Ile Leu G_y Glu Glu Leu 770 775 780			2352
	cgt ttt gcc agt ctc cat gac ctc cca ctc ctg gga aag ctg ctt ctg Gly Phe Ala Ser Leu His Asp Leu Arg Leu Leu Gly Lys Leu Leu Leu 785 790 795 800			2400
10	atg ggt gcc cgc act ctg cag ggg atc ccc cag atg att gga gag gtc Met Gly Ala Arg Thr Leu Gln Gly Ile Pro Gln Met Ile Gly Glu Val 805 810 815			2448
	atc agg aag gcc tca aag aat gac ttt ttt ctt cac tac atc ttc atg Ile Arg Lys Gly Ser Lys Asn Asp Phe Phe Leu His Tyr Ile Phe Met 820 825 830			2496
15	gag aal gcc lll gaa ctc ccc acL gga gcl gga lla cag ttg gaa ala Glu Asn Ala Phe Glu Leu Pro Thr Gly Ala Gly Leu Gln Leu Gln Ile 835 840 845			2544
	tct tca tct gga gtc att gct ccc gga gcc aag gct gga gta aaa ctg Ser Ser Ser Gly Val Ile Ala Pro Gly Ala Lys Ala Gly Val Lys Leu 850 855 860			2592
20	gaa gta gcc aac atg cag gct gaa ctg gtg gca aaa ccc tcc ctg tcc Glu Val Ala Asn Met Gln Ala Glu Leu Val Ala Lys Pro Ser Val Ser 865 870 875 880			2640
	gtg gag ttt gtg aca aat atg gcc atc atc att cgg gac ttc gct agg Val Glu Phe Val Thr Asn Met Gly Ile Ile Ile Pro Asp Phe Ala Arg 885 890 895			2688
25	agt ggg gtc caa atg aac acc aac ttc ttc cac gag tog ggt ctg gag Ser Gly Val Glr Met Asn Thr Asn Phe Phe His Glu Ser Gly Leu Glu 900 905 910			2736
	gct cat gtt gcc cta aaa gct ggg aag ctg aag ttt atc att cct tcc Ala His Val Ala Leu Lys Ala Gly Lys Leu Lys Phe Ile Ile Pro Ser 915 920 925			2784
	cca aag aga cca gtc aag ctg ctc agt gga gcc aac aca tta cat ttg Pro Lys Arg Pro Val Lys Leu Leu Ser Gly Gly Asn Thr Leu His Leu 930 935 940			2832
35	gtc tct acc acc aaa acg gag gtc atc cca cct ctc att gag aac agg Val Ser Thr Thr Lys Thr Clu Val Ile Pro Pro Leu Ile Clu Asn Arg 945 950 955 960			2880
	caq tcc tgg tca gtt tgc aag caa gtc ttt cct gcc ctg aat tac tcc Gln Ser Trp Ser Val Cys Lys Gln Val Phe Pro Gly Leu Asn Tyr Cys 965 970 975			2928
40	acc tca gcc gct tac tcc aac gcc agc tcc aca gac tcc gcc tcc tac Thr Ser Gly Ala Tyr Ser Asn Ala Ser Ser Thr Asp Ser Ala Ser Tyr 980 985 990			2976
	tat cgg ctg acc ggg gac acc aga tta gag ctg gaa ctg agg cct aca Tyr Pro Leu Thr Gly Asp Thr Arg Leu Glu Leu Glu Leu Arg Pro Thr 995 1000 1005			3024
45	gga gag att gag cag tat tct gtc agc gca acc tat gag ctc cag aga Gly Glu Ile Glu Gln Tyr Ser Val Ser Ala Thr Tyr Glu Leu Gln Arg 1010 1015 1020			3072
	caq gac aga gcc ttg gtg gat acc ctg aag ttt gta act caa cca gaa Glu Asp Arg Ala Leu Val Asp Thr Leu Lys Phe Val Thr Gln Ala Glu 1025 1030 1035 1040			3120
50	ggc gcg aag caa act gag got acc atg aca ttc aaa tat aat cgg cag Gly Ala Lys Gln Thr Glu Ala Thr Met Thr Phe Lys Tyr Asn Arg Gln 1045 1050 1055			3168
55				

EP 2 336 318 A1

5 agt atg acc ttg tcc agt gaa gtc caa att cag gat ttt gat gtt gcc 3216  
 Ser Met Thr Leu Ser Ser Glu Val Gln Ile Pro Asp Phe Asp Val Asp  
 1060 1065 1070

10 ctc gga aca atc ctc aga gtt aat gat gaa tct act gag ggc aaa aag 3264  
 Leu Gly Thr Ile Leu Arg Val Asn Asp Glu Ser Thr Glu Gly Lys Thr  
 1075 1080 1085

15 tct tac aga ctc acc ctg gac att cag aac aag aaa att act gag gtc 3312  
 Ser Tyr Arg Leu Thr Leu Asp Ile Gln Asn Lys Lys Ile Thr Glu Val  
 1090 1095 1100

20 gcc ctc atg ggc cac cta agt tgt qac aca aag gaa gaa aqa aaa atc 3360  
 Ala Leu Met Gly His Leu Ser Cys Asp Thr Lys Glu Glu Arg Lys Ile  
 1105 1110 1115 1120

25 aag ggt gtt att tcc ata ccc cgt ttg caa gcc gaa gcc aga agt gag 3408  
 Lys Gly Val Ile Ser Ile Pro Arg Leu Gln Ala Glu Ala Arg Ser Glu  
 1125 1130 1135

30 atc ctc gcc cac tgg tgg cct gcc aaa ctg ctt ctc caa atg gac tca 3456  
 Ile Leu Ala His Trp Ser Pro Ala Lys Leu Leu Leu Gln Met Asp Ser  
 1140 1145 1150

35 tct gct aca gct tat ggc tcc aca glu tcc aag agg gtg gca tgg caa 3504  
 Ser Ala Thr Ala Tyr Gly Ser Thr Val Ser Lys Arg Val Ala Trp His  
 1155 1160 1165

40 tat gat gaa gag aag att gaa ttt gaa tgg aac aca gcc acc aat gta 3552  
 Tyr Asp Glu Glu Lys Ile Glu Phe Glu Trp Asn Thr Gly Thr Asn Val  
 1170 1175 1180

45 gat acc aca aaa atg act tcc aat ttc cct gtg gat ctc tcc gat tat 3600  
 Asp Thr Lys Lys Met Thr Ser Asn Phe Pro Val Asp Leu Ser Asp Tyr  
 1185 1190 1195 1200

50 cct aag agc ttg cat atg tat gct aac aga ctc ctg gat cac aga gtc 3648  
 Pro Lys Ser Leu His Met Tyr Ala Asn Arg Leu Leu Asp His Arg Val  
 1205 1210 1215

55 cct caa aca gac atg act ttc cgg cac gtg ggt tcc aaa tta ata gtt 3696  
 Pro Gln Thr Asp Met Thr Phe Arg His Val Gly Ser Lys Leu Ile Val  
 1220 1225 1230

60 gca atg agc tca tgg ctt cag aag gca tct ggg agt ctt cct tat acc 3744  
 Ala Met Ser Ser Trp Leu Gln Lys Ala Ser Gly Ser Leu Pro Tyr Thr  
 1235 1240 1245

65 cag act ttg caa gac cac ctc aat agc ctg aag gag ttc aac ctc cag 3792  
 Gln Thr Leu Gln Asp His Leu Asn Ser Leu Lys Glu Phe Asn Leu Gln  
 1250 1255 1260

70 aac atg gga ttg cca gac tcc cac atc cca gaa aac ctc ttc tta aaa 3840  
 Asn Met Gly Leu Pro Asp Ser His Ile Pro Glu Asn Leu Phe Leu Lys  
 1265 1270 1275 1280

75 agc gat ggc cgc gtc aaa tat acc ttg aac aag aac agt ttg aaa att 3888  
 Ser Asp Gly Arg Val Lys Tyr Thr Leu Asn Lys Asn Ser Leu Lys Ile  
 1285 1290 1295

80 gag att cct ttg cct ttt ggt ggc aaa tcc tcc aga gat cta aag atg 3936  
 Glu Ile Pro Leu Pro Phe Gly Gly Lys Ser Ser Arg Asp Leu Lys Met  
 1300 1305 1310

85 tta gag act gtt agg aca cca gcc ctc cac ttc aag tct gtg gga ttc 3984  
 Leu Glu Thr Val Arg Thr Pro Ala Leu His Phe Lys Ser Val Gly Phe  
 1315 1320 1325

90 cal ctg cca tcl cga gag ttc caa gtc ccl act tll acc all ccc aag 4032  
 His Leu Pro Ser Arg Glu Phe Gln Val Pro Thr Phe Thr Ile Pro Lys

EP 2 336 318 A1

	1330		1335		1340																	
5	Llg	Lal	caa	ctg	caa	glg	cct	ctc	ctg	ggl	gll	cta	gac	ctc	tcc	acg						4080
	Leu	Tyr	Gln	Leu	Gln	Val	Pro	Leu	Leu	Gly	Val	Leu	Asp	Leu	Ser	Trp						1345
						1350					1355											1360
	aat	qtc	tac	agc	aac	ttg	tac	aac	tcg	tcc	gcc	tcc	tac	agt	qgt	qcc						4128
	Asn	Val	Tyr	Ser	Asn	Leu	Tyr	Asn	Trp	Ser	Ala	Ser	Tyr	Ser	Gly	Gly						1365
					1365					1370					1375							
10	aac	acc	agc	aca	gac	cat	ttc	agc	ctc	cgg	gct	cgt	tac	cac	atg	aag						4176
	Asn	Trp	Ser	Thr	Asp	His	Phe	Ser	Leu	Arg	Ala	Arg	Tyr	His	Met	Lys						1380
					1380					1385					1390							
	gct	gac	tct	gtg	gtt	gac	ctg	ctt	tcc	tac	aat	gtg	caa	gga	tct	gga						4224
	Ala	Asp	Ser	Val	Val	Asp	Leu	Leu	Ser	Tyr	Asn	Val	Gln	Gly	Ser	Gly						1395
									1400					1405								
15	gaa	aca	aca	Lal	gac	cac	aag	aal	acg	lcc	aca	cta	tca	tgl	gal	ggg						4272
	Glu	Trp	Thr	Tyr	Asp	His	Lys	Asn	Thr	Phe	Thr	Leu	Ser	Cys	Asp	Gly						1410
							1415						1420									
	tct	cta	cgc	cac	aaa	ttt	cta	gat	tcg	aat	atc	aaa	ttc	agt	cat	gta						4320
	Ser	Leu	Arg	His	Lys	Phe	Leu	Asp	Ser	Asn	Ile	Lys	Phe	Ser	His	Val						1425
						1430					1435					1440						
20	gaa	aaa	ctt	gga	aac	aac	cca	gtc	tca	aaa	ggg	tta	cta	ata	tcc	gat						4368
	Glu	Lys	Leu	Gly	Asn	Asn	Pro	Val	Ser	Lys	Gly	Leu	Leu	Ile	Phe	Asp						1445
					1445					1450					1455							
	gca	tct	agt	tcc	tgg	gga	cca	cag	atg	tct	gct	tca	ggt	cat	tgg	gac						4416
	Ala	Ser	Ser	Ser	Trp	Gly	Pro	Gln	Met	Ser	Ala	Ser	Val	His	Leu	Asp						1460
					1460				1465						1470							
25	tcc	aaa	asg	aaa	cag	cat	ttg	ttt	gtc	aaa	gaa	gtc	aag	att	gat	ggg						4464
	Ser	Lys	Lys	Lys	Gln	His	Leu	Phe	Val	Lys	Glu	Val	Lys	Ile	Asp	Gly						1475
							1480						1485									
	cag	ttc	aga	gtc	tct	tcg	ttc	tat	gct	aaa	ggc	aca	tat	ggc	ctg	tct						4512
	Gln	Phe	Arg	Val	Ser	Ser	Phe	Tyr	Ala	Lys	Gly	Thr	Tyr	Gly	Leu	Ser						1490
							1495						1500									
30	tgt	cag	agg	gat	cct	aac	act	ggc	cgg	ctc	aat	gga	gag	tcc	aac	ctg						4560
	Cys	Gln	Arg	Asp	Pro	Asn	Thr	Gly	Arg	Leu	Asn	Gly	Glu	Ser	Asn	Leu						1505
						1510					1515				1520							
	agg	ttt	aac	tcc	tcc	tac	ctc	caa	ggc	acc	aac	cag	ata	aca	gga	aga						4608
	Arg	Phe	Asn	Ser	Ser	Tyr	Leu	Gln	Gly	Thr	Asn	Gln	Ile	Thr	Gly	Arg						1525
										1530					1535							
35	tat	gaa	gat	gga	acc	ctc	tcc	ctc	acc	tcc	acc	tct	gat	ctg	caa	agt						4656
	Tyr	Glu	Asp	Gly	Thr	Leu	Ser	Leu	Thr	Ser	Thr	Ser	Asp	Leu	Gln	Ser						1540
									1545					1550								
40	ggc	atc	att	aaa	aat	act	gct	tcc	cta	aag	tat	gag	aac	tac	gag	ctg						4704
	Gly	Ile	Ile	Lys	Asn	Thr	Ala	Ser	Leu	Lys	Tyr	Glu	Asn	Tyr	Glu	Leu						1555
				1555					1560					1565								
	act	tta	asa	tct	gac	acc	aat	ggg	aag	tat	aag	aac	ttt	gcc	act	tct						4752
	Thr	Leu	Lys	Ser	Asp	Thr	Asn	Gly	Lys	Tyr	Lys	Asn	Phe	Ala	Thr	Ser						1570
							1575						1580									
45	aac	aag	atg	gat	atg	acc	ttc	tct	aag	caa	aat	gca	ctg	ctg	cgt	tct						4800
	Asn	Lys	Met	Asp	Met	Thr	Phe	Ser	Lys	Gln	Asn	Ala	Leu	Leu	Arg	Ser						1585
						1590					1595				1600							
	gaa	tat	cag	gct	gat	tac	gag	tca	ttg	agg	ttc	ttc	agc	ctg	ctt	tct						4848
	Glu	Tyr	Gln	Ala	Asp	Tyr	Glu	Ser	Leu	Arg	Phe	Phe	Ser	Leu	Leu	Ser						1605
									1610					1615								
50	gga	tca	cta	aat	tcc	cat	ggg	ctt	gag	tta	aat	gct	gac	atc	tta	ggc						4896
55																						

EP 2 336 318 A1

	Gly Ser Leu Asn Ser His Gly Leu Glu Leu Asn Ala Asp Ile Leu Gly	
	1620 1625 1630	
5	act gac aaa att aat agt ggt gct cac aag ggg aca cta agg att gcc Thr Asp Lys Ile Asn Ser Gly Ala His Lys Ala Thr Leu Arg Ile Gly	4944
	1635 1640 1645	
	caa gat gga ata tct acc agt gca acg acc aac ttg aag tgt agt ctc Gln Asp Gly Ile Ser Thr Ser Ala Thr Thr Asn Leu Lys Cys Ser Leu	4992
	1650 1655 1660	
10	ctg gtg ctg gag aat gag ctg aat gca gag ctt ggc ctc tct ggg gca Leu Val Leu Glu Asn Glu Leu Asn Ala Glu Leu Gly Leu Ser Gly Ala	5040
	1665 1670 1675 1680	
	tct atg aaa tta aca aca aat ggc cgc ttc agg gaa cac aat gca aaa Ser Met Lys Leu Thr Thr Asn Gly Arg Phe Arg Glu His Asn Ala Lys	5088
	1685 1690 1695	
15	ttc agt ctg gat ggg aaa gcc gcc ctc aca gag cta tca ctg gga agt Phe Ser Leu Asp Gly Lys Ala Ala Leu Thr Glu Leu Ser Leu Gly Ser	5136
	1700 1705 1710	
	gct tat cag gcc atg att ctg ggt gtc gac agc aaa aac att ttc aac Ala Tyr Gln Ala Met Ile Leu Gly Val Asp Ser Lys Asn Ile Phe Asn	5184
20	1715 1720 1725	
	ttc aag gtc agt caa gaa gga ctt aag ctc tca aat gac atg atg gcc Phe Lys Val Ser Gln Glu Gly Leu Lys Leu Ser Asn Asp Met Met Gly	5232
	1730 1735 1740	
	tca tat gct gaa atg aaa ttt gac cac aca aac agt ctg aac att gca Ser Tyr Ala Glu Met Lys Phe Asp His Thr Asn Ser Leu Asn Ile Ala	5280
25	1745 1750 1755 1760	
	ggc tta tca ctg gac ttc tct tca aaa ctt gac aac att tac agc tct Gly Leu Ser Leu Asp Phe Ser Ser Lys Leu Asp Asn Ile Tyr Ser Ser	5328
	1765 1770 1775	
	gac aag ttt tat aag caa act gtt aat tta cag cta cag ccc tat tct Asp Lys Phe Tyr Lys Gln Thr Val Asn Leu Gln Leu Gln Pro Tyr Ser	5376
30	1780 1785 1790	
	ctg gta act act tta aac agt gac ctg aaa tac aat gct ctg gat ctc Leu Val Thr Thr Leu Asn Ser Asp Leu Lys Tyr Asn Ala Leu Asp Leu	5424
	1795 1800 1805	
	acc aac act ggg aaa cta cgg cta gaa ccc ctg aag ctg cat ctg gct Thr Asn Asn Gly Lys Leu Arg Leu Glu Pro Leu Lys Leu His Val Ala	5472
35	1810 1815 1820	
	ggt aac cta aaa gga gcc tac caa aat aat gaa ata aaa cac atc tat Gly Asn Leu Lys Gly Ala Tyr Gln Asn Asn Glu Ile Lys His Ile Tyr	5520
	1825 1830 1835 1840	
40	gcc atc tct tct gct gcc tta tca gca agc tat aaa gca gac act gtl Ala Ile Ser Ser Ala Ala Leu Ser Ala Ser Tyr Lys Ala Asp Thr Val	5568
	1845 1850 1855	
	gct aag gtt cac ggt gtg gag ttt acc cat ggg ctc aac aca gac atc Ala Lys Val Gln Gly Val Glu Phe Ser His Gly Leu Asn Thr Asp Ile	5616
45	1860 1865 1870	
	gct ggg ctg gct tca gcc att gac atg agc aca aac tat aat tca gcc Ala Gly Leu Ala Ser Ala Ile Asp Met Ser Thr Asn Tyr Asn Ser Asp	5664
	1875 1880 1885	
	tca ctg cat ttc agc aat gtc ttc cgt tct gta atg gcc cgg ttt acc Ser Leu His Phe Ser Asn Val Phe Arg Ser Val Met Ala Pro Phe Thr	5712
50	1890 1895 1900	
	alg acc atc gal gca cal aca aal gcc aal ggg aaa ctc gct ctc leg Met Thr Ile Asp Ala His Thr Asn Gly Asn Gly Lys Leu Ala Leu Trp	5760
55		

EP 2 336 318 A1

	1905		1910		1915		1920	
5	gga gaa cal act	ggg cag clg	lal agc aaa	llc clg llg	aaa gca gaa		5808	
	Gly Glu His Thr	Gly Gln Leu Tyr Ser	Lys Phe Leu Leu	Lys Ala Glu				
		1925	1930	1935				
	oct ctg gca ttt act	ttc tct cat gat	tac aaa ggc	tcc aca agt cat		5856		
	Pro Leu Ala Phe Thr	Phe Ser His Asp Tyr	Lys Gly Ser Thr	Ser His				
		1940	1945	1950				
10	cat ctc gtg tct agg	aaa agc atc agt	gca gct ctt	gaa cac aaa gtc		5904		
	His Leu Val Ser Arg	Lys Ser Ile Ser	Ala Ala Leu Glu	His Lys Val				
		1955	1960	1965				
	agt gcc ctg ctt act	cca gct gag cag	aca ggc acc	tgg aaa ctc aag		5952		
	Ser Ala Leu Leu Thr	Pro Ala Glu Gln Thr	Gly Thr Trp Lys	Leu Lys				
		1970	1975	1980				
15	acc caa lll aac aac	aal gaa lac agc	cag gac llg gal	gcl lac aac		6000		
	Thr Gln Phe Asn Asn	Asn Glu Tyr Ser	Gln Asp Leu Asp	Ala Tyr Asn				
		1985	1990	1995	2000			
	act aaa gat aaa att	ggc gtg gag ctg	act gga cga	act ctg gct gac		6048		
	Thr Lys Asp Lys Ile	Gly Val Glu Leu Thr	Gly Arg Thr Leu	Ala Asp				
		2005	2010	2015				
20	cta act cta cta gac	ccc cca att aaa	gtg cca ctt tta	ctc agt gag		6096		
	Leu Thr Leu Leu Asp	Ser Pro Ile Lys	Val Pro Leu Leu	Leu Ser Glu				
		2020	2025	2030				
25	ccc atc aal alc aal	gal gcl lla gag	atg aga gal gcc	gll gag aag		6144		
	Pro Ile Asn Ile Asn	Asp Ala Leu Glu Met	Arg Asp Ala Val	Glu Lys				
		2035	2040	2045				
	ccc caa gaa ttt aca	att gtt gct ttc	gta aag tat gat	aaa aac caa		6192		
	Pro Gln Glu Phe Thr	Ile Val Ala Phe	Val Lys Tyr Asp	Lys Asn Gln				
		2050	2055	2060				
30	gat gtt cac tcc att	aac ctg cca ttc	ttt gag acc ttg	caa gaa tat		6240		
	Asp Val His Ser Ile	Asn Leu Pro Phe	Phe Glu Thr Leu	Gln Glu Tyr				
		2065	2070	2075	2080			
	ttt gag agg aat cga	caa acc att ata	gtt gta ctg gaa	aac gta cag		6288		
	Phe Glu Arg Asn Arg	Gln Thr Ile Ile	Val Val Leu Glu	Asn Val Gln				
		2085	2090	2095				
35	aga aac clg aag cac	alc aal all gal	caa lll gla aga	aaa lac aga		6336		
	Arg Asn Leu Lys His	Ile Asn Ile Asp	Gln Phe Val Arg	Lys Tyr Arg				
		2100	2105	2110				
	gca gcc ctg gga aaa	ctc cca cag caa	gct aat gat tat	ctg aat tca		6384		
	Ala Ala Leu Gly Lys	Leu Pro Gln Gln	Ala Asn Asp Tyr	Leu Asn Ser				
		2115	2120	2125				
40	ttc aat tgg gag aga	caa gtt tca cat	gcc aag gag aaa	ctg act gct		6432		
	Phe Asn Trp Glu Arg	Gln Val Ser His	Ala Lys Glu Lys	Leu Thr Ala				
		2130	2135	2140				
45	ctc aca asa aag tat	aga att aca gaa	aat gat ata caa	att gca tta		6480		
	Leu Thr Lys Lys Tyr	Arg Ile Thr Glu	Asn Asp Ile Gln	Ile Ala Leu				
		2145	2150	2155	2160			
	gat gat gcc aaa atc	aac ttt aat gaa	aaa cta tct caa	ctg cag aca		6528		
	Asp Asp Ala Lys Ile	Asn Phe Asn Glu	Lys Leu Ser Gln	Leu Gln Thr				
		2165	2170	2175				
50	tat atg ata caa ttt	gat cag tat atc	aaa gat agt tat	gat tta cat		6576		
	Tyr Met Ile Gln Phe	Asp Gln Tyr Ile	Lys Asp Ser Tyr	Asp Leu His				
		2180	2185	2190				
	gat ttg asa ata gct	att gct aat atc	att gat gaa atc	act gaa asa		6624		
55								

EP 2 336 318 A1

	Asp	Leu	Lys	Ile	Ala	Phe	Ala	Asn	Ile	Ile	Asp	Glu	Phe	Ile	Glu	Lys	
			2135					2200					2205				
5	tta	aaa	agt	ctt	gat	gag	cac	tat	cat	acc	cgt	gta	aat	tta	gta	aaa	6672
	Leu	Lys	Ser	Leu	Asp	Glu	His	Tyr	His	Thr	Arg	Val	Asn	Ser	Leu	Val	Lys
			2210				2215					2220					
	aca	atc	cat	gat	cta	cat	ttg	ttt	att	gaa	aat	att	gat	ttt	aac	aaa	6720
	Thr	Ile	His	Asp	Leu	His	Leu	Phe	Ile	Glu	Asn	Ile	Asp	Phe	Asn	Lys	2240
						2230					2235						
10	agt	gga	agt	agt	act	gca	toc	tgg	att	caa	aat	gtg	gat	act	aag	tac	6768
	Ser	Gly	Ser	Ser	Thr	Ala	Ser	Trp	Ile	Gln	Asn	Val	Asp	Thr	Lys	Tyr	
					2245					2250					2255		
	caa	atc	aga	atc	cag	ata	caa	gaa	aaa	ctg	cag	cag	ctt	aag	aga	cac	6816
	Gln	Ile	Arg	Ile	Gln	Ile	Gln	Glu	Lys	Leu	Gln	Gln	Leu	Lys	Arg	His	
				2260					2265					2270			
15	ata	cag	aat	ata	gac	atc	cag	cac	cta	gct	gga	aag	tta	aaa	caa	cac	6864
	Ile	Gln	Asn	Ile	Asp	Ile	Gln	His	Leu	Ala	Gly	Lys	Leu	Lys	Gln	His	
			2275					2280					2285				
	att	gag	gct	att	gat	gtt	aga	gtg	ctt	tta	gat	caa	ttg	gga	act	aca	6912
	Ile	Glu	Ala	Ile	Asp	Val	Arg	Val	Leu	Leu	Asp	Gln	Leu	Gly	Thr	Thr	
20			2290				2295				2300						
	att	tca	ttt	gaa	aga	ata	aat	gat	gtt	ctt	gag	cat	gtc	aaa	cac	ttt	6960
	Ile	Ser	Phe	Glu	Arg	Ile	Asn	Asp	Val	Leu	Glu	His	Val	Lys	His	Phe	
			2305			2310					2315				2320		
25	gct	ata	aat	ctt	att	ggg	gat	ttt	gaa	gta	gct	gag	aaa	atc	aat	gcc	7008
	Val	Ile	Asn	Leu	Ile	Gly	Asp	Phe	Glu	Val	Ala	Glu	Lys	Ile	Asn	Ala	
				2325					2330						2335		
	ttc	aga	gcc	aaa	gtc	cat	gag	tta	atc	gag	agg	tat	gaa	gta	gac	caa	7056
	Phe	Arg	Ala	Lys	Val	His	Glu	Leu	Ile	Glu	Arg	Tyr	Glu	Val	Asp	Gln	
			2340					2345					2350				
30	caa	atc	cag	gtt	tta	atg	gat	aaa	tta	gta	gag	ttg	gcc	cac	caa	tac	7104
	Gln	Ile	Gln	Val	Leu	Met	Asp	Lys	Leu	Val	Glu	Leu	Ala	His	Gln	Tyr	
			2355				2360						2365				
	aaq	ttg	aag	gac	act	att	cag	aaq	cta	agc	aat	gtc	cta	caa	caa	gtt	7152
	Lys	Leu	Lys	Glu	Thr	Phe	Gln	Lys	Leu	Ser	Asn	Val	Leu	Gln	Gln	Val	
			2370				2375					2380					
35	aag	ata	asa	gat	tac	ttt	gag	aaa	ttg	gtt	gga	ttt	att	gat	gat	gct	7200
	Lys	Ile	Lys	Asp	Tyr	Phe	Glu	Lys	Leu	Val	Gly	Phe	Ile	Asp	Asp	Ala	
			2385			2390					2395				2400		
	gtc	aag	aag	ctt	aat	gaa	tta	tct	ttt	aaa	aca	ttc	att	gaa	gat	gtt	7248
	Val	Lys	Lys	Leu	Asn	Glu	Leu	Ser	Phe	Lys	Thr	Phe	Ile	Glu	Asp	Val	
				2405					2410				2415				
40	aac	aaa	llc	cll	gac	atg	llg	ala	aag	aaa	lla	aag	lca	lll	gal	lac	7296
	Asn	Lys	Phe	Leu	Asp	Met	Leu	Ile	Lys	Lys	Leu	Lys	Ser	Phe	Asp	Tyr	
				2420				2425					2430				
	cac	caq	ttt	gta	gat	gaa	acc	aat	gac	aaa	atc	cgt	gag	gtg	act	caq	7344
	His	Gln	Phe	Val	Asp	Glu	Thr	Asn	Asp	Lys	Phe	Arg	Glu	Val	Thr	Gln	
45			2435				2440					2445					
	aga	ctc	aat	ggt	gaa	att	cag	gct	ctg	gaa	cta	cca	caa	aaa	gct	gaa	7392
	Arg	Leu	Asn	Gly	Glu	Ile	Gln	Ala	Leu	Glu	Leu	Pro	Gln	Lys	Ala	Glu	
			2450				2455					2460					
50	gca	tta	aaa	ctg	ttt	tta	gag	gaa	acc	aag	gcc	aca	gtt	gca	gtg	tat	7440
	Ala	Leu	Lys	Leu	Phe	Leu	Glu	Glu	Thr	Lys	Ala	Thr	Val	Ala	Val	Tyr	
			2465			2470					2475				2480		

55

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

ctg gaa agc cta cag gac acc aaa ata acc tts atc atc aat tgg tta 7488  
 Leu Glu Ser Leu Gln Asp Thr Lys Ile Thr Leu Ile Ile Asn Trp Leu  
 2485 2490 2495

cag gag gct tta agt tca gca tct ttg gct cac atg aag gcc aaa ttc 7536  
 Gln Glu Ala Leu Ser Ser Ala Ser Leu Ala His Met Lys Ala Lys Phe  
 2500 2505 2510

cga gag act cta gaa gat aca cga gac cga atg tat caa atg gac att 7584  
 Arg Glu Thr Leu Glu Asp Thr Arg Asp Arg Met Tyr Gln Met Asp Ile  
 2515 2520 2525

cag cag gaa ctt caa cga tac ctg tct ctg gta ggc cag gtt tat agc 7632  
 Gln Gln Glu Leu Gln Arg Tyr Leu Ser Leu Val Gln Val Tyr Ser  
 2530 2535 2540

aca ctt gtc acc tac att tct gat tgg tgg act ctt gct gct aag aac 7680  
 Thr Leu Val Thr Tyr Ile Ser Asp Trp Trp Thr Leu Ala Ala Lys Asn  
 2545 2550 2555 2560

cct act gac ttt gca gaq caa tat tct atc caa gat tgg gct aaa cct 7728  
 Leu Thr Asp Phe Ala Glu Gln Tyr Ser Ile Gln Asp Trp Ala Lys Arg  
 2565 2570 2575

atg aaa gca ttg gta gag caa ggg ttc act gtt cct gaa atc aag acc 7776  
 Met Lys Ala Leu Val Glu Gln Gly Phe Thr Val Pro Glu Ile Lys Thr  
 2580 2585 2590

atc ctt ggg acc atg cct gcc ttt gaa gtc agt ctt cag gct ctt cag 7824  
 Ile Leu Gly Thr Met Pro Ala Phe Glu Val Ser Leu Gln Ala Leu Gln  
 2595 2600 2605

aaa gcl acc llo cag aca ccl gal lll ala gtc ccc cla aca gal llg 7872  
 Lys Ala Thr Phe Gln Thr Pro Asp Phe Ile Val Pro Leu Thr Asp Leu  
 2610 2615 2620

agg att cca tca gtt cag ata aac ttc aaa gac tta aaa aat ata aaa 7920  
 Arg Ile Pro Ser Val Gln Ile Asn Phe Lys Asp Leu Lys Asn Ile Lys  
 2625 2630 2635 2640

atc cca tcc agg ttt tcc aca cca gaa ttt acc atc ctt aac acc ttc 7968  
 Ile Pro Ser Arg Phe Ser Thr Pro Glu Phe Thr Ile Leu Asn Thr Phe  
 2645 2650 2655

cac att cct tcc ttt aca att gac ttc gta gas atg aaa gta aag atc 8016  
 His Ile Pro Ser Phe Thr Ile Asp Phe Val Glu Met Lys Val Lys Ile  
 2660 2665 2670

aic aga acc all gac cag alg ctg aac agl gag ctg cag tgg ccc gll 8064  
 Ile Arg Thr Ile Asp Gln Met Leu Asn Ser Glu Leu Gln Trp Pro Val  
 2675 2680 2685

cca gat ata tat ctc agg gat ctg aag gtg gag gac att cct cta ggg 8112  
 Pro Asp Ile Tyr Leu Arg Asp Leu Lys Val Glu Asp Ile Pro Leu Ala  
 2690 2695 2700

aga atc acc ctg cca gac ttc cgt tta cca gac atc gca att cca gaa 8160  
 Arg Ile Thr Leu Pro Asp Phe Arg Leu Pro Glu Ile Ala Ile Pro Glu  
 2705 2710 2715 2720

ttc ata atc cca act ctc aac ctt aat gat ttt caa gtt cct gac ctt 8208  
 Phe Ile Ile Pro Thr Leu Asn Leu Asn Asp Phe Gln Val Pro Asp Leu  
 2725 2730 2735

cac ata cca gaa ttc cag ctt ccc cac atc tca cac aca att gaa gta 8256  
 His Ile Pro Glu Phe Gln Leu Pro His Ile Ser His Thr Ile Glu Val  
 2740 2745 2750

cct act ttt ggc aag cta tac agt atc ctg aaa atc caa tct cct ctt 8304  
 Pro Thr Phe Gly Lys Leu Tyr Ser Ile Leu Lys Ile Gln Ser Pro Leu  
 2755 2760 2765

ttc aca tta gat gca aat gct gac ata ggg aat gga acc acc tca gca 8352



EP 2 336 318 A1

Phe Thr Leu Asp Ala Asn Ala Asp Ile Gly Asn Gly Thr Thr Ser Ala  
 2770 2775 2780

5 aac gaa gca ggt atc gca gct tcc atc act gcc aaa gga gag tcc aaa 8400  
 Asn Glu Ala Gly Ile Ala Ala Ser Ile Thr Ala Lys Gly Glu Ser Lys  
 2785 2790 2795 2800

tta gaa gtt ctc aat ttt gat ttt caa gca aat gca caa ctc tca aac 8448  
 Leu Glu Val Leu Asn Phe Asp Phe Gln Ala Asn Ala Gln Leu Ser Asn  
 2805 2810 2815

10 cct aag att aat ccg ctg gct ctg aag gag tca gtg aag ttc tcc agc 8496  
 Pro Lys Ile Asn Pro Leu Ala Leu Lys Glu Ser Val Lys Phe Ser Ser  
 2820 2825 2830

aag tac ctg aga acg gag cat ggg agt gaa atg ctg ttt ttt gga aat 8544  
 Lys Tyr Leu Arg Thr Glu His Gly Ser Glu Met Leu Phe Phe Gly Asn  
 2835 2840 2845

15 gct att gag gga aaa tca aac aca gtg gca agt tta cac aca gaa aaa 8592  
 Ala Ile Glu Gly Lys Ser Asn Thr Val Ala Ser Leu His Thr Glu Lys  
 2850 2855 2860

aat aca ctg gag ctt agt aat gga gtg att gtc aag ata aac aat cag 8640  
 Asn Thr Leu Glu Leu Ser Asn Gly Val Ile Val Lys Ile Asn Asn Gln  
 2865 2870 2875 2880

20 ctt acc ctg gat agc aac act aaa tac ttc cac aaa ttg aac atc ccc 8688  
 Leu Thr Leu Asp Ser Asn Thr Lys Tyr Phe His Lys Leu Asn Ile Pro  
 2885 2890 2895

25 aaa ctg gac ttc tct agt cag gct gac ctg cgc aac gag atc aag aca 8736  
 Lys Leu Asp Phe Ser Ser Gln Ala Asp Leu Arg Asn Glu Ile Lys Thr  
 2900 2905 2910

ctg ttg aaa gct ggc cac ata gca tgg act tct tct gga aaa ggg tca 8784  
 Leu Leu Lys Ala Gly His Ile Ala Trp Thr Ser Ser Gly Lys Gly Ser  
 2915 2920 2925

30 tgg aaa tgg gcc tgg ccc aga ttc tca gat gag gga aca cat gaa tca 8832  
 Trp Lys Trp Ala Ser Pro Arg Phe Ser Asp Glu Gly Thr His Glu Ser  
 2930 2935 2940

caa att agt ttc acc ata gaa gga ccc ctc act tcc ttt gga ctg tcc 8880  
 Gln Ile Ser Phe Thr Ile Glu Gly Pro Leu Thr Ser Phe Gly Leu Ser  
 2945 2950 2955 2960

35 aat aag atc aat agc aaa cac cta aga gta aac caa aac ttg gtt tat 8928  
 Asn Lys Ile Asn Ser Lys His Leu Arg Val Asn Gln Asn Leu Val Tyr  
 2965 2970 2975

gaa tct ggc tcc ctc aac ttt tct aaa ctt gaa att caa tca caa gtc 8976  
 Glu Ser Gly Ser Leu Asn Phe Ser Lys Leu Glu Ile Gln Ser Gln Val  
 2980 2985 2990

gat tcc cag cat gtg ggc cac agt gtc cta act gct aaa ggc atg gca 9024  
 Asp Ser Gln His Val Gly His Ser Val Leu Thr Ala Lys Gly Met Ala  
 2995 3000 3005

45 ctg ttt gga gaa ggg aag gca gag ttc act ggg agg cat gat gct cat 9072  
 Leu Phe Gly Glu Gly Lys Ala Glu Phe Thr Gly Arg His Asp Ala His  
 3010 3015 3020

tta aat gga aag gtt att gga act ttg aaa aat tct ctt ttc ttt tca 9120  
 Leu Asn Gly Lys Val Ile Gly Thr Leu Lys Asn Ser Leu Phe Phe Ser  
 3025 3030 3035 3040

50 gcc cag cca ttt gag atc acg gca tcc aca aac aat gaa ggg aat ttg 9168  
 Ala Gln Pro Phe Glu Ile Thr Ala Ser Thr Asn Asn Glu Gly Asn Leu  
 3045 3050 3055

55

EP 2 336 318 A1

aaa gtt cgt ttt cca tta agg tta aca ggg aag ata gac ttc ctg aat 9216  
 Lys Val Arg Phe Pro Leu Arg Leu Thr Gly Lys Ile Asp Phe Leu Asn  
 3060 3065 3070

5 aac tat gca ctg ttt ctg agt ccc agt gcc cag caa gca agt tgg caa 9264  
 Asn Tyr Ala Leu Phe Leu Ser Pro Ser Ala Gln Gln Ala Ser Trp Gln  
 3075 3080 3085

gta agt gct agg ttc aat cag tat aag tac aac caa aat ttc tct gct 9312  
 Val Ser Ala Arg Phe Asn Gln Tyr Lys Tyr Asn Gln Asn Phe Ser Ala  
 3090 3095 3100

10 gga aac aac gag aac att atg gag gcc cat gta gga ata aat gga gaa 9360  
 Gly Asn Asn Glu Asn Ile Met Glu Ala His Val Gly Ile Asn Gly Glu  
 3105 3110 3115 3120

gca aat ctg gat ttc tta aac att cct tta aca att cct gaa atg cgt 9408  
 Ala Asn Leu Asp Phe Leu Asn Ile Pro Leu Thr Ile Pro Glu Met Arg  
 3125 3130 3135

15 cta cct tac aca ata atc aca act cct cca ctg aaa gat ttc tct cta 9456  
 Leu Pro Tyr Thr Ile Ile Thr Thr Pro Pro Leu Lys Asp Phe Ser Leu  
 3140 3145 3150

tgg gaa aca aca gcc ctg aag gaa ttc ttg aac acg aca aag caa tca 9504  
 Trp Glu Lys Thr Gly Leu Lys Glu Phe Leu Lys Thr Thr Lys Gln Ser  
 3155 3160 3165

20 ttt gat tta agt gta aaa gct cag tat aag aaa aac aaa cac agg cat 9552  
 Phe Asp Leu Ser Val Lys Ala Gln Tyr Lys Lys Asn Lys His Arg His  
 3170 3175 3180

25 ccg atc aca aat cct ctg gct gtg ctt tgl gag ttt atc agt cag acc 9600  
 Ser Ile Thr Asn Pro Leu Ala Val Leu Cys Glu Phe Ile Ser Gln Ser  
 3185 3190 3195 3200

atc aaa tcc ttt gac agg cat ttt gaa aaa aac aca aac aat cca tta 9648  
 Ile Lys Ser Phe Asp Arg His Phe Glu Lys Asn Arg Asn Asn Ala Leu  
 3205 3210 3215

30 gat ttt gtc acc aaa ccc tat aat gaa aca aaa att aag ttt gat aag 9696  
 Asp Phe Val Thr Lys Ser Tyr Asn Glu Thr Lys Ile Lys Phe Asp Lys  
 3220 3225 3230

tac aaa gct gaa aaa tct cac gac gag ctg ccc agg acc ttt caa atc 9744  
 Tyr Lys Ala Glu Lys Ser His Asp Glu Leu Pro Arg Thr Phe Gln Ile  
 3235 3240 3245

35 ccl gga tcc act gtt cca gtt gtc aal gtt gaa gtg tcl cca ttc acc 9792  
 Pro Gly Tyr Thr Val Pro Val Val Asn Val Glu Val Ser Pro Phe Thr  
 3250 3255 3260

ata gag atg tgg gca ttc ggc tat gtg ttc cca aaa gca gtc agc atg 9840  
 Ile Glu Met Ser Ala Phe Gly Tyr Val Phe Pro Lys Ala Val Ser Met  
 3265 3270 3275 3280

40 cct agt ttc tcc atc ata ggt tct gac gtc cgt gtg cct tca tac aca 9888  
 Pro Ser Phe Ser Ile Ile Gly Ser Asp Val Arg Val Pro Ser Tyr Thr  
 3285 3290 3295

45 tta atc ctg cca tca tta gag ctg cca gtc ctt cat gtc cct aga aat 9936  
 Leu Ile Leu Pro Ser Leu Clu Leu Pro Val Leu His Val Pro Arg Asn  
 3300 3305 3310

ctc aag ctt tct ctt cca gat ttc aag gaa ttg tgt acc ata agc cat 9984  
 Leu Lys Leu Ser Leu Pro Asp Phe Lys Glu Leu Cys Thr Ile Ser His  
 3315 3320 3325

50 atc ttt att cct gcc atg ggc aat atc acc tat gat ttc tcc ttt aaa 10032  
 Ile Phe Ile Pro Ala Met Gly Asn Ile Thr Tyr Asp Phe Ser Phe Lys  
 3330 3335 3340

55

EP 2 336 318 A1

tca agt gtc atc aca ctg aat acc aat gct gac ctt ttt aac cag tca 10080  
 Ser Ser Val Ile Thr Leu Asn Thr Asn Ala Glu Leu Phe Asn Gln Ser  
 3345 3350 3355 3360  
 5  
 gat att gtt gct cat ctc ctt tct tca tct tca tct gtc att gat gca 10128  
 Asp Ile Val Ala His Leu Leu Ser Ser Ser Ser Ser Val Ile Asp Ala  
 3365 3370 3375  
 ctg cag tac aaa tta gag ggc acc aca aga ttg aca aga aaa agg gga 10176  
 Leu Gln Tyr Lys Leu Glu Gly Thr Thr Arg Leu Thr Arg Lys Arg Gly  
 3380 3385 3390  
 10  
 ttg aag tta gcc aca gct ctg tct ctg agc aac aaa ttt gtg cag gct 10224  
 Leu Lys Leu Ala Thr Ala Leu Ser Leu Ser Asn Lys Phe Val Glu Gly  
 3395 3400 3405  
 agt cat acc agt act gtg agc tta acc acg aac aat atg gaa gtg tca 10272  
 Ser His Asn Ser Thr Val Ser Leu Thr Thr Lys Asn Met Glu Val Ser  
 3410 3415 3420  
 15  
 gtg gca aaa acc aca aaa ccg gaa att cca att ttg aga atg aat ttc 10320  
 Val Ala Lys Thr Thr Lys Pro Glu Ile Pro Ile Leu Arg Met Asn Phe  
 3425 3430 3435 3440  
 aag caa gaa ctt aal gga aal acc aag tca aaa cct act gtc tct tcc 10368  
 Lys Gln Glu Leu Asn Gly Asn Thr Lys Ser Lys Pro Thr Val Ser Ser  
 3445 3450 3455  
 20  
 tcc atg gaa ttt aag tat gat ttc aat tct tca atg ctg tac tct acc 10416  
 Ser Met Glu Phe Lys Tyr Asp Phe Asn Ser Ser Met Leu Tyr Ser Thr  
 3460 3465 3470  
 25  
 gct aaa gga gca gtt gac cac cag ctt agc ttg gaa agc ctc acc tct 10464  
 Ala Lys Gly Ala Val Asp His Lys Leu Ser Leu Glu Ser Leu Thr Ser  
 3475 3480 3485  
 tac ttt tcc att gag tca tct acc aaa gga gat gtc aag ggt tct gtt 10512  
 Tyr Phe Ser Ile Glu Ser Ser Thr Lys Gly Asp Val Lys Gly Ser Val  
 3490 3495 3500  
 30  
 ctt tct cgg gaa tat tca gga act att gct agt gag gcc aac act tac 10560  
 Leu Ser Arg Glu Tyr Ser Ser Gly Thr Ile Ala Ser Glu Ala Asn Thr Tyr  
 3505 3510 3515 3520  
 ttg aat tcc aag agc aca cgg tct tca gtg aag ctg cag gcc act tcc 10608  
 Leu Asn Ser Lys Ser Thr Arg Ser Ser Val Lys Leu Gln Gly Thr Ser  
 3525 3530 3535  
 35  
 aaa att gat gat atc tgg aac ctt gaa gta aaa gaa aat ttt gct gga 10656  
 Lys Ile Asp Asp Ile Trp Asn Leu Glu Val Lys Glu Asn Phe Ala Gly  
 3540 3545 3550  
 gaa gcc aca ctc caa cgc ata tat tcc ctc tgg gag cac agt acg aaa 10704  
 Glu Ala Thr Leu Gln Arg Ile Tyr Ser Leu Trp Glu His Ser Thr Lys  
 3555 3560 3565  
 aac cac tta cag cta gag ggc ctc ttt ttc acc aac gga gaa cat aca 10752  
 Asn His Leu Gln Leu Glu Gly Leu Phe Phe Thr Asn Gly Glu His Thr  
 3570 3575 3580  
 40  
 agc aaa gcc acc ctg gaa ctc tct cca tgg cac atg tca gct ctt gtt 10800  
 Ser Lys Ala Thr Leu Glu Leu Ser Pro Trp Gln Met Ser Ala Leu Val  
 3585 3590 3595 3600  
 cag gtc cat gca agt cag ccc agt tcc ttc cat gat ttc cct gac ctt 10848  
 Gln Val His Ala Ser Gln Pro Ser Ser Phe His Asp Phe Pro Asp Leu  
 3605 3610 3615  
 50  
 ggc cag gaa gtg gcc ctg aat gct aac act aag aac cag aag atc aga 10896  
 Gly Gln Glu Val Ala Leu Asn Ala Asn Thr Lys Asn Gln Lys Ile Arg  
 3620 3625 3630  
 55

EP 2 336 318 A1

tgg aaa aat gaa gtc cgg att cat tcc ggg tct ttc cag agc cag gtc 10944  
 Trp Lys Asn Glu Val Arg Ile His Ser Gly Ser Phe Gln Ser Gln Val  
 3635 3640 3645

5 gag ctt tcc aat gac caa gaa aag gca cac ctt gac att gca gga tcc 10992  
 Glu Leu Ser Asn Asp Gln Glu Lys Ala His Leu Asp Ile Ala Gly Ser  
 3650 3655 3660

tta gaa gga cac cta agg ttc ctc aaa aat atc atc cta cca gtc tat 11040  
 Leu Glu Gly His Leu Arg Phe Leu Lys Asn Ile Ile Leu Pro Val Tyr  
 3665 3670 3675 3680

10 gac aag agc tta tgg gat ttc cta aag ctg gat gtc acc acc agc att 11088  
 Asp Lys Ser Leu Trp Asp Phe Leu Lys Leu Asp Val Thr Thr Ser Ile  
 3685 3690 3695

ggt agg aga cag cat ctt cgt gtt tca act gcc ttt gtg tac acc aaa 11136  
 Gly Arg Arg Gln His Leu Arg Val Ser Thr Ala Phe Val Tyr Thr Lys  
 3700 3705 3710

15 aac ccc aat gcc tat tca ttc tcc atc cct gta aaa gtt ttg cct gat 11184  
 Asn Pro Asn Gly Tyr Ser Phe Ser Ile Pro Val Lys Val Leu Ala Asp  
 3715 3720 3725

aaa ttc att att cct ggg ctg aaa cta aat gat cta aat tca gtt ctt 11232  
 Lys Phe Ile Ile Pro Gly Leu Lys Leu Asn Asp Leu Asn Ser Val Leu  
 3730 3735 3740

20 gtc atg cct acg ttc cat gtc cca ttc aca gat ctt cag gtt cca tgg 11280  
 Val Met Pro Thr Phe His Val Pro Phe Thr Asp Leu Gln Val Pro Ser  
 3745 3750 3755 3760

25 tgc aaa ctt gac ttc aga gaa ata caa atc tat aag aag ctg aga act 11328  
 Cys Lys Leu Asp Phe Arg Glu Ile Gln Ile Tyr Lys Lys Leu Arg Thr  
 3765 3770 3775

tca tca ttt gcc ctc acc cta cca aca ctc ccc gag gta aaa ttc cct 11376  
 Ser Ser Phe Ala Leu Thr Leu Pro Thr Leu Pro Glu Val Lys Phe Pro  
 3780 3785 3790

30 gaa gtt gat gtg tta aca aaa tat tcc caa cca gaa gac tcc ttg att 11424  
 Glu Val Asp Val Leu Thr Lys Tyr Ser Gln Pro Glu Asp Ser Leu Ile  
 3795 3800 3805

35 ccc ttt ttt gag ata acc gtg cct gaa tct cag tta act gtg tcc cag 11472  
 Pro Phe Phe Glu Ile Thr Val Pro Glu Ser Gln Leu Thr Val Ser Gln  
 3810 3815 3820

ttc acg ctt cca aaa agt gtt tca gat gcc att gct gct ttg gat cta 11520  
 Phe Thr Leu Pro Lys Ser Val Ser Asp Gly Ile Ala Ala Leu Asp Leu  
 3825 3830 3835 3840

40 aat gca gta gcc aac aag atc gca gac ttt gag ttg ccc acc atc atc 11568  
 Asn Ala Val Ala Asn Lys Ile Ala Asp Phe Glu Leu Pro Thr Ile Ile  
 3845 3850 3855

gag cct gag cag acc att gag att ccc tcc att aag ttc tct gta cct 11616  
 Val Pro Glu Gln Thr Ile Glu Ile Pro Ser Ile Lys Phe Ser Val Pro  
 3860 3865 3870

45 gct gga att gtc att cct tcc ttt caa gca ctg act gca cgc ttt gag 11664  
 Ala Gly Ile Val Ile Pro Ser Phe Gln Ala Leu Thr Ala Arg Phe Glu  
 3875 3880 3885

cta gac tct ccc gta tat aat gcc act tgg agt gcc agt ttg aaa aac 11712  
 Val Asp Ser Pro Val Tyr Asn Ala Thr Trp Ser Ala Ser Leu Lys Asn  
 3890 3895 3900

50 aaa gca gat tat gtt gaa aca gtc ctg gat tcc aca tgc agc tca acc 11760  
 Lys Ala Asp Tyr Val Glu Thr Val Leu Asp Ser Thr Cys Ser Ser Thr  
 3905 3910 3915 3920

55

EP 2 336 318 A1

5  
gta cag ttc cta gaa tat gaa cta aat gtt ttg gga aca cac saa atc 11808  
Val Gln Phe Leu Glu Tyr Glu Leu Asn Val Leu Gly Thr His Lys Ile  
3925 3930 3935

gaa gat ggt acc tta gcc tct aag act aaa gga aca ctt gca cac cgt 11856  
Glu Asp Gly Thr Leu Ala Ser Lys Thr Lys Gly Thr Leu Ala His Arg  
3940 3945 3950

gac ttc agt gca gaa tat gaa gaa gat ggc aaa tat gaa gga ctt cag 11904  
Asp Phe Ser Ala Glu Tyr Glu Glu Asp Gly Lys Tyr Glu Gly Leu Gln  
3955 3960 3965

gaa tgg gaa gga aaa gcg cac ctc aat atc aaa aqc cca gcg ttc acc 11952  
Glu Trp Glu Gly Lys Ala His Leu Asn Ile Lys Ser Pro Ala Phe Trp  
3970 3975 3980

gat ctc cat ctg cgc tac cag aaa gac aag aaa ggc atc tcc acc tca 12000  
Asp Leu His Leu Arg Tyr Gln Lys Asp Lys Lys Gly Ile Ser Thr Ser  
3985 3990 3995 4000

gca gcc tcc cca gcc gta ggc acc gtg ggc atg gat atg gat gaa gat 12048  
Ala Ala Ser Pro Ala Val Gly Thr Val Gly Met Asp Met Asp Glu Asp  
4005 4010 4015

gac gac lll lcl aaa lgg aac ltc lac lac agc ccl cag lcc lcl cca 12096  
Asp Asp Phe Ser Lys Trp Asn Phe Tyr Tyr Ser Pro Gln Ser Ser Pro  
4020 4025 4030

gat aaa asa ctc acc ata ttc aaa act gag ttg agg gtc cgg gaa tct 12144  
Asp Lys Lys Leu Thr Ile Phe Lys Thr Glu Leu Arg Val Arg Glu Ser  
4035 4040 4045

gat gag gaa act cag atc aaa gtt aat tgg gaa gaa gag gca gct tct 12192  
Asp Glu Glu Thr Gln Ile Lys Val Asn Trp Glu Glu Glu Ala Ala Ser  
4050 4055 4060

ggc ttg cta acc tct ctg aaa gac aac gtg ccc aag gcc aca ggg gtc 12240  
Gly Leu Leu Thr Ser Leu Lys Asp Asn Val Pro Lys Ala Thr Gly Val  
4065 4070 4075 4080

30  
ctt tat gat tat gtc aac aag tac cac tgg gaa cac aca ggg ctc acc 12288  
Leu Tyr Asp Tyr Val Asn Lys Tyr His Trp Glu His Thr Gly Leu Trp  
4085 4090 4095

ctg aga gaa gtg tct tca aag ctg aga aga aat ctg cag aac aat gct 12336  
Leu Arg Glu Val Ser Ser Lys Leu Arg Arg Asn Leu Gln Asn Asn Ala  
4100 4105 4110

35  
gag tgg gtt tat caa ggg gcc att agg caa att gat gat atc gac gtg 12384  
Glu Trp Val Tyr Gln Gly Ala Ile Arg Gln Ile Asp Asp Ile Asp Val  
4115 4120 4125

agg ttc cag aaa gca gcc agt ggc acc act ggg acc tac caa gag tgg 12432  
Arg Phe Gln Lys Ala Ala Ser Gly Thr Thr Gly Thr Tyr Gln Glu Trp  
4130 4135 4140

40  
aag gac aag gcc cag aat ctg tac cag gaa ctg ttg act cag gaa gcc 12480  
Lys Asp Lys Ala Gln Asn Leu Tyr Gln Glu Leu Leu Thr Gln Glu Gly  
4145 4150 4155 4160

45  
caa gcc agt ttc cag gga ctc aag gat aac gtg ttt gat gcc tgg gta 12528  
Gln Ala Ser Phe Gln Gly Leu Lys Asp Asn Val Phe Asp Gly Leu Val  
4165 4170 4175

oga gtt act caa aaa ttc cat atg aaa gtc aag aag ctg att gac tca 12576  
Arg Val Thr Gln Lys Phe His Met Lys Val Lys Lys Leu Ile Asp Ser  
4180 4185 4190

50  
cgc att gat lll ctg aac ltc ccc aga ltc cag lll ccg ggg aaa cc 12624  
Leu Ile Asp Phe Leu Asn Phe Pro Arg Phe Gln Phe Pro Gly Lys Pro

55

EP 2 336 318 A1

	4195	4200	4205	
5	egg ala lac acI agg gag gaa cll Lgc acI atg lIc atg agg gag gla Gly Ile Tyr Thr Arg Glu Glu Leu Cys Thr Met Phe Met Arg Glu Val 4210 4215 4220	12672		
	egg acg gta ctg tcc caq gta tat tcc aaa gtc cat aat ggt tca gaa Gly Thr Val Leu Ser Gln Val Tyr Ser Lys Val His Asn Gly Ser Glu 4225 4230 4235 4240	12720		
10	ata ctg ttt tcc tat ttc caa gac cta gtg att aca ctt cct ttc gag Ile Leu Phe Ser Tyr Phe Gln Asp Leu Val Ile Thr Leu Pro Phe Glu 4245 4250 4255	12768		
	tta agg asa cat aaa cta ata gat gta atc tcc atg tat agg gaa ctg Leu Arg Lys His Lys Leu Ile Asp Val Ile Ser Met Tyr Arg Glu Leu 4260 4265 4270	12816		
15	Llg aaa gal lla lca aaa gaa gcc caa gag gla lll aaa gcc aLL cag Leu Lys Asp Leu Ser Lys Glu Ala Gln Glu Val Phe Lys Ala Ile Gln 4275 4280 4285	12864		
	tct ctg aag acc aca gag gtg cta cgt aat ctt cag gac ctt tta caa Ser Leu Lys Thr Thr Glu Val Leu Arg Asn Leu Gln Asp Leu Leu Gln 4290 4295 4300	12912		
20	ttc att ttc caa cta ata gaa gat aac att aac cag ctg aaa gag atg Phe Ile Phe Gln Leu Ile Glu Asp Asn Ile Lys Gln Leu Lys Glu Met 4305 4310 4315 4320	12960		
	aaa ttt act tat ctt att aat tat atc caa gat gag atc aac aca atc Lys Phe Thr Tyr Leu Ile Asn Tyr Ile Gln Asp Glu Ile Asn Thr Ile 4325 4330 4335	13008		
25	ttc aat gat tat atc cca tat gtt ttc aaa ttg ttg aaa gaa aac cta Phe Asn Asp Tyr Ile Pro Tyr Val Phe Lys Leu Leu Lys Glu Asn Leu 4340 4345 4350	13056		
	tgc ctt aat ctt cat aag ttc aat gaa ttt att caa aac gag ctt cag Cys Leu Asn Leu His Lys Phe Asn Glu Phe Ile Gln Asn Glu Leu Gln 4355 4360 4365	13104		
30	gaa gct tct caa gag lta cag cag atc cat caa tac att atg gcc ctt Glu Ala Ser Gln Glu Leu Gln Gln Ile His Gln Tyr Ile Met Ala Leu 4370 4375 4380	13152		
	cgt gaa gaa tat ttt gat cca agt ata gtt ggc tgg aca gtg aaa tat Arg Glu Glu Tyr Phe Asp Pro Ser Ile Val Gly Trp Thr Val Lys Tyr 4385 4390 4395 4400	13200		
35	tat gaa ctt gaa gaa aag ata gtc agt ctg atc aag aac ctg tta gtt Tyr Glu Leu Glu Glu Lys Ile Val Ser Leu Ile Lys Asn Leu Leu Val 4405 4410 4415	13248		
	gct ctt aag gac ttc cat tct gaa tat att gtc agt gcc tct aac ttc Ala Leu Lys Asp Phe His Ser Glu Tyr Ile Val Ser Ala Ser Asn Phe 4420 4425 4430	13296		
	act tcc caa ctg tca agt caa gtt gag caa ttt ctg cac aga aat att Thr Ser Gln Leu Ser Ser Gln Val Glu Gln Phe Leu His Arg Asn Ile 4435 4440 4445	13344		
45	cag gaa tat ctt agc atc ctt acc gat cca gat gga aaa ggg aaa gag Gln Glu Tyr Leu Ser Ile Leu Thr Asp Pro Asp Gly Lys Gly Lys Glu 4450 4455 4460	13392		
	aaq att gca gac ctt tct gcc act gct cag gaa ata att aaa aqc cag Lys Ile Ala Glu Leu Ser Ala Thr Ala Gln Glu Ile Ile Lys Ser Gln 4465 4470 4475 4480	13440		
50	gcc att gcg acc aag aaa ata att tct gat tac cac cag cag ttt aga Ala Ile Ala Thr Lys Lys Ile Ile Ser Asp Tyr His Gln Gln Phe Arg 4485 4490 4495	13488		
55				

EP 2 336 318 A1

tat aaa ctg caa gat ttt tca gac caa ctc tct gat tac tat gaa aaa 13536  
 Tyr Lys Leu Glu Asp Phe Ser Asp Gln Leu Ser Asp Tyr Tyr Glu Lys  
 4500 4505 4510  
 5  
 ttt att gct gaa tcc aaa aga ttg att gac ctg tcc att caa aac tac 13584  
 Phe Ile Ala Glu Ser Lys Arg Leu Ile Asp Leu Ser Ile Gln Asn Tyr  
 4515 4520 4525  
 cac aca ttt ctg ata tac atc acg gag tta ctg aaa aag ctg caa tca 13632  
 His Thr Phe Leu Ile Tyr Ile Thr Glu Leu Leu Lys Lys Leu Gln Ser  
 4530 4535 4540  
 10  
 acc aca gtc atc aac ccc tac atg aag ctt gct cca gga gaa ctt acg 13680  
 Thr Thr Val Met Asn Pro Tyr Met Lys Leu Ala Pro Gly Glu Leu Thr  
 4545 4550 4555 4560  
 atc atc ctc taa tttttttaa agaaatctc atttattctt cttttccaat 13732  
 Ile Ile Leu \*  
 15  
 tgaactttca cataqcacag aaaaaattca aactgcctat attgataaaa ccatacagtg 13792  
 agccagcctt gcagtaggca gtagactata agcagaagca cataagaaat ggacctgcac 13852  
 caaagctggc accagggctc ggaaggtctc tgaactcaga aggatggcat tttttgcaag 13912  
 20  
 ttaaagaaa tcagatctg agttattttg ctaaaacttg gggaggagga acaataaat 13972  
 ggagtcttta ttgtgtatca t 13993  
 <210> 319  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 319  
 gcctcagctt gcttcgccc 20  
 30  
 <210> 320  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 320  
 gctcactggt cagcatctgg 20  
 40  
 <210> 321  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 45  
 <400> 321  
 tgagaatctg ggcgaggccc 20  
 <210> 322  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 55

EP 2 336 318 A1

5  
 <400> 322  
 g-ccttcata ttgccatcc 20  
 <210> 323  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

10  
 <400> 323  
 cctccctcat gaacatagtg 20  
 <210> 324  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

15  
 <400> 324  
 gacgtcagaa cctatgatgg 20  
 <210> 325  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

20  
 <400> 325  
 tgagtgagtc aatcagcttc 20  
 <210> 326  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

25  
 <400> 326  
 gccttctgct tgaattacaa 20  
 <210> 327  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

30  
 <400> 327  
 gcgccttctg cttgagttac 20  
 <210> 328  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

35  
 <400> 328  
 tcgcgccttc tgcttgagtc 20  
 <210> 329  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

40  
 <400> 329  
 <210> 329  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

45  
 <400> 329  
 <210> 329  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

50  
 <400> 329  
 <210> 329  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

55



EP 2 336 318 A1

<220>  
 <223> Antisense Oligonucleotide  
 5 <400> 329  
 cttcgcgcct tctgcttgag 20  
 <210> 330  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 330  
 agtctgcttc gcgccttctg 20  
 <210> 331  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 331  
 tcagtctgct tcgcgccttc 20  
 <210> 332  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 332  
 cctcagtctg cttcgcgcct 20  
 <210> 333  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 333  
 agcctcagtc tgcctcgcgc 20  
 <210> 334  
 <211> 43445  
 <212> DNA  
 <213> H. sapiens  
 30 <400> 334  
 accaagacag cgtcaggac tggttctcct cgtggctccc aattcagtc aggagaagca 60  
 gaqattttgt ccccatggtg ggtcatttga aaaaagcacc cctggtcagg gcagccttct 120  
 35 <400> 334  
 cagaccctga ggcgctgggc atggcccac tgagacacag gaagggcgc gccagagcac 180  
 tgaagacgct tggggaagg aaccacactg ggaccacgcc cctgggtgct gggctecat 240  
 40 <400> 334  
 cccaggtggg cccctccc gaggtcttc aaggctcaaa gagaagccag tctagaaaag 300  
 caaacaggtc aggccggga ggcgccttt gaccctttg caatcctggc gctcttcag 360  
 45 <400> 334  
 cctgggcttc ctataaatg ggtgcccgg ccggccgcgc attcccacg gacctcgg 420  
 50 <400> 334  
 gctcagctgc ccttcgcg tctcgcgc gaggagccg cccagccag caggccgc 480  
 55

EP 2 336 318 A1

aggccgaggc caggccgcag cccaggagcc gccccaccgc agctggcgat ggaccccgcc 540  
aggcccgccg cglggcgcl gclggcgclg cclggcgclg lglgclglc gclggcgccg 600  
5 gccaggggccg gtgagtgccg ggcgcctctg cgggcgcaga gggagcggga gggagccggc 660  
ccacagaggt tggccggcc agcctggcc tagcccaag ggaagccagc cacagcccc 720  
aggcgagtg gggggattg accagctggc ggcctgca ggcctaggat gggggggcg 780  
ggatggagg gctgaggag ggtctccgg agcctgcct cctctgaaa gg-gaaaact 840  
10 g-gccggtg tccccctgc gggccctagc acccctggg aagacgtggg aagctcacag 900  
at-tctttct cctg-cttac agaagaggaa at-gctggaa atgtcagcct gg-ctgtcca 960  
agtaaggcat ctgcgcctg ggcgtggaag ggcgccagc cccgtgcact ctctacacc 1020  
15 cgggccccg agggccccc acclacagc cclgagalg calcglgglg lcccllclc 1080  
tgaccccaag aagcaagtcc cctgagcctc tccccacacc caagggatgc caactctct 1140  
ctacotggcc tctgttctg tccaaaagt tcagcctggg ggcgggggag ggaagggatt 1200  
g-ctctccgc tggcctgtc acactttgaa gaaacatcac tgtctgttt atcagtgcact 1260  
20 agtcattgat tcgaagcatg tgagggtgag gaaatactga cttaacott tg-gaaagaa 1320  
tcgaacctcc acccccttcc tatttaactg acccctggg gttaaaggaa ctggcctcca 1380  
agccgcacc tgtg-gctg agcccgggg cggacttctg atggggcagc accgccact 1440  
25 aglggccglc lglcalcac gcagclggac tcaggaccca galgllclll llcllcaall 1500  
gtcagaaaa ttcctctcaa ctacagtga aacctccaga aattcttttc taggagttg 1560  
t-aaqtaagt tacccttaa- gcttaatgaa ctttgcctta agta-ttgct agtcttaag 1620  
tcacggaatt accgctgtc caagctaaaa aagcattaga gatagtacta tt-gcgtaat 1680  
30 g-tgtcatct cttaatttgc cagagggtct c-catgcaga tttctgagc ccattactt 1740  
gacacttgc actcccttc ctgtgcctca gatgagatat tcaagacatg ccagccaatt 1800  
taaacattag cctcagcaaa aacatactg agaagtcaa tctataaagg asaattaagt 1860  
a-aaagtcaa ttaaaaaata atttgagttg aattaccatt ttaattctc ta-gccactg 1920  
35 cccclclclg ccagaaallg gclglccllg cgagagclal llclglclag lggclgacgl 1980  
at-tctcccc agt-tagaag atgcgaccg atcaagcac ctccggaagt acacatacaa 2040  
ctatgaggct gagagttcca gtggagtccc tgggactgct gattcaagaa gtgccaccag 2100  
40 gatcaactgc aagg-tatgga gcatgcaggc aggagggacc tagagcccac agctttcccc 2160  
cagccctgtt ccagcggggc cccaacacgc gaccttccc gaggggtgtc actgagcaaa 2220  
ccgagaacat ccagaaactg ttgtaactg atcaaaagcac tgggactttg cc-ctgtttg 2280  
taagtccgc acat-gctga gatgtgtct gccccacca aatttcgcaa gtcagaagta 2340  
45 t-ttccggtt aact-cccag atgcaatagg aatccatgat ctgatttagc agcagtg-gg 2400  
g-ctgtagat ttcagcgtga gagaggcca ctaggtgagc tatgggaggc aggcaac-cg 2460  
caatcgcact gtgaaatgca gttttataa t-taagtcaa acagaatctg ttgctgaaaa 2520  
50 a-gaatggaa agaagaaaa aatataaaca tacagtttgt tctaaaataa aactttgctt 2580  
at-tattgaga ctgctgttac tcatgttaca tacatgtgga gcagatctac aggctgotat 2640  
tggggtttg gtggggaaga gaagtcagc tgagcagtca cctt-tttta gagagtaacc 2700

55

EP 2 336 318 A1

tagctcttgt atgtgctgtc caatatggta gacatgagcc acattgggct atttaaaagg 2760  
 5 aatgaaatta aaaactcata ttogttgtca cattagctgc atttcaactg ctcaacagcc 2820  
 accctggcta ctggctccca tattgaacag cacacatgta caacatttct ataaagttat 2880  
 ttgaatagtg ctggataata agtaggaatc cgttgaact ccagctatat gcaagctct 2940  
 aaataggccc taatagatac aaccagtttt ttgggtgaca ttaaggagac atttgcctgtg 3000  
 10 gaaaogaagg atggcctcc toctgcttcc tgtttttctt ctccacttcc actcctagtc 3060  
 tqcaqccctt ctatttaacc acagctcttt acaattaaag tqagtaactt taqaaccaat 3120  
 aaaaggacat cctccttccc atgcttaggg gcaaacctaa gaaatgtgtt accogggagg 3180  
 gggaaaaagt cagcaatagg actaagtcta ggttgggtgc cagagaaccc aggaggcatg 3240  
 15 ttgataaggc atgtggtgtt gaggcgcagg cagtgggttt cccagcacca ttccctttgg 3300  
 tqctctgatt agacattaaq ccttggcctt caggggccac ctctcattct tqatagaca 3360  
 cctcaatgct ctgcacccc gaattctcag gttgagctgg aggttcccca gctctgagc 3420  
 ttcatctctg agaccagcca gtgcacctg aaagaggtgt atggcttcaa cctgagggc 3480  
 20 aaagccttgc tgaagaaaac caagaactct gaggagtttg ctgcagccat gtcacgcta 3540  
 gcatgttgtt acatgagcac acgcatgtgt gttgtccgc tgaggatga acctgttgt 3600  
 ttgcaccagg caccgatgtg actgtaagta ttgtattcc gtatccatcg tggatcaggg 3660  
 25 aallactgag llllcacaa calcaaaaag agagaagcal lagllaaccl lccctagcla 3720  
 ggttccctta attatcattt tcatgtgttt ctaaaaatct catgctttaa accctttgag 3780  
 attataaaac tqacatgctt tqttaaaaca agtqaattct tatttaaaqa actagtaag 3840  
 actagtctt ggtgcttctt ggtgtgggt cccagaggca ctgctgctg tggccggcac 3900  
 30 atggcggggc agggctgtt cacgcaggg cagaggagca ccaaggcttc ggtggctccc 3960  
 cctcctaggc tggcattcag ccaactgacg ctgatggcc actgcagctg catctctgct 4020  
 gactggtcag ggcacatgtc gcacccattg taaatatttt caacatccc cctgctcat 4080  
 cctcaatcac agttttagg gtctaggtg tgtatgaata caggcaggat agagttgta 4140  
 35 acllgglagc alcagaaaac tclgtclgla ltaglclgll llcalgtclg clalaaagac 4200  
 atacctgaga ctgggcaat tcaaaaagaa aggtttattg gactcacagt tccactggc 4260  
 tggggaggtc tcacaatcac ggcggaaggt gagggacagc aagtcacatc ttatgtagat 4320  
 40 ggtggctggc aaagagagc tgtgcagaga aactcctgtt tttagaacca tcagatccc 4380  
 cgacacccat ctgcaatcac gagaacagca cgggaaagac ctgccccat gatcaatca 4440  
 cctccccccg ggtccctccc acaacaagtg ggaattatga gagctacgag acgaaatttg 4500  
 ggtggggagc cagagccaaa ccatatcacc acccttccc atttttcagt ttgctaacc 4560  
 45 attagattca gatgccagtc cttcttggc aaaataggct gtgaggcttc ttctttct 4620  
 atgctttatt ttctccaaga cttaactgta tatgaggag aggggtatgg tggcagagg 4680  
 aaagagtgtt ttatttttg gtcttggtc ttctccaaat acacaagaca ctctgttct 4740  
 50 tgaagaggag ggtctcca gtttgcctct tcatgacttt aactgtcttt tttaaaaatt 4800  
 gacatacaat aattataca atttattgag aacatagtga tattttgata caagtaagt 4860  
 atgggtgatc gatcagagta attagcatac ccactatctc aaacatttat cttttctctg 4920

55

EP 2 336 318 A1

5 tgttgggaac tttctgagag agtgtaggct ctgggagata agtccgtoac cttttccccc 4980  
 lgalglaacc agag\_ggclg cagccagglc c\_cagaaact cagagaglac ccaglgggaa 5040  
 a\_cccctaaga ccaaaagtcag catgggcttc agccatggcc tgacaccata caaaagaatg 5100  
 actgtccaac aagtqtatca aaataaagctc caattcactg gtatgcaaca aatgcgaatt 5160  
 aatgtaacaa caagatattc atctgctttt acccatcata ctgcaaaact ggaaaacagt 5220  
 gatagcaccct gttctctggca ggccagtggag gaaaagtgtg ctgtcctggag ctgctggcgg 5280  
 10 aaacgagagc catcaggcaa tatctactgt aatttaaaat acttaatacc ctctgacaca c340  
 gatatttttag tcttgggac tctagcccat gaaaataaaa gcagtaatgc gtgaagatag 5400  
 gcacataagg atgtctgttc tggattggtt tgtgtggttt aaaaaaatc cagaaagaga 5460  
 15 cagggcaaal gccacaaa\_ ggggcaelgl c\_gaalaaal lalalllagc ca\_ggaa\_gg 5520  
 aatgtttctgc atgcagcttc taaaaaatc tgttagagct gtaccaagtq ac\_cagaagq 5580  
 a\_ttttgtga agta\_aatta agtgagaaaa acaagataaa agtatgcata atacaatgcc 5640  
 acttgataa aacaaacaa\_ ggcaaaatct t\_tgatgact ctgtttgcac tcacccaatg 5700  
 20 ttacagagga ttgatgagc gtgcagcaac aaatggaaca accactcggg tgtccgtatg 5760  
 gggagatgg gcaaaagagc tgatatgggt cgagaacaga gcagggctgg atgagccaag 5820  
 caaaaaaagt taanaacacag ctggacctgg tggctcatgc ctgtagtccc agcaacttgg 5880  
 25 cagggccgag agggagaaic acclgagglc aggaglllgc gaccagcccg gccaca\_gg 5940  
 tgaaaaactgt ctctactaaa aatacaaaaa tttagctgggt gtgatggcac atgccagtag 6000  
 tcttagctac tcccagggc\_ gaggcaggag aatcaactga tcccaggaq tggaggttc 6060  
 agtgagctga ggttgcgca ttgcactcca gcccgggcga ccgagcgaga ctccatttca 6120  
 30 aaaaaagaa aagaaaaaag aaaaaagaa aaaaaagaa tcacaaaaac ttatgtatat 6180  
 g\_gcataact ttttgaaaa\_ gtatgtctat c\_gtagctat attctatatt tacaataaa 6240  
 tgatgtcaga agaacaattg gttaaaaaaa tatgagaaa\_ gaaacttcag tcccaccag 6300  
 ctacttcca gcaagttgta atggagaagg acatttccgt gaccatctc tcctggggac 6360  
 35 agglalgagc lcaagclggc callccagaa ggaagcagg llllccllla ccggagaaa 6420  
 gatgaacctc ctacatccc\_ gaacatcaag aggggcatca tttctgccc ctgggttccc 6480  
 ccagagacag aagaagccaa gcaagtgtt t\_tctggtga ggatttagaa agctgatagc 6540  
 40 agtggccctt gaaactcacc ttcattgttt agagaccagt cctaccata\_ acaaagcaga 6600  
 t\_cactgagtc agctccatga ctagtacat aggaagccct ggattgggt gaaatacgg 6660  
 tgcccaggt tctctctgc ccttaggctc actgacagat catcccagc aggottaica 6720  
 ggttgggtct aattttaaaa cagtcattga cagtcctgq ccaccccacc ctgcttttg 6780  
 45 t\_tgatgctt caccigtgt\_ tctgggta tgggtgacac agtaaatcct gtgtgattt 6840  
 taaacaccas aaataatgg atctgttgc t\_gtctcttt acgaatttca ggttccactg 6900  
 tgagacagaa ttcaattcac ct\_cagtccc tgagcacttt tgtgtgtct\_ aattctcta 6960  
 cgacaccate atgggagaag acaccgctgc aacctgcgg\_ ggctttctg cagaccacc 7020  
 50 t\_taaactgg tttctctc\_ ccaactggg c\_ggcccagg\_ actagcaaga ccacactctg 7080  
 calaggaag\_ aaaaagaa\_ ccccccaaa cclagallc\_ lllcllll\_ lclllcaga 7140

55

EP 2 336 318 A1

tcccccccc atcccccca agtaccsaag gatgttgccc gtgttgaata ca-gtgg-gt 7200  
 calclllcllc clccalagga laccgllglal ggaacclgcl ccacclcacll laccglaag 7260  
 5 acgaggaagg gcaatgtggc aacagaaata tccactgaaa gagacctggg gcagtgtgat 7320  
 cgttcaaac ccactccac aggcacacac ccacttctc tcacaaaag ca-ggtaagt 7380  
 cccatgtcag cactgtcgtg cacagcaagg agcatctct tattaataca attccagaac 7440  
 10 tttgagcta gtgggcaac ttgaggacag cctgcccctg ctgtttttta tacagaccag 7500  
 agataggacc ctgagcaggc acgggaaggt ctgcccaggc ttcacggcct gggatcagtt 7560  
 gagccaaggc ttgagtcagg ctctctctc ccagcccaga gctctgtctt tctctctg 7620  
 ctctgtcac tggaccaaa ctgctctaa tctcatcact tgagagtaat gactactcac 7680  
 15 c-clgagaag gllccgggca tggalglagg ccagcaaaa ccccllclgl cl-llllc-gc 7740  
 acaaggactc ctctgcccag ctccaagcct ctggcctttg aagaagtccc aagacctgtg 7800  
 tctccccct ctccctcact ccactgaagt gagtgactta gactgtccca gcttctgtc 7860  
 ctccacccc cagtaccacc ctgaccaac atggcccac tgcaccggc ctggagacc 7920  
 20 ctctctctc tgttaactgg ggccatggag caccatatta cctgagcctg ccagaccct 7980  
 gcaacatctt cctgatatg agcccagcc tgtctcagtg aacatgaata accgggcaa 8040  
 tcaactgtcat gctgggccc gttcctgtc atgtctctt gggttgaaa cagggact 8100  
 25 galgaccalg aglcccacag ccagaagagg a-aaalgcacl ggcllaggg cl-llllc-ga 8160  
 gcatctgctg tttctcaac ccactctgg ccagcaccac ggaagggaca gtggcagatg 8220  
 aaccatggc ctcccccctc gcatgctcc agtctaagc aggaqccaag tcaataaagt 8280  
 atacgtggta tactcaataa ggtgataagc tgaacagtgc agacaagsag tctgggct 8340  
 30 gaccaggaag gagaagaac tattcatgta cctcagcggg caacatttca tggaaagt 8400  
 ggagcaggaa ccaaaaaac gcaagaata tgaatgaa agagacatgt aagaatggc 8460  
 ttttggcaa aaaaaagtta ctgacaggt ctgtgaggg ctatgtgtg gcatggcat 8520  
 ctggaggata caaagttag acattgtcca ctgagggtgg aaaaagagg gtctacagct 8580  
 35 tgactcagct ttggggatc ccacttctg cccccctg tctaaatgtc aagtaccag 8640  
 t-tatctctt tctcagct- tatctagtgg tacaggactc ctgctccct- ctacctgaa 8700  
 ggtaaatgct tttaacagaa gatcacagga ctgatcaaaa tgcctgtctc caatctctt 8760  
 catagaccog cccctgtca actctgatca ccagcagcca gctctgtcag tacacactg 8820  
 40 acgclaaag gaagcalglg ccagaagcca clgcaaggc gcaacacccl lccclgcll 8880  
 tctctacaa gtaggctatg tgatgcacc ctgattgtc attaatggg tcaagtgtga 8940  
 ctgaacact ctcaagtct ctgttccag caaacctgt cctgggagg agaatcag 9000  
 45 agggataaac tgcggccc- cctgtccc- ctttttaagc gaacaggcca ttggcagaa 9060  
 aagtctagg catgcaaac actccagac caacaaaagc tatctaagac ccattctta 9120  
 agggctgtag atccagaaaa cctgaggatc actgcagggt accctgtta gaaaagctt 9180  
 catggaagat ttgggatac- gactggaac ttgtgtatc aatccactt tgaaacctga 9240  
 50 taatcaatga atatatttg agtaactgac atattcttg ctctatgttg tggaaatc 9300  
 gaaagaall lgaagcally cclagllcc laccllggc cclccagac laglggag 9360

55

EP 2 336 318 A1

tataaggcac gcatgtcttt ttgatgggag gataactagc gtgaccagga agaggtggat 9420  
g\_lallcall cagggccaac aalggclgga llaccocalg clllgaaaga lgggcaggac 9480  
5 tgggtagat gcacagacag ggaaaacctt caacatggaa agaatagtat gtcttgcca 9540  
tccgtgacat ggtcgtctt cttggttacc aggaataagt atggatggt accacaagt 9600  
acacagactt tgaacttga agacacacca aagatcaaca gcogcttctt tggtaaggt 9660  
aagagttct gtccacatag ttgotggaaa actactcaae gatggccta tcatggotta 9720  
10 gccacttgct gagccctgtt aatgtctgc tgactaaca gtgatacaga cactggtgtt 9780  
ctggctacct ctatgagaa agcaactca ttcctgatg tcaagttgca atggcataaa 9840  
ggaaaagaag ttccaaagc tacttaggca ttgtaaata gaaaactgga atcctaagtt 9900  
15 laacalgaca lallgalag aalgcacac acccalctlg lgalaagalc cagagclgcl 9960  
ccagacgagq tggaccaact qggagagaac ctccagagtc tggccagata gtaacctcag 10020  
gagtcagctt tttagaggtag aaggaactct aacaactctca agtccaaacc ttaccagta 10080  
ttgtattgta ttatctctg tccaaattcc ttctgtaca ttaacctat gtctttttg 10140  
20 ccalagcae cclgagcl cggclgglag agatlgall llalacctal lccacagag 10200  
agacagtga acagagaggc tttagagttg atgtagtca ggcgcagaa tcttagaggg 10260  
ggaaaataa gtgcaagctt gtaactaag ccaggactat tctcattaca ccacattcc 10320  
atgatgactt ttaacctct tctggcata ggtccagtc ggtggggag aggataaaa 10380  
25 agtgtctccc ctcccacaa gotgctggtt gacccaatta gaagaaatgg tgataagcac 10440  
ccatgtgctt ggtcccagtt gtaacctgt caacagtgc acctctcac caattatttc 10500  
aagctaaggg taacctgat atagactcag acaagtctgg attccactt agctctact 10560  
cttagacct gagagctct gggaaacct agttgctcat ctctgggtca cactctcca 10620  
30 tctctgggtc tcatctctt gtctctctc tgggactcag agctgagtc cagggatgag 10680  
caatttcat gcccacaaa ctctgtggt ctcaagca qgctqaatt taactataa 10740  
ttgaacaata atgcaacct acagggatag gatgatgagt cagtgaaac aagcaatca 10800  
35 cctatggcag agccagatc agcaggcatt gaatacagga tagtttctt ccttttccc 10860  
ctgtgctgat actccacaa ttccagcttc cagtagaca agatattggt gagatgaaga 10920  
aagctagagt tctctgaca ctctccatct tccaggtact aagaagatgg gctctgatt 10980  
tgagagcacc aatccacat cactccaaa gcaggccga gctgtttga agactctca 11040  
40 ggaacgaa aaccaacca lclclgagca aatalccag agagclaelc lclcaa\_aa 11100  
gotggttact gactgagag gctcagtg tgaagcagtc acatctctc tggcacagct 11160  
gattgaggt tccaggtat taatggtac agtcaactt ttataaac tcatgtaac 11220  
tgactgaact tccaaacct ggcaactgg agaactcag ggaccatttg gatatcaatc 11280  
45 cagttaatca attagtcaat cagttcatga tggctggata gagaactatc agctgctgcg 11340  
ctgagttcca tgaacacac acggcctac tgtgttcaag gcagctatg atttgtgtgt 11400  
taaacagae ggagaatag tccacattt tgatgggtae ctttaattc ctaggctat 11460  
50 tgcaggtgct ctccagaagc ttatagctg cgggagagag aactcagacg aaaaataaa 11520  
lalgalclcl claccclca aggcactggc lllaagclcl algaagglga gagaaggac 11580

55

EP 2 336 318 A1

tgaggccagg aatgagacc agctaagtgt gccagggcat attctgtgtg ctggccaaag 11640  
 gactlglala acagcllcl lglglclaca galccacagl cccclcllgg aacllllclc 11700  
 gattgggctt ctccgtggg taatattcct aaggaaagca tcatggttct gagctcaaag 11760  
 5 tgggttttg aagttagatn tgaatagtga atgaggtagat taagggtctn cctggcagag 11820  
 gacacaccat gagcaatatn ttatgtgcc tgaaggtagt ctgtataact ttatccatgt 11880  
 cttctctctc agccccatca ctttacaagc cttgggtcag tgtggacagc ctcagtctc 11940  
 10 cactcacatc ctccagtggc tgaacagtgt gcatgccaac ccccttctga tagatgtggt 12000  
 cacctaactg gtgccctca tcccagacc cccagcacag cagctgcag agatcttcaa 12060  
 catggcgagg gatcagcgca gccgagccac ctgtatggc ctgagccacg cgtcaacaa 12120  
 gggagtctc acacgtatn tctctccta ggagcagagg aacatcttgc aactctctgc 12180  
 15 atctctgtat taaaactgaa cccctcttc cactttcaa ctctgtctc tactcttctg 12240  
 tttttcttg atcatttttg gggtaatgac tgaanaaag aatcagcaa acacaaatg 12300  
 aatttttaa aataatttc ctacattata ttataaagt tttgaaat agcaaatgt 12360  
 20 acagaallic acagggaaaa cccclagaaa accagclalc lccclalall laaglylal 12420  
 tatatttct ttatcacata tacatccatc cattaattca tottatttct tgaagcatt 12480  
 caaagttaat tcaaacatc aacacacttt cccctaagta ttacagcttg catattata 12540  
 acttcagttc aataatagt agcagtttt tctctgaat tttttgttt gtctgtttg 12600  
 25 ttttttttg ttgtgttg tttttgaga tggctcact gtgtcacc caagctggagt 12660  
 cagtgatgca gtccaggtc actgaagcct caaattctc ggtgaagt atctccac 12720  
 cccagcctc tgaatagct ggaccacagg tcatgctac catgccctgg caattttg 12780  
 tattcttgg agatacagg ttcaccatg tggctcagg tagcaggtt tctcttctg 12840  
 30 gaaattttt ggtctttc tttttacatt ttatataaa tttatgtga acaagtcaa 12900  
 tttgttaca tgaatagat gtgcagtag taagtcagg cttcaggt atccatcacc 12960  
 cagacaacat atagtgtac cactaagtaa ttctcacc tccatctcc tccacttca 13020  
 35 cactctctg gtctcaattg tctatcttc cacacactat gtcttctgt gcacattatt 13080  
 tcactccac ttataaatga caacacgca ttttctct tctgtgactg tctgtttca 13140  
 ctaagacaa tgaactccag tccatccat gttgtgca atgacatgat tttattctt 13200  
 ttatggcca atagtattt attgctata ctttccat ttttaatca atcgtccatt 13260  
 40 galagacac lagglgal ccalglclll cclallglga alaglgclgl galaaacala 13320  
 tgggtgcagg tttctttg atataatgat tctttctc ttaggtatat acccagtaat 13380  
 gggattgtg gattatgt tagttctatt ttagttct tgagaaatct ctgtattgtt 13440  
 tccatagtg gttgactta tttcaactc catcaacagt gattaactgt tctctttct 13500  
 45 cgtatctc accaacaac gttattttt gcttttgaa taatggcct ccagactct 13560  
 gaaagatgt atctcattg ggttttaatt tacattctc taatgattag taatgttatg 13620  
 catttttca taccctatn gccatttga tctctctt tgaaaaaa gtctattcat 13680  
 50 gctcttgc tacttttaa tgggattatt tgggggatt tttgttgag ttgtttgaat 13740  
 tctctgtaca tccggatn tagtaccoca tggatgaat agtttgcaa tttttctc 13800  
 cactctcag gttaccacc tgttgattat tgttttact gtgcagaaac ttttaactt 13860

55

EP 2 336 318 A1

aattaagttc tatttgtcta ttttttgttt ttgttgtott tgcctttgag gtctttatca 13920  
 ogaattcttt gtctaggcca atgtccagag aagttttccc taggttttct tcttgcaatt 13980  
 5 tttatagtctc aggtcttata ttttaagtcct tgatccatct tgagttgatt tttttatatg 14040  
 gggacagata ggagccagc tttattcttc tgcataggc aatccatct tcccagacc 14100  
 acttattgae aagggtgtcc tttccctagt gcatgttttt gtcatttttg tcaaaagacc 14160  
 10 gttgactgta agtatgtgac tttattctcg ggttcagtat tctgttccat tcatctatgt 14220  
 ctctattttt atgccagtac catgctgttt agattactat agccttqttg taataatcga 14280  
 agtcaggtaa tgtgatgcc ccagctatgt tctttttgct taaaattgct tcagctatct 14340  
 aggcctcttt tggatccac atgaatttta taattatttt ttctaattca caagtttggg 14400  
 15 ttttaagaca aacctaacg gggttaccaa gctctgactc tcttctctta ttctgtagct 14460  
 atcataagac aaacctaca gggaccagg agctgctgga cattgctaac taactgatgg 14520  
 aacagattca agatgactgc actggggatg aagattacac ctatttgatt ctgogggtaa 14580  
 20 tctcagctcll llalalgaca lcalcllll cagaagcacl lllcllggac accllllact 14640  
 tccctctctc gcaacctgac ggttctctgt tctttttctt caatgcaggt caatggaaat 14700  
 atgggocaaa ccattgagca gtttaactca caactcaagt ctccaactct gaaatgtgtc 14760  
 caaagtacac agccatcac gatgatccag aaagctgcc tccaggtctc ggggaaaatg 14820  
 25 gagcctaaag acaaggtaaa gtccaccaga agaggtctgc aagtgaactt ttattaacaa 14880  
 ggatttggaa ggtactaggg gaatgagact ctagatttca tctactgac ttattctgct 14940  
 gtttcttcc tttcttccc tcttctctc ctctctctc cctccctcc ctctctctt 15000  
 tcttctctc ctctctctc tgcagatgga atctcactct attgccagg ctggagtgca 15060  
 30 gggcatgat ctggctcac tgcacttct cctctctggg tccaagcaat tctctctgcc 15120  
 tcagcctcct gagtaactgg gattacaggc atgtgcatt acaccagct aatttttgta 15180  
 ttttttagta gagaaggag tttgccatgt tggccaggct ggtcttggc tctgacctc 15240  
 aggtgatccg cctgctcag ccttgcaag tctgggatt acagggctga gccactgac 15300  
 35 cgggcclcll clglcllcll allgcaall tcaacaagc lallgacttg actgcllclg 15360  
 agtatgtgac gtgagagaaa tacttgactt tctgtctatg tcaacatgca gaactgaga 15420  
 tgtttttgct tctacctgc cactaccag atgaccato cctctcatca tggaaaaaca 15480  
 40 tcttaattt tcccccaata agcttaggct aggatagcca acttggccc ctcttaggtg 15540  
 caaagactcc agaactttg aaactacct atttattagc cccaaactct tactaccct 15600  
 tctcatcttt atcccacac taaaataact tacgttaaaa caacttgatt ttcacttagt 15660  
 ggtgatctc caaacaatc acaacttgg cataatttat gtgtttteat ggaattgaat 15720  
 45 tcaacaggca tccacaggc tttttctggg aaccttact tgatagtgc ctaggaaaca 15780  
 ctggcaagaa gattcaatac cagcattga agaacgatta cagagaaatt agacctgac 15840  
 taaqaaaqa gtaqcaqac aatgccagt tttgccaggc atgttctgtg ttctgaccac 15900  
 50 aggacagtga taacctctc ctcttttgac tgcaggacca ggaggttctt ctccagactt 15960  
 tcttgatga tctctctcg ggagataagc gactggctgc ctatcttatg ttgatgagga 16020  
 gcttccaca ggcagatat acaaaattg tccaaattct accaaggaa cagaatgagc 16080

55



EP 2 336 318 A1

aagtgaagac ctttgtggc tcccatattg ccaatatott gaactcagaa gaattggata 16140  
 tccaagagta agtaagagc attocacca tataccactg agggccotga gcgggaatc 16200  
 5 caaccotagg ttttggcata gccactgtct gcccttgctt ctgaaacaaa cacttgtgca 16260  
 aatgtgtagc agatctagac ccaaagactt agggtaaatg aatcaagac attttggtag 16320  
 tgattggaaa tccatattta cttgggggtgc aagagtcaaa ggataataac atgggtgtgc 16380  
 10 agctcaaaat atacctcttc ttatctagtc tgaaaaagtt agtgaaagaa gctctgaaag 16440  
 aatctcaact tccaactgtc atggacttca caaaattctc tcggaactat caactctaca 16500  
 aatctgtttc tottccatca cttgaccocag cctcagccaa aatagaaggg aatcttatat 16560  
 ttgatccaac taactacct cctaaagaaa gcatgtgtac aactaccctc actgectttg 16620  
 15 gatttgcttc agctgacctc atcgaggtaa ggtggaagag tttgaggttc totagcccat 16680  
 tttgtacagc atcataaana gagagtccct gggagccagg agctaccocag aggaaaacta 16740  
 agaaccacca ggcacttccc aocatgattc tgagggtttc ttctctccc ccttcccgc 16800  
 cttctctct ccccgctagg ggtccactga agcatgactt cttaacatta atagaaatgc 16860  
 20 aggcctggcg aggtggctca ctctgtaat cccagcactt tgggaggccg agggggctgg 16920  
 atcatgaggt caggatatcg acaccatcct ggctaacaag gtgaaagccc atctctacta 16980  
 aaaatacaaa aatagcccg ggcgtggtg caggcacctg tagtcccgc tacttgggag 17040  
 25 galgaggcag gagaalgcg tgaaccagc aggcctgagc tgcaglgagc cgagagallg 17100  
 cgcactcgc ctcagcctg ggcgacagag caagactcca tctcaaaaa aaaaaaaaa 17160  
 aaaaaaattg aatgcaaat gtctcgtctt taagtccaa agcgaaggaa gcatatgtc 17220  
 tgctagtca gatcgtctc asatctcaaa tcactccaa ctctgaatcc ttgttgaat 17280  
 30 tatttgcct atctgaacc tagctgctc ttctagaaa aagcaagtaa taaggtaag 17340  
 atctagtga gatttaata aagcagctcc tgtgaaatgc taaggtaagc tctggcctg 17400  
 tggattcac atactgttt agataaagg acatcaagag tgggactac taggctgca 17460  
 tacaacaaag aaactgatg ccattttctt gctgatttt ctctctcaga ttggcttga 17520  
 35 aggaaaaggc lllgagccaa callggaagc tclllllggg aagcaaggal llcccaga 17580  
 cagtgtcaac aaagctttg actgggttaa tggtaagtt cctgatggtg tctctaagg 17640  
 cttagtgga cacttggc ataccaaga tgataaacat gacaggtgt gtatttga 17700  
 40 agtatctct taaggaaagc tttgggtctc aatgcaaaa caattcttt ctaagcaag 17760  
 aagctctca aataactatc aactgaaggg ataactatgg tttttatca ccagaccgc 17820  
 tgggtaag gccagtatc tctgcagtta aagatctct gaattcagtg tcccagaaa 17880  
 ccagactcac aataagtae ctaggataac aagagtatga actctgggt ggtgtggtg 17940  
 45 gttcatgct gtaaaccag cactttggga ggccaagggt ggcagatcac aaggtcagga 18000  
 atttgagacc agcctggcca acatactgaa acccgtctc tactaaaaa acaaaaaaac 18060  
 taqctggca tqtagtgc tqctgtaat cctagctact cgggagctg acacagaga 18120  
 50 atgtcttga cccgggagc ggaggttga gtagcogag atcacgcgt tactctcag 18180  
 cccgggtgac agtgtagac tgtatctaa aaaaaaaaa agtatgaact ctggcatag 18240  
 attaattct aactccctg tottgaagct ggcgcactt ggggaagttg gttgatatta 18300  
 55

EP 2 336 318 A1

5 tgtgtatctg tttctgtctg tatcccagac tactaataac agtccaaacc tcacaaggtt 18360  
 atttaagac aatgaaataa ggcattctaa atgccaaagca cagtgcctga tgcctggcatt 18420  
 ggttgttcaa taagcagaca ctattaacag tctctaaatta atatcttcaat tattattaac 18480  
 tgctgtcttt ggcctctcac cccatcagtg cactagcaaa tgagaccaaa ctccacattt 18540  
 gaagctagca atgagcccc atttaaggag ggaataggt tgtatgatct ggagcttatt 18600  
 ctgaatttt ttgcaccca aagtgtggtc tggtcagaaa tacagcttct caagcttcac 18660  
 10 ccacaatcta ctgaatcaga agcgcatttt agcaagacct catgtgactt gtatgcacat 18720  
 tcaactttgc agagcaaggc agtaatttac cctccaggc tcaactgtga gcacgagctc 18780  
 catcttctaa tttctgacc cccacttgag gcagaggatc tttgatctgc ttgagctctg 18840  
 15 tcaglllccac allllllll lcccaalgcc tgggcalcca lclclgaga lclclllcl 18900  
 tctgagaaqa acttqtctag gatcaagtgt tttctaaact tctggtgat ttatataaca 18960  
 gctacatttt ctaagaaac aactttagt ctcaactggt caaagaagag aaggctaagc 19020  
 agggaaacggg tgggggatag aggatcttct aatcttgagg atcctggcat acgggagaat 19080  
 20 agggacccct cctccatcc caccacatct tactatgtct acagattttt taattaagaa 19140  
 tagctttagg agtgccacta tccctgacaa gaccttagtt cttaactctc tgccttagagg 19200  
 aattagcctg gactccagtg tctccctggt cctcacctgg agcatttttt aggccatcc 19260  
 25 tggclgcalc agacagglcc cacallggga acgaaaggl glllgacal gclgacalc 19320  
 cactggccat tttactacta aactctcagg atatggtaa tggaaatag ctcagctctg 19380  
 agaaqtgat taaagatttg asatccaaag aagtcccgga agccagagcc tacctccca 19440  
 tcttgggaga ggagcttggc tttgccagtc tccatgaact ccagctcctg ggaagctgc 19500  
 30 tctgatggg tgcctccac ctgcagggga tccccagat ggtaatcag caggcccccac 19560  
 tggggccca tgagaccaga cgttggtttt ttttagatc gccagactc cctaccgac 19620  
 ccagctgca aagcccga aagatgctgt actttctca gagatggagg ttgccttga 19680  
 atttactga agatgactc tggatccat ggaatgta acatttagaa ataaagctat 19740  
 35 lcalaalgll agclglall llaagagcal laalllall alclggaaa caalgllcgg 19800  
 tataccttc ctacctttg ctgaaggtcc tttattttt atttttatt ttttaattt 19860  
 ttgagatgga gtctgctcc caggctggag tgcagtgata caatctcgc tcactgcaac 19920  
 40 tctgccttc gggctcaagc aattctcctg cctcagctc ccaagtagct gggactctgg 19980  
 acgtgcacca gcctgcccg ctatcttctg tatctttagt agagacaagc ctgttgaca 20040  
 ccattgcaag ctggcttcca actcctgacc tcaagtgac cctcagctg ggcctccac 20100  
 agtgcctgaa taacaggtc ggcactgc acctgacctg aaggtcttt taagattgaa 20160  
 45 atgatacaat gattataaaa gaaagtattt ggcaactat aattcactat ctaataatgc 20220  
 tataattttt attactaat ctaaaaagga aatataaaa tgtactcta tgccttatt 20280  
 aaaaaaatgt tgactttaag aaaaaggtc tcaagctatt ttatgaaa attatttaa 20340  
 aaataaaac caatgcaaa tgatgttac atcatctca taggcctttg gttcaaaa 20400  
 50 atgtatttt taataatata atacatttca agtacactt caettacagt cagactccag 20460  
 aacaccaga llaagcalg gcalalalga lacllaaagl ccalaagcl ctgaggcca 20520

55

EP 2 336 318 A1

gcaatattct taagagcctt ctgagtcacc ttgaaatga catgatatct atctagtga 20580  
 a.LLcLLaLa Lcclgallca clgaaaacgg laaaaacalc aglllgalcl LLaLLLaLca 20640  
 5 aactattcag ctcaLcaaaa tatgctagtc ctctctttcc agataaagag gaattacct 20700  
 ccaatgtatg gqacgttqta attaacaaaa ccgactttaa aaacacttac ttttatttgc 20760  
 tctcccttgt tggctctaca gattggagag gcatcagga agggctcaaa gaatgactt 20820  
 tttcttcaact acatcttcaL ggagaatgoc ttggaactcc ccaactggagc tggattacag 20880  
 10 ttgcaaatat ctctcatctcg agtcattgct cccggagcca aggtctggagt aaaactcgaa 20940  
 gtagccaaacg taagattctg tttgcctttt gattttcttag gttattactt tctccaggg 21000  
 tgcatttctt gttaaaaacL atttaaaaat ggttttccac ttcaagacaa aa-gcttcat 21060  
 15 callglaalc accLcallaL LLLLLLaLga aaaacLLcaL gLLccacca gaalgcacl 21120  
 cctcactagc tccagtatg gtatggccat aagacaagaa ctcaqttctc tcaacaatg 21180  
 agtattctca tcatcttttL aatctggttL tgcctcactg taactcaggt gctttctagt 21240  
 tctgggtagt atacLccaaL tctagagaac tgagaactcg ctctcttct tccaaaacaa 21300  
 20 tcccagtaat gtttccaaag gctctagtta tccaggaaat ctctgcccgg aggtgagaaa 21360  
 ggtggttga tctgactgac aggggactga agtatttaat gaactctaat aggttgttL 21420  
 ctgacttata gatgcaggct gaactggtg caaaaacctc cgtgtctgtg gactttgga 21480  
 25 caaalalggg calcalcalL ccggacLLcg claggagLgg gLccagalg aacaccaacL 21540  
 tcttccacga gtcgggtctg gaggtcactg ttgocctaaa agctgggaag ctgaagtta 21600  
 tcatctcttc cccaaaagaa ccagtcagc tctcagtgg aggttaattc tLcagccaa 21660  
 gctgcttag ccagttgaa agagagaaca gagaatgtac ctgcagaatt ttgccaggct 21720  
 30 aaacagttga ttgagatcaL tcaagtcctg aggaagcagg agaggagttag aaaggaaaga 21780  
 tccgggtta ctatctttaa tcttagccta gacttactac ataactacat aatLaccctt 21840  
 ctctactttL tcaactttta ctaaaactgc ctttatcttt ctgctttgag acttattaag 21900  
 acctactgct taatagttL ttattaagtt ctgatttttt gttatctatt tcttttga 21960  
 35 atgaagaaac aatagctctg gtagatcat ctctggaaaL ttaaLatttt ccccccaaa 22020  
 aaatacctaL gaacatattg atttgaggta gctaggtagg taaagcatga aactcctaac 22080  
 ctctgataa tggaaacag cctcttttgg agagttccat ttt.aagtggc acctcaacc 22140  
 atgattttg cttagttttc atattttaga cacattcatg tgttcattca aaaataat 22200  
 40 LLaallggcc agccacggLg gLLcalgccl claalccLag caclLLggga gccccaegLg 22260  
 gatggatgc ttgagccctg gtgtttggat accagcctgg gcaacatggc aaaaccocat 22320  
 ctctacaaaL aaaatLaaatL aaatacaaaa atagccagt cgtggtgqca catgctgta 22380  
 45 gctccagcta ctcagaaggc tgagatggga ggatcaactg agcccaagag ttcaagcctt 22440  
 cagtgaacca tgcctgcaac actgcactcc agcctgggag acagagcaag atctgtctc 22500  
 acaaaaaaca aaaatagta t.ttttaattg cctaataatL accacgtatg ttgagtga 22560  
 cacacaaggt cctgacctL tgaacgctta cattttataL gggagacaca caattaagca 22620  
 50 agcagtaac atagagtaag gctaaagtta tagaaagtat tagagtacca tgaatttL 22680  
 Lalcalglag cclgclag LcagggcaLg callclgaag caagLgLacl Lgacclgala 22740

55

EP 2 336 318 A1

actgaggact gtgtcagagc ctttaggca aaggagaaag gagtgagtgt tccaggcaaa 22800  
 aggaaaagca tglalaggcc tgaagglaaa ggaalalggc lcaaggaaac ggaagaaglg 22860  
 5 cagaatggta aggggctcag agatgatggg gagaggtagg caggggagag agcatgccca 22920  
 cctgcgaaaag ccatcctaag gactttggac tcttttgaag qcacaggaqt tcaaaaacggg 22980  
 agcagaaata agataggggc gatgttttag aagaaatact ctgactctag tgggaagat 23040  
 10 gggtgagaag gaggcacagc tggacacgaa gagaccattg gacatctctt acgatccat 23100  
 ggggctaaga gctcataatg gctgcagtg gagaaaagcc aggtatagaa aggagtgcgc 23160  
 agattctaca acttctaag aggcagaatc acaagtactg ggtgattaac tgggtatggg 23220  
 gacaaggcaa aagaaagaag acaagaggaa gaggggccc ttcattttaa taagaactac 23280  
 15 aglgggagag clccgglll caaggaaagl cacaaallca gllllggalg tclglalll 23340  
 gatgtcctcc tatcaacaa ccaqtttaqa aatctaqtq tcaataqac ctatqgact 23400  
 gagcccagta aagaggcttg gctccacat atggatttgg gaatcattag tctacagagg 23460  
 20 tggttgtggc taaacagcaa ctggtataga gtagacatg agagatgagg acagaaact 23520  
 ggagaagaca aacalalaaa ggaagaaggg caalaaccag caalagalla gaagaagga 23580  
 ccagagaagc agaaggagaa ccaagccat aaaaggctac agaagccaaa gagcagccac 23640  
 agggagatc acccatggg taggcgaaa gggcattag gactccaqca catcagcaaa 23700  
 ctttggctct gggccaccoc caacttgag aacaatact tggaggacaa tggctactt 23760  
 25 caaagaaagc atccctagaa caaacccggc caatgttga cttctctca tctactaagt 23820  
 ttttaagta acacttgaa ggaaggctgc acatctctt cagaatgag aggcctcagc 23880  
 gcttagtct ggtcatgagc gcgcaactct atggaaggct tctgggaggc caaggaagat 23940  
 30 gaaacctaaa tatgcccatc gtagtgagga gcaaggaggc cattagagac atgatgaaa 24000  
 gcattttcag gagaaggagc gagcagtcag agcacattgg gaggaagtag agactgcaaa 24060  
 ggcagacaa ccttqatgc gaggaaatg aaaaagcaag aaaaagaaqa aaggagata 24120  
 ggggaggggc acaggggaag agacttgagc gctttaatg caggtggaag gaagcagta 24180  
 35 gagagtaggc gattcatac gcaagagaca gttctcttg cctgcattg taggaagaa 24240  
 ggggcacact gaagtccagc ccagtgatc agctatttaa catctctgag cctctgctc 24300  
 tgtaaaatga gaaccataag cctactgttg tgggattac aggtaacaga tggaaagaa 24360  
 40 tcagccagaa gcttcagagc cactctcatg gcttgcatg ttgatgtct tctaatatt 24420  
 allgllllc caglaaalla aalagllaga calagglglg gaclgaggga agacagcagg 24480  
 acaaggggt atttgaccoc tgagaattg tgatgtccat tttgatccat gacttgcca 24540  
 taactcaggt atttttgttc tccaccagca acacattaca tttgtctct accaccaaaa 24600  
 45 gggaggatg cccacctctc attgagaca ggcagtcctg gtcagtttgc aagcaagct 24660  
 ttcctggcct gaattactgc acctcaggcg ctactcca cccagctcc acagactccg 24720  
 cctcctacta tccctganc ggggacacca ggttagagat gctcagtgcc tgaccagca 24780  
 tttctcacc tccacatca tggccacctc gcatggcaca ggaaaaaata ctctgtctg 24840  
 50 taagaccctg tcaatagcc tctgggttg caccatctt ggtatttaa agcaggctcc 24900  
 tclggccaac acalggglg tcacllllg clclcllglg calgggalgg galcacagca 24960

55

EP 2 336 318 A1

cagatcccaa tttgctocta attcagtgtc catgtttctg agcctccaga cccatcgcta 25020  
 lgagcllcccl ggagcccacc aalglgcllg aagccllccac cglaccllagg lggclccclg 25080  
 5 ttttcagccc ccaagttcca gtgcttgctc tcagctttgc tgaacaacc agccaaotcc 25140  
 tgctctgctt gtccaaagtc ttgggaatcc ttgtgtctgc ccttgcoctg gctctctgta 25200  
 ggactgaggg atcaaaaaga tcactttagt taagggaag agacaatggt aaaataagga 25260  
 ccatattttt gttgcatttg aggctgaatt gttttgggaa cataatcacc atccttgaaa 25320  
 10 gctctaacat tatgcactgc cttoattgta atgtctttag attagagctg gaactgaggc 25380  
 ctacagqaga gattgaqcaq tattctgtca ccqcaaccta tqagctccag acagagcaca 25440  
 gaggccttgg ggataccctg asgtttgtaa ctcaagcaga aggtgagtat tcaaaaacaca 25500  
 gctgcctcat ctctgctcgc agtctcaggt tcagaattcc tgaggagaag acatgtcatt 25560  
 15 taacctatft aacaaatagg ttaactgagt acccaactaag cggcaggcct atctaaagac 25620  
 ctgggttaac tgagtaacca ataagcggca ggcctattct aagacotggg gctagaacag 25680  
 tgaacaatgg agtctctgoc ttcatggaag ttcagtgaa caaccaca agtaatat 25740  
 20 lggaalalca galaaglac ggaggagaaa acagagcgle gactlglcla lggaggccla 25800  
 ggagtaggag ggaggaagaa gggcaggaa agcagtgcat ttggaataat aagggaagt 25860  
 ctccctqgta aagtgaqcat aagqagcct atcaaaaata agagqagaag cctgtctqta 25920  
 agactgttac caggcagagc gccacagcag tccaaggcc ctgaggctca cccactccta 25980  
 25 ccatgtttcc aggaaggaa ggaagacagt atggctggag cagaaagccc agggagaaa 26040  
 gaggtagaag atgaggacag agagatatgg agaggtgaag gaaggataat ctcataggcc 26100  
 atggtaagaa ctttgcttt ttctatgaat taaacgaaag ccattgggga gtcctcaga 26160  
 ttgtatftat gtttatgctt agaaaagact atgggcagac aagggcagag aaactaatat 26220  
 30 ctaggttatc acaataatcc aggcaggaat cagtgttctt ttggatcagg gcaatgacag 26280  
 aaqagatatg aqaagqqaat gqattctgoc catatfttga agattagcct gacaagattt 26340  
 gctgatacag tggatgttca gtgtaagagg aaaaggggaa tgaagacaaa cctaaggttt 26400  
 35 tggccocggg caactgaaaa atggaacttc cattttattgc gatggaacgg gctactegag 26460  
 gaggcaggtt tagggaatcg gagaaatfta ggtgttcact ttggaaaaaa aatataatag 26520  
 ggatagcag gagcaggttt tagggaatgg ggcacattta ggtgttcact ttggaaaaat 26580  
 ttttatatag ggatagcata tcacagaatt aaactaggaa gaaaatcca tgatagaaag 26640  
 40 cactlgaggc gcagggcaag clggggcaal aglglcllgl aaacallgl llacgaagga 26700  
 tataaaatgg accagoccat ggattgaagg acgcccggga atcttgttac aaagaaagg 26760  
 ggagttgggg agatggagcc caggccaagg ccagcaagga accaggaag gcatcttggg 26820  
 tagaaagtae tatagagatg tctgtcttc ctggcccaga agggctgoga gcttttctg 26880  
 45 tcccaaaaac aagcaagtg ctcccatftt cagggccttt gcatctctga cctctgct 26940  
 ggaatgtgct cctccagaa ctcagcgtgg ctccaacctc tttcattct ggtctctgcc 27000  
 cacatgtgcc cttaacagag aqaattctc tgaccacca gatatgaaata acacttctc 27060  
 50 tatcccttcc ttttatoctc gtatccagtt ttaactctct tcataacatt caataaccac 27120  
 tgacatgagc aagtacttg tttattgctc ctacacctcc cccactagaa ggtaaagccc 27180  
 atgaaagcaa ggatcccca gtaccaagag cagtgcocag cacacaatag gctcataaca 27240

55

EP 2 336 318 A1

5 ggcaatccat aaagacttgc atacatgaac acaactgagt ttaaaattat cagtaaaaga 27300  
 gacccattaa aaaattttaa tgagaacaaa aaaattoagt aaaatcctga acgtgtttt 27360  
 tgtttaagca cattgattcc ttggagtttc tetacctttt cctccttttc cttccaaaac 27420  
 atagcttctt tatttattta tttatttatt tgtttgttta tttatttatt tatttatata 27480  
 ttattttttt gagatggagt ctgcctcttt tgcccaggct gcagtgcaagt ggtgccatct 27540  
 10 cggctcactg caagcccccg ctcccgggtt catgccatto tctgcctca gctcctgag 27600  
 taqctqqgac tacagqcaac caccacacgc cccqgctaata tttttgtatt tttaqtaqag 27660  
 acggggtttc accatgtag ccagaatggt ctgatctcc tgacctcatg atctgcccgc 27720  
 ctggcctcc caaagtgtg ggattacagg tgtgagccac cgcaccggc ccaaaaacata 27780  
 15 gcttcttacc acacatctcc tgattctctt atacactcgt ccaggtgoga agcagactga 27840  
 ggctaccatg acatcaaaa ataatcggca gagtatgacc ttgtccagtg aagtcacaaat 27900  
 tcoggatttt gatgtgacc tggaaacaat cctcagagtt aatgatgaat ctactgaggg 27960  
 20 caaaacgclcl lacagaclea cclggacal lcagaacaag aaaallacly agglcgcccl 28020  
 catgggccac ctaaggtaaa gaaggccgag ggtcatctga cctgcactgc aggcctgggt 28080  
 ggttcttttc attatctctc ttcaacttca tacctgacca agccatgttc tccctagtc 28140  
 tacaatcaga gtggcagaga gggccctcaa caattttttt tttttttgag atggagtctc 28200  
 25 actctgtcac caggtcggag tgcagtggca caatctggc tcaatgcaac ctccgcctcc 28260  
 ccagttcaag tgatctccc gcttaagcct cccaaggagc tggaaactata ggtgcatgcc 28320  
 accacacccc gctaattttt atatttttag tagagacagg gtttaccat attgaccagg 28380  
 atggctctga tctcctgacc tctgtatcca cctgccttgg cctcccaag tctgggatt 28440  
 30 acaggtgtaa gccactgcac ccggccaagc tctcaacatt ttaacctct ggcctatcc 28500  
 agttggattt tctaccatc tctcaggcac ttaactattca tgtatcaagc acagtgcctg 28560  
 g-gctttaa gaaatatac cggctctcac aataaactgc gaggtcactg tgagtttcc 28620  
 tgtttcatgg ataaggaaa ggtagctcag aggggttaaa tcaattggtc acaatcacag 28680  
 35 agclaglaaa lagcagaga ggalccaac agllllcaa aacllclcl lclclclaaa 28740  
 cctgtttgca aagtcctaa tttgtctga atgttgctt tagaagtga tgagttgat 28800  
 ctgtggctgt ttctctgac cctccttcta tctggttttg atcaccocaa atggaaotc 28860  
 40 tgtttaatcc tgcaatctc cattgaaagg acaaaatcat tggtgccaac tgattttctt 28920  
 taccatagtt gtgacacaaa ggaagacaga aaaatcagg gtgttatttc catccccgt 28980  
 t-gcaagcag aagccagaag tgagatctc gcccaactgt cgcctgcaa ac-gcttctc 29040  
 caaatggact catcctac agcttatggc tccacagttt ccaagagggc ggcctggcat 29100  
 45 tatggtatgt gtctcttccc ctgtgtgagc acttccaaag taatgcaggt gt-gagacct 29160  
 g-ggttacag gctcaactag taccattcac aactatttcc tacgtatttt cagatgaaga 29220  
 caagattgae tttcaatga acacagccac caatgtagat accaaaaaaa tcaactocaa 29280  
 50 t-tccctgtg gatcctccg attatcctaa gagcttgcat atgtatgcta atagactcct 29340  
 ggatcacaga gtcctcaca cagacatgac t-tccggcac gtgggttoca aa-taatagt 29400  
 tgtaagtatg agtc-gccag tcaataaata catggatata agtctaatc acatcctcaa 29460

55

EP 2 336 318 A1

5 cctgagcta ggtgcaggaa ggtttccaaa gatgtataag goatgottcc tccccccag 29520  
 ggaattcttg gggagaaaaa azaactttca caagtgtgta gttaccocagt tacacaaagc 29580  
 tgaatgtgat acatatcaaa gagatgctac taagtagaac agttctttgc ctagtggat 29640  
 caaaggaagc ttcaggacac cagctaggag cctgactatg ttagacattc ctttataaa 29700  
 tatggacagt gatcagtgac tggcaacgaa gattcataat tttctgttat ttatttttaa 29760  
 10 cttcagtgcc attgcccagc ttaataatta acttgtcaaa tcggattttt tgcctaagt 29820  
 tcattgctct ttgaagctca tccaagccca tacccttaa aatctctctt caatttcag 29880  
 gcaatgagct catggottca gaaggcatct gggagtcttc cttatacca gactttgcaa 29940  
 gaccacctca atagcctgaa ggagtccaac cccagaaca tgggattgcc agacttocac 30000  
 15 acccagaaa acctctctc aaaaaggtaa aagaagaaag cagcaaggct totgaacca 30060  
 tcaaaagtaa atgaaagatc ttaacatagca tgatttagac atttttttaa atttttaag 30120  
 gaaataatth aagcatttta aggagattaa taactatagc acaaacactg tggcatcttt 30180  
 gcattagtaa acatgagaac accaaccttg tcaggaagaa tctaagaag toattagagg 30240  
 20 atctcgtac tttcaccta agatatttta ttcagtaca cctgttataa gcaattctc 30300  
 cctctgactg tgaagaattc agaatggcta gaggcgttat tgactacagg ctgctgta 30360  
 agctagagag agtcagaaca gccattgagc actaaatgga ggcagcattc tgagaaata 30420  
 cllaaccca ggclactga cllocalacc talgllclll ccacaaalca agllglclca 30480  
 25 atcagttta gcaattttg atcaagtac cctatgtgc aaaatgctag actaggtaca 30540  
 ctgagaagat aqaaactgg taaggtatag ccttttcttt caagaagata ccattgagac 30600  
 acaacaaat gagaataac taattatata agcaaaatta tgacatgctc ttgagaaag 30660  
 30 ctgcaaggga ctatgtaac gtaagaatga gacaaattgg ctatgactta gctgggatgg 30720  
 taatgataag gactggccc tagaagagct tctcaggat ttgagtgttt gacaggtgga 30780  
 cgtaaaagca aaggggtcca gccataggag tagcacaag aaaagtgcag agtggcttg 30840  
 ggaatggggc aagtacaata ttgtgtgtaa ggtcagaggc agagaacttt gaatgactga 30900  
 35 lgclgacly lgggaglyl alclllgllg lcalllcag cgalggcccg gcaaalala 30960  
 ccttgaaca gaacagttt azaattgaga tccctttgcc ttttgggtgc aaatcctca 31020  
 gagatctaa gatcttagag actgttagga caccagcct ccacttcaag tctgtggat 31080  
 40 tccatctgcc atctcagag tccaagtcc cacttttac cattccaag ttgtatcaac 31140  
 tgcaagtgcc tctcctggc gttctagacc tctccacga tgtctacgc aactgtaca 31200  
 actggtccg ctcctacag ggtggcaaca ccagcacaga ccatttcagc ctccggctc 31260  
 ctaccacat gaaggctgac tctgtggtt acctgcttc ctacaatgt caaggtgagc 31320  
 45 tatgctcag taaaggtgc accgggctag tccatggcag gctcaagag gagagctcc 31380  
 tccagggag aaagactt ggctttctag cagataatct tcttctac ttggaactc 31440  
 ttattttat tcaacaaata gsaatattta taaacatat caoqtgtatt aaatattca 31500  
 50 ctaggcagta acagaaagta gacagataag ccagcaatta taattcagtg tgagaggc 31560  
 tatgataaag ttagtatac aagtatacag tagagtggac gcactcaaca agggaaacta 31620  
 acaaaagcct gtgggtgca ggcaaggct cctggaggaa tgcctttgc tatcagattt 31680

55

EP 2 336 318 A1

tatctttgca ttacagatgg aggagtctat tgcacaattg gccacagaaa atggggcttt 31740  
 atattgaaa gactttcaac atagagattg cctggaaat gtactgotta atttaacaa 31800  
 5 tgtcttttca tttttatgtn aggatctgga gaaacaacat atgaccacaa gaatacgttc 31860  
 acactatcat atgatgggtc tctacggcac aaattttctag attcgaatat caaattoagt 31920  
 catgtagaaa aactgggaaa caaccagtc tcaaaagggt tactaatatt ccatgcatct 31980  
 agttcctggg gaccacagat gtctgcttca gttcatttgg actccaaaaa gaaacagcat 32040  
 10 ttgtttgtca aagaagtcaa gattgatggg cagttcagag tctcttctgt ctatgctaaa 32100  
 ggcacataty gctgtctty tcagaggat cctaaccactg gccggctcaa tggagatcc 32160  
 aacctgaggt ttaactctc ctactccaa ggcaccaacc agataacagg aagatatgaa 32220  
 15 galggaacc cctccctcac cctccctccl galctgcaaa glygcaltcal laaaaalcal 32280  
 ccttccctaa agtatgagaa ctacagactg actttaaaat ctgacaccaa tgggaagtat 32340  
 aagaactttg ccactctaa caagatggat atgacctct cttaagcaaaa tgcactctg 32400  
 cgttctgaat atcaggctga ttaogagtca ttgaggttct tcagctgct ttctggatca 32460  
 20 cttaattccc atggtcttga gttaatgct gacatcttag gcactgacaa aattaatagt 32520  
 ggtgctcaca aggcagacac aaggattggc caagatggaa tatctaccag tgcacagacc 32580  
 aacttgaagt gtatctctc ggtgctggag aatgagctga atgacagac tggcctctct 32640  
 25 ggggcalcll lgaallaac aacaaalggc cglctcaggg aacacaalgc aaaallcagl 32700  
 ctggatggga aagccgccct cacagagcta tcactgggaa gtgcttatca gccatgatt 32760  
 ctgggtgtcg acaccaaaaa ctttttcaac tccaaggta gtaagaagg acttaactc 32820  
 tcaaatgaca tgatgggtc atatgctgaa atgaaatttg accacacaaa cagtctgaa 32880  
 30 atgacaggct tatcactgga cttctcttca aaacttgaca acatttaacg ctctgacaag 32940  
 tttataagc aaactgtaa ttacagcta cagccctatt ctctggtaac tactttaac 33000  
 agtgacctga aatacaatgc tctggatctc accaacaatg gaaactacg gctagaacc 33060  
 ctgaagctgc atgtggctgg taacctaaaa ggagctacc aaaataatga aataaaacac 33120  
 35 auctalgcca cctcctctc tgccllclca gcaagclala aagcagacac tcllgclaa 33180  
 gttcagggty tggagtttag ccatoggtc aacacagaca tgcctgggt gcttcagcc 33240  
 atgacatga gcacaaacta taattoagac tcactgcatt tcagcaatgt ctccgtct 33300  
 40 gtaatggccc cgtttaccat gaccatgat gcacatacaa atggcaatgg gaaactogct 33360  
 cctggggag aacatactgg gcagctgtat agcaaatcc tgttgaagc agaacctctg 33420  
 gcatttactt tctctcatga ttacaaaggc tccacaagtc atcaactogt gcttaggaaa 33480  
 45 agcatcagty cagctcttga acacaaagtc agtgcctgc ttactccagc tgagcagaca 33540  
 ggcacctggc aactcaagac ccaatttaac acaatgaat acagccagga ctggatgct 33600  
 tacaacacta aagtaaaaat tggcgtggag ctactggac gaactctggc tgaactaact 33660  
 ctactagact ccccaattaa agtgcactt ttactcagty agccatcaa taccattgat 33720  
 gctttagagc tgagagatgc cgttgagaag cccaagaat ttacaattgt tcttttga 33780  
 50 aagtatgata aaaaccaaga tgttcactcc actaacctcc ctttttttga gaccttcaa 33840  
 gaalalilly agaggaalog acaaacall aagllglac lggaaaacgl acagagaac 33900  
 55



EP 2 336 318 A1

ctgaagcaca tcaatttga tcaatttga agaaaataca gagcagccct gggaaaactc 33960  
 ccacagcaag claa\_galla lclgaallca lccaallggg agagacaagl lccacalgcc 34020  
 5 aaggagaaac tgactgctcc cacaanaaag tatagaatta cagaaaatga tatacasatt 34080  
 ccattagatg atgcacaaa ccaactttaat caaaaactat ctcaactgca gacatatatg 34140  
 atacaatttg atcagtata taaagatagt tatgatttac atgattttaa aaagctatt 34200  
 10 gctaataatta ttga\_gaaa cattgacaaa taaaaaagtc ttgatgagca ctatcatatc 34260  
 cgtgtaaatt tagtaaaaac aatccatgat ctacatttgt ttattgaaa tactgatttt 34320  
 aacaaaagtg gaagtagtac tgcactctgg atcaaaaatg tggatactaa gtaccaaaac 34380  
 agaatccaga tacaagaaaa actgcagcag ctaagagac acatacagaa tatagacatc 34440  
 15 cagcacclag clggaagll azaacaacac a\_lgaggcla llga\_gllag aglgcLLLLa 34500  
 gatcaattqg qaactacaa ttcaattgaa aqaataaatg acattcttga gcatgtcaaa 34560  
 cactttgtta taaactta tggggatttt gaagtagctg agaaaatcaa tgcottcaga 34620  
 gccaaaagtc atgagttaa cgagaggtat gaagtagacc aacaaatcca ggttttaag 34680  
 20 gataaattag tagagttggc ccaccactac aagttgaagg agactattca gaagctaagc 34740  
 aatgtctac aacaagttaa gataaaagat tactttgaga aattggttgg atttattgat 34800  
 gatgctgta agaagctta tgaattctt ttaaaacat tcattgaaga tcttaacaaa 34860  
 25 lcccllgaca lgllgaaaa gaaallaaag lcalllgall accaccagll lglagalgaa 34920  
 accaatgaca aaatccgtga ggtgactcag agactcaatg gtgaaattca ggcctcggaa 34980  
 ctaccacaaa aagctgaagc attaaaactg ttttagagq aaaccaagcc cacagttgca 35040  
 ggtatctgg aaagcctaca ggacacaaa ataaccttae tcactcaattg gtacaggag 35100  
 30 gctttaagtt cagcatctt ggctcacatg aaggccaaat tccagagagc cttagaagat 35160  
 acacgagacc gaatgtatca aatggacatt cagcaggaac tcaacgata ccgtctctg 35220  
 ctaggccagq ttaaaagcac actgtcacc tacatttctg attggtggac tctgctgct 35280  
 aagaacotta ctgactttgc agagcaatat tctatcaag attgggctaa acgtatgaaa 35340  
 35 gcattgtag agcaagggt cactgttct gaaatcaaga ccatccttgg gaccatgct 35400  
 gcotttgaag tcaacttca ggccttcag aaagctacct tccagacacc tgattttata 35460  
 gcccctaa cagattttag gattccatca gtcagatae acttcaaga cttaaaaaat 35520  
 ataaaaatcc catccaggct tccacacca gaatttacca tcttaaacac ctccacatt 35580  
 40 cllccllla caal\_gacl lglagaaalg aaaglaaaga lcalcagac ca\_lgaaccag 35640  
 atgctgaaca gtgagctgca gtggccggt ccagatatat atctcagga tctgaagggt 35700  
 gaggacatto ctctagcag aatccctg ccagacttcc gtttaccaga aatgcaatt 35760  
 45 ccagaattca taatcccaac tctcaactt aatgattttc aagttctga cctcacata 35820  
 ccagaattcc agctcccca catctcacac acaattgaag tactacttt tggcaagcta 35880  
 tacagtattc tgaanaatca atctctctt tccacattag atgcaaatgc tgacataggg 35940  
 aatggaacce cctcagcaaa cgaagcgggt atcgcagctt ccactactgc caaaggagag 36000  
 50 tccaaattag aagtctcaa ttttgatttt caagcaaatg cacaactctc aaaccctaag 36060  
 a\_laalccgc lggccclgaa ggaglcaglg aagllclcca gcaaglaccl gagaaccgag 36120

55

EP 2 336 318 A1

catgggagtg aaatgctgt- ttttggaat gctattgagg gaaaatcaaa cacagtggca 36180  
 aglllacaca cagaaaaaaa cacacggag cllaglaalg gaglgallgl caagalaaac 36240  
 5 aatcagotta coctggatag caaactaaa tacttcaca aattgaacat ccccaactg 36300  
 cacttctcta gtcaggctca cctgcgcaac cagatcaaga cactgttgaa acctggccac 36360  
 atagcatgga ctctctctgg asaagggtca tggaaatggg cctgcccag atctcagat 36420  
 gagggaaac atgaatcaca aattagtttc accatagaag gaccctcac ttcctttgga 36480  
 10 cgtccaata agatcaatag caaacaccta agagtaaacc aaaacttggg ttatgaactt 36540  
 ggctccctca actttctaa acttgaatt caatcacaag tcgattccca gcatgtgggc 36600  
 cacagtgttc taacgctaa aggcattgca cgttttgag aaggaaggc agagttaact 36660  
 15 gggaggcag algc\_call\_ aaalggaag g\_lallgga clllgaaaa llclcllllc 36720  
 ttttcagccc agccatttca gacacggca tccacaaaac atgaaaggaa ttgaaaagt 36780  
 cgttttccat taaggttaac aggaagata gacttctga ataactatgc acgtttctg 36840  
 agtcccagtg cccagcaagc aagttggca gaaagtgca ggttcaatca gtataagac 36900  
 20 aacccaaaal lclcgclgg aaacaacgag aacallalgg aggccalgl aggaalaaal 36960  
 ggagaagca atctggattt cttaaacatt ccttaacaa ttcctgaaa ggtctacct 37020  
 tacacaataa tcacaactcc tcaactgaaa catctctctc tatgggaaa aacaggctg 37080  
 aaggaattct tgaaacgac acagcactca ttgatttac gtgtaaacgc tcagtataag 37140  
 25 aaaaaaaa acaggcattc cctcacaat cctttggctg tctttgtga gttatcagt 37200  
 cagagcatca aatccttga caggcatttt gaaaaaaca gaaacaatgc attagatttt 37260  
 gtcaccaaat cctataatga acaaaaatt aagtttgata agtacaagc tgaaaaact 37320  
 cagcagagc tcccaggac ctttcaatt cctggataca ctgttccagt tgcattggt 37380  
 30 gaagtgtctc cttcaccat agagatgtcg gcattcggct atgtgttcc aaaaagcagc 37440  
 agcatgcta gtttctccat cctagttct cagctcctg tgccttata cacattaatc 37500  
 ctgccatcat tagagctgc agtcttcat gtcctagaa atctcaagct ttctcttca 37560  
 35 gatttcaagg aattgtgtac ctaagccat atttttatc ctgccatgg caatattacc 37620  
 tatgatttct cctttaaac aagtgctac aactgaata ccaatgctga ac-ttttaac 37680  
 cagtcagata ttgtgctca tctctttct tcatcttcat ctgtcattga tgcactgcag 37740  
 tacaaattag agggcaccac aagattgaca agaaaaggg gattgaagtt agccacagct 37800  
 40 c\_glclclga gcaacaaal\_ lglggagggl aglcalaaca glaclglgag claaaccag 37860  
 aaaaatagtg aagtgtcag- ggcaacaacc acaaaagccc aaattccaat tt-gagaatg 37920  
 aatttcaagc aagaactta tggaaatacc aagtcaaac ctactgtctc ttctccatg 37980  
 45 gaatttaagt atgattcaa ttctccatg ctgtaactc cggc-aaag agcagttgac 38040  
 cacaagctta gcttggaaag cctcactct tacttttcca ttgagtcac taccaaagga 38100  
 gatgtcaagg gttcggttc- ttctcgggaa tattcaggaa ctattgctag tgaggcaac 38160  
 acttacttga attccaagag cacacggtct tcagtgaagc tgcaggccac ttccaaaatt 38220  
 50 gatgatctct ggaacctga agtaaaagaa aattttctg gagaagccac actccaacgc 38280  
 a\_alalccc lclggagca caglacgaaa aaccacllac agclagagg cc-cllllc 38340

55

EP 2 336 318 A1

accaacggag aacatacaag caaagccacc cgggaactct ctccatggca aatgtcagct 38400  
 cllgllcagc lccalgcagc lcagcccagl lclllccalg allllccclga ccllggccag 38460  
 5 gaagtggccc tgaaigctaa cactaagaac cagaagatca gatggaasaa tgaagtcogg 38520  
 atcattctg ggtctttcca gagccaggtc cagctttcca atgaccaaga aaaggcacac 38580  
 ctgacattg caggatcctc agaaggcac ctaaggttcc tcaaaaatat catcctacca 38640  
 gctatgaca agagcttatg ggatttccca aagctggatg taaccaccag catggtagg 38700  
 10 agacagcacc ttctgtttc aactgccttt ggttacacca aaaaccccaa tggctattca 38760  
 ttctccatcc ctgtaaaac tttgctgat aaattcatta ttctgggct gaaactaat 38820  
 gatctaaatt cagtctttc catgcctacg ttccatgtcc catttacaga tcttcagggt 38880  
 ccctcgtgca aacttgactc ccgagacata caaatctata agaagctgag aacttcaica 38940  
 15 ttgcccctca acctaccaac actcccagag gtaaaattcc ctgaagtga tgggtaaca 39000  
 aaatattctc aaccagaaga ctcttgatt ccttttttg agataacgtt gctgaaact 39060  
 cagttaactg tgtcccagt cagcttcca aaaagtgttt cagatggaat tggctgttg 39120  
 20 galclaaalg caglagccaa ccagalogca gaclllgagl lgcccaccal calcglcccl 39180  
 gagcagacca ttgagattcc ctccattaag ttctctgtac ctgctggaat tgcattct 39240  
 tctttcaag cactgactcc acgctttgag ctgagctctc cctgtataa tgcacttg 39300  
 agtgccagtt tgaanaacaa agcagattat gttgaaacag tctggattc cacatgagc 39360  
 25 tcaaccgtac agttcctaga atatgacct aatggtaaga aatatoctgc ctctctct 39420  
 agatactgta tattttcaac gagagttatg agtaanaat tatgtattta gtgtgagta 39480  
 gatgtacaat tactcaatg ccaaaaattt taagtaagas aagagatca tctatacct 39540  
 acacgtaaa accaaaactg agaaaacta ggtcattca agacaaacag cttaaagaa 39600  
 30 aatggatttt tctgtaatta ttttaggact aacaatgtct tttactatt tttttaaa 39660  
 taagtgtgag ctgtacattg catatttta acacaagtga aatactctc taqgataqaa 39720  
 ttctcccagt ttccacaatg aaaaaccaa cgtctactg ttatgaatct aataaaac 39780  
 35 aaaaactctc ctatacagt ttgggaccac acaaaatgc agatggtagg ttagectct 39840  
 agactaaagg aacattgca cccgtgact tcagtgcaga atatgaagaa gatggcaaat 39900  
 atgaaggact tccatagga gcttttatg aattgaaac ttataccttt tgaanaact 39960  
 40 ttgtgatttt ctccatctcc atacccttt cgtgatagct catctgtttt tctgcttca 40020  
 cgggaalggg aggaaaagcg cccclcaala lcaaaaagccc agcglccacc galclccalc 40080  
 tgcctacca gaaagacaag aaaggcatct ccacctcagc agcccccoca gccgtaggca 40140  
 ccgtgggcat gqatagga gaaqatgac acttttctaa atggaacttc tactacagcc 40200  
 45 ctccagtaac taccacctaa tgagtgcac gcccacaagc gcgagtggag aatggggca 40260  
 gatacattta atccaggacc aatattcag agattccca aactaggtga aagacaggcg 40320  
 gtaagcaact tcttctctga ggaatattc tctagaaagt attacaatga gtccttgatt 40380  
 cattttaatg tttagatgca ccatgacat cccatcagca ctattattta ttaattctg 40440  
 50 gcaaatccag gaagatgag gttatactc acatctaaa tcataggcaa gctcagccat 40500  
 aggcagggtat tattttcag agaggactgg ttctgtagt atttaaaac ttaaaattct 40560  
 tcccacaat agaattgcta gatgagatc acaaaattcc tctcatgtca tttacaagct 40620

55

EP 2 336 318 A1

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

cggccagggc caaatcaagg gtgacattac cagaggagaa gaccaaaacat ggtctataga 40680  
cgttactaa aagtctgta tgggcttggg gaatgogtac tgaigtggg atctggggc 40740  
tctgcagggt gggcccaac ttgccttttt tgcattttct tcttttcta tctgtaattt 40800  
cctgactctt ctctctctc ctctctctt tctctctctt actctctctt cagtctctag 40860  
tcttaggaag gcttaattt taagtgtcac aatgtaaatg acaaacagca agcgtttttg 40920  
ttaaactcct tctggggcat gtgataaaga gaaattaaca acagtagact taatttaacca 40980  
taaaacaaa acatqaactg acatatgaaa cataaatccc tttcaqtata tcaaaagattc 41040  
tctgatcttt atttttaact gctaataaag ttttagtga ctatattgtg taattgagat 41100  
aattgaaaac atgtatttt ttttttctc tctgtttagt cctctccaga taaaaactc 41160  
accatattca aaactgagtc gagggctcgg gaactctgat aggaactca gatcaaatg 41220  
aattgggaag aagaggcagc ttctggcttg caaacctctc tgaagagaaa cgtgcccag 41280  
ggcacagggg tctcttatga ttatgtcaac aagtaacct gggaaacac acggctcacc 41340  
cggagagaag tglcccaaaa gctgagaaga aatcctcaga acaatcctga ggggllal 41400  
caaggggcca ttaggcaaa tgaatgatac gaagttaggt tccagaaagc agccagtggc 41460  
accactgga cctaccaaga gtggaaggac aagcccaga atctgtaaca ggaactctg 41520  
actcaggaag gccaaagccag tttccaggga ccaaggata acgtgcttcg tggcttggtg 41580  
cgagttactc aagaattcca tatgaaagtc aagcatctga ttgactcact caatgatatt 41640  
cgaactctc ccagattcca gttctcgggg aaacctgga tataactag ggaggaaactt 41700  
tgcaactatg tcataaggga ggtagggacg gtaactctcc aggtatattc gaaagtccat 41760  
aatggttcag aaatactgtc tctctatttc caagacctag tgattacact tctcttag 41820  
tcaaggaaac ataaactaac agatgtaac tcgatgtata gggaaactgt gaaagattta 41880  
tcaaaaqaag cccaagagct atttaaqcc atcagctctc tcaagaccac agaggtccta 41940  
cgtaatcttc aggacctttt caaattcatt tccaactaa tagaagataa cattaacag 42000  
cgaagagaga tgaattttac ttatcttatt aattatctc aagatgagat caacacaatc 42060  
ccaglgall alalcccala tglulllaaa tgltgaaag aaaacctctg cc\_laalcll 42120  
cataagttca atgaatttat tcaaaaagag cctcaggaag ctctcaaga gtacagcag 42180  
atccatcaat acattatgct ccttcgtgaa gaatattttg atccaagtat agttggctgg 42240  
acagtgaat attaagaac tgaagaaag atagtcagtc tgatcaagaa ccgttagtt 42300  
gctcttaagg acttccattc tgaatatatt gtcagtgcct ctaactttac ttcccaactc 42360  
tcaagtcaag ttgagcaat tctgcacaga aatattcagg aatacttag cctccttacc 42420  
gatccagatg gaaaaggaa agagaagatt gcagagcttt ctgccactgc tcaggaaata 42480  
ataaaagcc aggcatttc gacgaagaaa ataatctctg attaccaaca gcagtttaga 42540  
tataaactgc aagaattttc agaccaactc tctgattact atgaaaaatt taatgctgaa 42600  
tcaaaaagat tgaatgacc gtccattcaa aactaccaca cttttctgat atacatcag 42660  
gagttactga aaaagctga atcaaccaca gcatgaacc cctacatgaa gctgctcca 42720  
ggagaactta ctatctcc ctaatttttt aaaagaaatc ttcattttat ctcttttcc 42780  
aattgaactt tccatagca cagaaaaaat tcaaacctgc tataatgata aaaccatata 42840

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

```

g-gagccagc cttccagtag gcagtagact ataagcagac gcacatatga acggacctg 42900
caccnaagct ggcaccaggc ctggcagggt cctggaactc agaaggatgg cactttttgc 42960
aagttaaaga aaatcagga- ctgagttatt ttgctaaact tgggggagga ggaacaaata 43020
aatggagtct ttattgtgta tcataccact gaatgtggct catttgtatt gaaagacagt 43080
gaaacgaggg cattgataaa atgttctggc acagcaaaa- ctctagaaca catagtgtga 43140
tttaagtaac agaa-aaaaa tggaaacgga gaaattatgg agggaaatat ttgcaaaaa 43200
tatttaaaaa gatcaagtaa ttgtgttttt a-aattaaat atttataa- taaaatattt 43260
a-aattaa- tattataa- taaatatttt a-aattaa- tattatae- taaatatttt 43320
a-aattaaag tattataa- taaatatttt a-aattaa- tattatae- taaatatttt 43380
a-aattaa- tattataa- taaatatttt a-aattaa- tattatae- taaatatttt 43440
a-aat. 43445

<210> 335
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 335
lclglaagac aggagaaaqa 20

<210> 336
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 336
atttctctt ctgtaaqaca 20

<210> 337
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 337
gatgccttac ttggacagac 20

<210> 338
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 338
agaaatagct ctcccaagga 20

<210> 339
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
  
```

EP 2 336 318 A1

5 <223> Antisense Oligonucleotide  
 <400> 339  
 g-cgcatcctt ctaacgtggg 20  
 <210> 340  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 340  
 tctccatac cttgcagttg 20  
 <210> 341  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 341  
 tggctcactgt ctaaccatct 20  
 <210> 342  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 342  
 caqttgaaat gcaqctaata 20  
 <210> 343  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 343  
 tgcagactag gagtgaaagt 20  
 <210> 344  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 344  
 aggaggatgt cctttattg 20  
 <210> 345  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 345  
 atcagagcac caaagggaaat 20  
 <210> 346  
 <211> 20

40  
 45  
 50  
 55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 346  
 ccagctcaac ctgagaattc 20  
  
 <210> 347  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 347  
 calgacllac ctggacatcg 20  
  
 <210> 348  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 20  
  
 <400> 348  
 cctcagcggg cacacacaca 20  
  
 <210> 349  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 349  
 gtcacatccg tgcctgggtc 20  
  
 <210> 350  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 35  
  
 <400> 350  
 cagtgcctct gggaccccac 20  
  
 <210> 351  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 351  
 agctgcagtg gccgatcagc 20  
  
 <210> 352  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 50  
  
 <400> 352  
  
 55

EP 2 336 318 A1

	gacctcccc gccacgtgga	20
5	<210> 353 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 353 tctgatcacc atacattaca	20
	<210> 354 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 354 atttcccaact gggtactctc	20
20	<210> 355 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 355 ggctgaagcc catgctgacc	20
	<210> 356 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 356 gttgacagc cattcttttg	20
35	<210> 357 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 357 cacllglllg acagcalle	20
	<210> 358 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 358 attttaaatt acactagata	20
50	<210> 359 <211> 20 <212> DNA <213> Artificial Sequence	
55		



EP 2 336 318 A1

	<220> <223> Antisense Oligonucleotide	
5	<400> 359 c-gttctcca cccatcatcag	20
	<210> 360 <211> 20 <212> DNA <213> Artificial Sequence	
10	<220> <223> Antisense Oligonucleotide	
	<400> 360 gagctcatac ctgtcccaga	20
15	<210> 361 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
20	<400> 361 tccaagggcc actcctatca	20
	<210> 362 <211> 20 <212> DNA <213> Artificial Sequence	
25	<220> <223> Antisense Oligonucleotide	
	<400> 362 ccagtatttc acgccaatcc	20
30	<210> 363 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
35	<400> 363 ggcaggaggc acctcgggca	20
	<210> 364 <211> 20 <212> DNA <213> Artificial Sequence	
40	<220> <223> Antisense Oligonucleotide	
	<400> 364 ttttaaatt agaccaacc	20
45	<210> 365 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
50	<400> 365 tgactgtttt aaaattagac	20
	<210> 366	

55

EP 2 336 318 A1

<211> 20  
<212> DNA  
<213> Artificial Sequence

5 <220>  
<223> Antisense Oligonucleotide

<400> 366  
cccagcaaac acaggtgaag 20

10 <210> 367  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

15 <400> 367  
gagtgtagtc ttgctagtc 20

<210> 368  
<211> 20  
<212> DNA  
<213> Artificial Sequence

20 <220>  
<223> Antisense Oligonucleotide

<400> 368  
ctatgcagag tgtggtattg 20

25 <210> 369  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

30 <400> 369  
agaagatgca accacatgta 20

<210> 370  
<211> 20  
<212> DNA  
<213> Artificial Sequence

35 <220>  
<223> Antisense Oligonucleotide

<400> 370  
acacggtatc ctatggagga 20

40 <210> 371  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

45 <400> 371  
tgggacttac catgcctttg 20

<210> 372  
<211> 20  
<212> DNA  
<213> Artificial Sequence

50 <220>  
<223> Antisense Oligonucleotide

55

EP 2 336 318 A1

	<400> 372	
	ggttttgctg cctacatcc	20
5	<210> 373	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
10	<400> 373	
	acaaggagtc cttctgcaga	20
	<210> 374	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Antisense Oligonucleotide	
	<400> 374	
	atgttcactg agacaggctg	20
20	<210> 375	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
25	<400> 375	
	gaaggtccat ggttcactg	20
	<210> 376	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Antisense Oligonucleotide	
	<400> 376	
	atagactgg aagcatcctg	20
35	<210> 377	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
40	<400> 377	
	gagattggag acgagcattc	20
	<210> 378	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Antisense Oligonucleotide	
	<400> 378	
	catgacctac ttgtaggaga	20
50	<210> 379	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
55		

EP 2 336 318 A1

	<220>		
	<223> Antisense Oligonucleotide		
5	<400> 379	tggatttggg tacacaagtc	20
	<210> 380		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
10	<220>		
	<223> Antisense Oligonucleotide		
	<400> 380	actcaatata tattcattga	20
15	<210> 381		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
20	<400> 381	caaggaagca caccatgtca	20
	<210> 382		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
25	<220>		
	<223> Antisense Oligonucleotide		
	<400> 382	atacttatttc ctggttaacca	20
30	<210> 383		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
35	<400> 383	ggtagccaga acaccagtgc	20
	<210> 384		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Antisense Oligonucleotide		
	<400> 384	actagaggtg gccagaacac	20
45	<210> 385		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
50	<400> 385	accacctgac atcacaggtc	20
55			

EP 2 336 318 A1

5 <210> 386  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 386  
 tactgtgacc tatgccagga 20

10 <210> 387  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 387  
 cgaagtgcta ctgttgacat 20

15 <210> 388  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 388  
 tccagacttg tctcagtcta 20

20 <210> 389  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 389  
 tctaagaggt agaçctaaag 20

25 <210> 390  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 390  
 ccagagatga gcaacttagg 20

30 <210> 391  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 391  
 ggccatgtaa attçctcatc 20

35 <210> 392  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

40  
 45  
 50  
 55

EP 2 336 318 A1

	<400> 392	
	aaagaaacLa LcclgLaLlc	20
5	<210> 393	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
10	<400> 393	
	t-cttaqtac ctgcaaatg	20
	<210> 394	
	<211> 20	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 394	
	cattagatac ctggacacc	20
20	<210> 395	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
25	<400> 395	
	gattcatgga actcagcgca	20
	<210> 396	
	<211> 20	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 396	
	c-ggagagca cctgcaatag	20
35	<210> 397	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
40	<400> 397	
	tgaaggtag agaaatcata	20
	<210> 398	
	<211> 20	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
	<400> 398	
	ggaaactcac ttgt-gacog	20
50	<210> 399	
	<211> 20	
	<212> DNA	

55

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
 <400> 399  
 aggtqcaaga tgttcctctg 20  
 <210> 400  
 <211> 20  
 <212> DNA  
 10 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 400  
 tgcacagagg tgcaagatg 20  
 15 <210> 401  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 20 <223> Antisense Oligonucleotide  
 <400> 401  
 cacaagagta aggagcagag 20  
 <210> 402  
 <211> 20  
 <212> DNA  
 25 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 402  
 gatggatggt gagaaattac 20  
 30 <210> 403  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 35 <223> Antisense Oligonucleotide  
 <400> 403  
 tagacaattg agactcagaa 20  
 <210> 404  
 <211> 20  
 <212> DNA  
 40 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 404  
 atgtgcacac aaggacatag 20  
 45 <210> 405  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 50 <223> Antisense Oligonucleotide  
 <400> 405  
 acatacaaat ggcaataggc 20  
 55

EP 2 336 318 A1

5  
 <210> 406  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 406  
 taggcaaagg acatgaatag 20

10  
 <210> 407  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 407  
 tttatgatagc tacagaataa 20

15  
 <210> 408  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 408  
 ctgagallac ccgcagaatc 20

20  
 <210> 409  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 409  
 catgtatgtc atataaaaca 20

25  
 <210> 410  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 410  
 ttccaatga cctgcattga 20

30  
 <210> 411  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 411  
 agggatggtc aatcggtag 20

35  
 <210> 412  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

55



EP 2 336 318 A1

<223> Antisense Oligonucleotide  
 <400> 412  
 ggctaataaa taggtagt- 20  
 5  
 <210> 413  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 10  
 <400> 413  
 tcctagagca ctatcaagta 20  
 <210> 414  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 414  
 ccLcclggLc clgcagLcaa 20  
 20  
 <210> 415  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 25  
 <400> 415  
 catttgacaca agtctttgt- 20  
 <210> 416  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 416  
 ctgacacacc atgttattat- 20  
 35  
 <210> 417  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 40  
 <400> 417  
 cttttcaga ctacataaga 20  
 <210> 418  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 418  
 tcacacttac ctcgatgagg 20  
 50  
 <210> 419  
 <211> 20  
 55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 419  
 aagaaaatgg catcaggttc 20  
  
 <210> 420  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 420  
 ccaagccaal ctgagaaaqa 20  
  
 <210> 421  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 421  
 aaatacacac ctgcctcatgc 20  
  
 <210> 422  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 422  
 ctccacaaat acacacctgc 20  
  
 <210> 423  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 423  
 agtgaagt.t.tggtctcatc 20  
  
 <210> 424  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 424  
 ttgctagctt caaagtggaa 20  
  
 <210> 425  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 425  
  
 40  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 425  
  
 45  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 425  
  
 50  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 425  
  
 55

EP 2 336 318 A1

	tcaagaataa gctccagatc	20
5	<210> 426 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 426 gcatacaagt cacatgaggt	20
	<210> 427 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 427 tacaagggtgt ttcttaagaa	20
20	<210> 428 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 428 atgcagccag gatgggccta	20
	<210> 429 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 429 ttaccatata ctgagaggtt	20
35	<210> 430 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 430 gcaaaagglag aggaaggla	20
	<210> 431 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 431 aaqgaccttc aqcaaaqgta	20
50	<210> 432 <211> 20 <212> DNA <213> Artificial Sequence	
55		

EP 2 336 318 A1

	<220>		
	<223> Antisense Oligonucleotide		
5	<400> 432	cataggagta catttatata	20
	<210> 433		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
10	<220>		
	<223> Antisense Oligonucleotide		
	<400> 433	attatgataa aatcaatttt	20
	<210> 434		
15	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
20	<400> 434	agaaatttca ctacatagac	20
	<210> 435		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
25	<220>		
	<223> Antisense Oligonucleotide		
	<400> 435	agcatatttt gatgagctga	20
	<210> 436		
30	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
35	<400> 436	gaaaggaagg aclagcatal	20
	<210> 437		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Antisense Oligonucleotide		
	<400> 437	cctctccaat ctgtagacc	20
	<210> 438		
45	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Antisense Oligonucleotide		
50	<400> 438	cggataact cagaccttg	20
	<210> 439		

55

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 439  
 agtcagaaaa caacctatct 20  
 <210> 440  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 440  
 cagcctgcat ctataagtca 20  
 <210> 441  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 441  
 aaagaattac cctccactga 20  
 <210> 442  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 442  
 tctttcaaac tggctaggca 20  
 <210> 443  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 443  
 gcctggcaaa attctgcaag 20  
 <210> 444  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 444  
 ctacctcaaa tcaatattgt 20  
 <210> 445  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 443  
 gcctggcaaa attctgcaag 20  
 <210> 444  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 444  
 ctacctcaaa tcaatattgt 20  
 <210> 445  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 444  
 ctacctcaaa tcaatattgt 20  
 <210> 445  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 55

EP 2 336 318 A1

	<400> 445 tgctttacct acctagctac	20
5	<210> 446 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 446 accttggtg tctcactcaa	20
	<210> 447 <211> 20 <212> DNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 447 atgcattccc tgactagcac	20
20	<210> 448 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 448 catctctgag cccctacca	20
	<210> 449 <211> 20 <212> DNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 449 gctgggcattg ctctctccc	20
35	<210> 450 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 450 gctttcgcag ctgggcattg	20
	<210> 451 <211> 20 <212> DNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 451 actcctttct atacctggc	20
50	<210> 452 <211> 20 <212> DNA <213> Artificial Sequence	
55		

EP 2 336 318 A1

	<220>	
	<223> Antisense Oligonucleotide	
5	<400> 452 a-tctgcctc tttagaaagt	20
	<210> 453 <211> 20 <212> DNA <213> Artificial Sequence	
10	<220>	
	<223> Antisense Oligonucleotide	
	<400> 453 ccaagcctct ttacgggc	20
15	<210> 454 <211> 20 <212> DNA <213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
20	<400> 454 cactcatgac cagactaaga	20
	<210> 455 <211> 20 <212> DNA <213> Artificial Sequence	
25	<220>	
	<223> Antisense Oligonucleotide	
	<400> 455 acctccaga agcctccat	20
30	<210> 456 <211> 20 <212> DNA <213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
35	<400> 456 tccatatgaa atctcctac	20
	<210> 457 <211> 20 <212> DNA <213> Artificial Sequence	
40	<220>	
	<223> Antisense Oligonucleotide	
	<400> 457 tatttaattt actgagaaac	20
45	<210> 458 <211> 20 <212> DNA <213> Artificial Sequence	
	<220>	
	<223> Antisense Oligonucleotide	
50	<400> 458 laalglttg ctgggaaga	20
55		

EP 2 336 318 A1

5 <210> 459  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 459  
 catctctaac ctgggtgtccc 20

10 <210> 460  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 460  
 ctgccatgct aggtggccat 20

15 <210> 461  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 461  
 agcaaattgg gatcctgtgct 20

20 <210> 462  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 462  
 tctggaggct cagaaacatg 20

25 <210> 463  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 463  
 tgaagacagc gagccaccta 20

30 <210> 464  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 464  
 aggattccca agactttgga 20

35 <210> 465  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

40  
 45  
 50  
 55



EP 2 336 318 A1

5 <400> 465  
cagcLcLaal claaagaca 20

<210> 466  
<211> 20  
<212> DNA  
<213> Artificial Sequence

10 <220>  
<223> Antisense Oligonucleotide

<400> 466  
caataactcac ctteLqcttg 20

<210> 467  
<211> 20  
<212> DNA  
<213> Artificial Sequence

15 <220>  
<223> Antisense Oligonucleotide

<400> 467  
aLctctctgt cctcatcttc 20

20 <210> 468  
<211> 20  
<212> DNA  
<213> Artificial Sequence

25 <220>  
<223> Antisense Oligonucleotide

<400> 468  
ccaactcccc ctttctttg 20

<210> 469  
<211> 20  
<212> DNA  
<213> Artificial Sequence

30 <220>  
<223> Antisense Oligonucleotide

<400> 469  
tctgggccag gaagacacga 20

35 <210> 470  
<211> 20  
<212> DNA  
<213> Artificial Sequence

40 <220>  
<223> Antisense Oligonucleotide

<400> 470  
tattgtgtgc tgggcactgc 20

<210> 471  
<211> 20  
<212> DNA  
<213> Artificial Sequence

45 <220>  
<223> Antisense Oligonucleotide

<400> 471  
tgcttgcac ctggacgag 20

50 <210> 472  
<211> 20  
<212> DNA

55

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
 <400> 472  
 ccttctttac cttaggtgac 20  
 <210> 473  
 <211> 20  
 <212> DNA  
 10  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 473  
 gctctctctg ccacctctga 20  
 15  
 <210> 474  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 20  
 <400> 474  
 aacttctaaa gccaacattc 20  
 <210> 475  
 <211> 20  
 <212> DNA  
 25  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 475  
 tgtgtcacia ctatggtaa 20  
 30  
 <210> 476  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 35  
 <400> 476  
 agacacatac cataatgcca 20  
 <210> 477  
 <211> 20  
 <212> DNA  
 40  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 477  
 tctcttcat ctgaaaatac 20  
 45  
 <210> 478  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 50  
 <400> 478  
 tgaggatgta attagcact 20  
 55

EP 2 336 318 A1

5  
 <210> 479  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 479  
 agctcattgc ctacaaaatg 20

10  
 <210> 480  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 480  
 gttctcatgt ttactaatgc 20

15  
 <210> 481  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 481  
 gaallgagac aactlgall 20

20  
 <210> 482  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 482  
 ccqccatcg ctgaaatgaa 20

25  
 <210> 483  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 483  
 catagctcac cttgcacat 20

30  
 <210> 484  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 484  
 cggcgcacc tttacctgag 20

35  
 <210> 485  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

40  
 <210> 485  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

45  
 <210> 485  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

50  
 <210> 485  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>

55

EP 2 336 318 A1

5 <223> Antisense Oligonucleotide  
 <400> 485  
 tctccagatc ctaacataaa 20  
 <210> 486  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 486  
 t-gaatgacc cttagattttc 20  
 <210> 487  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 487  
 aaaaaccall lllcLLaaag 20  
 <210> 488  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 488  
 caqctcacac ttattttaa 20  
 <210> 489  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 489  
 g-tcccaaac ctgtatagga 20  
 <210> 490  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 490  
 agctccatac tgaagtctc 20  
 <210> 491  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

35 <220>  
 <223> Antisense Oligonucleotide  
 <400> 491  
 caattcaata aaagctccat 20  
 <210> 492  
 <211> 20

40 <220>  
 <223> Antisense Oligonucleotide  
 <400> 491  
 caattcaata aaagctccat 20  
 <210> 492  
 <211> 20

45 <220>  
 <223> Antisense Oligonucleotide  
 <400> 491  
 caattcaata aaagctccat 20  
 <210> 492  
 <211> 20

50 <220>  
 <223> Antisense Oligonucleotide  
 <400> 491  
 caattcaata aaagctccat 20  
 <210> 492  
 <211> 20

55

EP 2 336 318 A1

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 5  
  
 <400> 492  
 gtttcaaaa ggtataaggc 20  
  
 <210> 493  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 493  
 LcccaLLcc clgaaagcag 20  
  
 <210> 494  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 15  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 494  
 tggtatttac ctgagggctg 20  
  
 <210> 495  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 495  
 aTaaataata gtgcTgatgg 20  
  
 <210> 496  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 496  
 cTatggctga gcttgcctaT 20  
  
 <210> 497  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 497  
 cTctctgaaa aataTaccct 20  
  
 <210> 498  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 498

EP 2 336 318 A1

5 t-gatgtatc tcacttagca 20  
 <210> 499  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 499  
 10 tagaaccatg tttggtcttc 20  
 <210> 500  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 15 <223> Antisense Oligonucleotide  
 <400> 500  
 tttctcttta tcacatgccc 20  
 <210> 501  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 20 <223> Antisense Oligonucleotide  
 <400> 501  
 tatagtacac taaaacttoa 20  
 <210> 502  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 25 <223> Antisense Oligonucleotide  
 <400> 502  
 ctggagaggc ctaaacagag 20  
 <210> 503  
 <211> 568  
 <212> DNA  
 <213> H. sapiens  
 <220>  
 <221> misc\_feature  
 <222> 44, 99, 156, 168  
 <223> n = A,T,C or G  
 <400> 503  
 ccaaaagatt gattgactgc ccattcaag ctacaagcaa tttntgatat acatcaogta 60  
 gttactgaae aagctgcaat caacacagtt catggaccnc taccatgag ct-gctccag 120  
 45 gagaacttct atcaatctct taatttttta aaaganatct tcattttatc ttctttttcca 180  
 atgaaacttt cacatagcac agaaaaaatt caaactgcoct atattgataa aaccatacag 240  
 tqagccagcc ttqcaqtaeq csgtagacta taagcagaag cacatgatgaa ctggacotgc 300  
 accaaaagctg gcaccagggc toggaaggtc tetgaaactca gaaggatggc at-tttttgca 360  
 50 agttaaagaa aatcaggatc tqagttatct tgctaaaact gggggaggag gaacaaaata 420  
 atggagtctt tattgtgta- cataccactg aatgtggctc atttgtanta aaagacagtg 480  
 55

EP 2 336 318 A1

aaacgagggc attgataaaa tgttctggca cagcaaaaac totagaacac atagtgtgat 340  
 ttaagtaaca gaataaaaaa ggaaaacgg 368

5

<210> 504  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10

<220>  
 <223> Antisense Oligonucleotide

<400> 504  
 acatthttatc aatgcctctg 20

15

<210> 505  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

20

<400> 505  
 gccagaacat tttatcaatg 20

<210> 506  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

25

<220>  
 <223> Antisense Oligonucleotide

<400> 506  
 agaggttttg ctgtgccaga 20

30

<210> 507  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

35

<400> 507  
 cagaggtlll tgctglgcca 20

<210> 508  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

40

<220>  
 <223> Antisense Oligonucleotide

<400> 508  
 tctagaqgtt ttgctgtacc 20

45

<210> 509  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

50

<400> 509  
 aatcacacta tgtcttctag 20

<210> 510

55

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 510  
 aaatcacact atgtgttcta 20  
 <210> 511  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 511  
 taaatcacac tatgtgttc 20  
 <210> 512  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 512  
 cttaaatacac actatgtgt 20  
 <210> 513  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 513  
 tattctgtta cttaaatacac 20  
 <210> 514  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 514  
 tggtagcctc agtcgcttc 20  
 <210> 515  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 515  
 agtctgcttc ggccttctg 20  
 <210> 516  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 50  
 <400> 516  
 gcgccagggc cgaagaggaa 20  
 55



EP 2 336 318 A1

<210> 517  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 5  
 <400> 517  
 caqgtatgag ctcaagctcg 20  
 <210> 518  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 10  
 <400> 518  
 catcctgaac atcaagaggg 20  
 <210> 519  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 15  
 <400> 519  
 eggcagtgtg atcgcttcaa 20  
 <210> 520  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 20  
 <400> 520  
 cacttgctct catcaaaggc 20  
 <210> 521  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 25  
 <400> 521  
 cacactggac gctaagagga 20  
 <210> 522  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 30  
 <400> 522  
 cgctgagcca cgcggtaaac 20  
 <210> 523  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 35  
 <400> 523  
 tgtccaaatt ctaccatggg 20  
 <210> 524  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 40  
 <400> 524  
 cagctgacct catcgagatt 20  
 <210> 525  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 45  
 <400> 525  
 gccaagtcc tgatggtgtc 20  
 <210> 526

55

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> H. sapiens  
 5 <400> 526 agctgcttct gatgggtgac 20  
 <210> 527  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 10 <400> 527 cggcatcacc attccggacc 20  
 <210> 528  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 15 <400> 528 cctactatcc gctgaccggg 20  
 <210> 529  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 20 <400> 529 cggccaacta agttgtgaca 20  
 <210> 530  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 25 <400> 530 agaacatggg attcccagac 20  
 <210> 531  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 30 <400> 531 ccccaacttc agtcctggg 20  
 <210> 532  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 35 <400> 532 cagagcttgg cctcctctggg 20  
 <210> 533  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 40 <400> 533 tggccgcttc agggaaacaa 20  
 <210> 534  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 45 <400> 534 cagctgagca gacaggcacc 20  
 <210> 535  
 <211> 20

55

EP 2 336 318 A1

	<212> DNA	
	<213> H. sapiens	
5	<400> 535 gggagagaca agtttcacac	20
	<210> 536	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 536 gtactgcac cctgcattcaa	20
	<210> 537	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 537 ctgaaggtgac tcaagactc	20
	<210> 538	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 538 tgcagagca atatctatc	20
	<210> 539	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 539 aaacattggt acaaccaagg	20
	<210> 540	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 540 ccgctggctc tgaaggagtc	20
	<210> 541	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 541 tctagtcagg ctgacctgg	20
	<210> 542	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 542 gggccacagt gttcaactg	20
	<210> 543	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 543 aatcaagtgt catcacactg	20
	<210> 544	
	<211> 20	
	<212> DNA	
50		
55		

EP 2 336 318 A1

	<213> H. sapiens	
	<400> 544	
5	gggtagtcat aacagtactg	20
	<210> 545	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 545	
10	agagcacacg gtcttcagtg	20
	<210> 546	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 546	
15	ttacagctag agggcctctt	20
	<210> 547	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 547	
20	caccgtgggc atggatatgg	20
	<210> 548	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 548	
25	gggaatctga tgacgaaact	20
	<210> 549	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 549	
30	tgtcaacaag taccactggg	20
	<210> 550	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 550	
35	acctgggata tacactaggg	20
	<210> 551	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 551	
40	ccaagtatag ttggctggac	20
	<210> 552	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 552	
	tacatgaagc ttgctccaag	20
	<210> 553	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
50		
55		

EP 2 336 318 A1

5	<p>&lt;400&gt; 553            atgtcagcct ggtctgtcca            &lt;210&gt; 554            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
10	<p>&lt;400&gt; 554            gcacctccgg aagtacacac            &lt;210&gt; 555            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
15	<p>&lt;400&gt; 555            ctgcagcttc atccgaaga            &lt;210&gt; 556            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
20	<p>&lt;400&gt; 556            tgagggcaac gccttgctga            &lt;210&gt; 557            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
25	<p>&lt;400&gt; 557            ccattccaga agggaaagcag            &lt;210&gt; 558            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
30	<p>&lt;400&gt; 558            cgaggaagg caatgtggca            &lt;210&gt; 559            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
35	<p>&lt;400&gt; 559            ccttgtcaac tctgacagc            &lt;210&gt; 560            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
40	<p>&lt;400&gt; 560            agcagccagt cctgtcagta            &lt;210&gt; 561            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
45	<p>&lt;400&gt; 561            agcatgtggc agaagccatc            &lt;210&gt; 562            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
50	<p>&lt;400&gt; 562            gagagcacc aatccacalc            &lt;210&gt; 563            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
55		

EP 2 336 318 A1

<210> 563  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 5  
 <400> 563  
 cctcagtgat gaaccagtca 20  
 <210> 564  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 10  
 <400> 564  
 gatagatgtg gtcacctacc 20  
 <210> 565  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 15  
 <400> 565  
 cctcagcaca gcagctgccg 20  
 <210> 566  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 20  
 <400> 566  
 gattctgctg gtcactggaa 20  
 <210> 567  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 25  
 <400> 567  
 caaagccatc actgatgatc 20  
 <210> 568  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 30  
 <400> 568  
 agaaagctgc catccaggct 20  
 <210> 569  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 35  
 <400> 569  
 caggaggctc ttcttcagac 20  
 <210> 570  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 40  
 <400> 570  
 gagtccttca caggcagata 20  
 <210> 571  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 45  
 <400> 571  
 tgccaatata ttgaactcag 20  
 <210> 572  
 <211> 20

55

EP 2 336 318 A1

	<212> DNA	
	<213> H. sapiens	
5	<400> 572 catcgagatt ggctgggaag	20
	<210> 573	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 573 ggagctggat tacagttgca	20
	<210> 574	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 574 caacatqcaq qctcaactcg	20
	<210> 575	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 575 acattacatt tggctctctac	20
	<210> 576	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 576 ctcagggcgt tactccaag	20
	<210> 577	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 577 cggacaccag attaqagctg	20
	<210> 578	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 578 gagctccaga gagaggacag	20
	<210> 579	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 579 atcggcagag tatcaccttg	20
	<210> 580	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 580 caaggtggtt atttcatac	20
	<210> 581	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
50	<400> 581 caaggtggtt atttcatac	20
	<210> 582	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
55	<400> 582 caaggtggtt atttcatac	20

EP 2 336 318 A1

5	<p>&lt;400&gt; 581            gactcactcctg ctacagctta            &lt;210&gt; 582            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
10	<p>&lt;400&gt; 582            gcaaatcctc cagagatcta            &lt;210&gt; 583            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
15	<p>&lt;400&gt; 583            cctctcctggg tgttctagac            &lt;210&gt; 584            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
20	<p>&lt;400&gt; 584            a_gaaggctg actcctgctc            &lt;210&gt; 585            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
25	<p>&lt;400&gt; 585            gggaccacag atgtctgctc            &lt;210&gt; 586            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
30	<p>&lt;400&gt; 586            cgggccggct caatggagag            &lt;210&gt; 587            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
35	<p>&lt;400&gt; 587            gctgcgttct gaatatcagg            &lt;210&gt; 588            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
40	<p>&lt;400&gt; 588            tgctgacatc ttaggcactg            &lt;210&gt; 589            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
45	<p>&lt;400&gt; 589            aagtgtagtc tctggtgct            &lt;210&gt; 590            &lt;211&gt; 20            &lt;212&gt; DNA            &lt;213&gt; H. sapiens</p>	20
50	<p>&lt;400&gt; 590            caaaattcag tctggatggg</p>	20
55		



EP 2 336 318 A1

5 <210> 591  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 591  
 gggaaactac ggctagaacc 20

10 <210> 592  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 592  
 ctgcctgtgg ctgctaaccct 20

15 <210> 593  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 593  
 ccatgacctat cgatgcacat 20

20 <210> 594  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 594  
 atgggaaact cgctctctgg 20

25 <210> 595  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 595  
 agtcatcatt tcgtgtctag 20

30 <210> 596  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 596  
 gaatacagcc aggacttggc 20

35 <210> 597  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 597  
 ggcgtggagc ttacaggagc 20

40 <210> 598  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 598  
 gagatgagag atgcccgttga 20

45 <210> 599  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 599  
 agtcttgatg agcactatca 20

50 <210> 600

55

EP 2 336 318 A1

	<211> 20	
	<212> DNA	
	<213> H. sapiens	
5	<400> 600	
	ctaaagtacca aatcagaatc	20
	<210> 601	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 601	
	ctccatqagt taatcgaag	20
	<210> 602	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 602	
	aggccacagt tgcagtgtat	20
	<210> 603	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 603	
	tctgattggg ggactcttgc	20
	<210> 604	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 604	
	caagtcagtc ttcaggctcc	20
	<210> 605	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 605	
	ccagattctc agatgagggc	20
	<210> 606	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 606	
	catctgtcat tgatgcactg	20
	<210> 607	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 607	
	aggagatgtc aagggttcgg	20
	<210> 608	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 608	
	ggaaactattg ctagcgaggc	20
	<210> 609	
	<211> 20	
	<212> DNA	
50		
55		

EP 2 336 318 A1

<213> H. sapiens  
 <400> 609  
 cctctccat ggcaaatgtc 20  
 5 <210> 610  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 610  
 caccgtgact tcagtcgaga 20  
 10 <210> 611  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 611  
 acLgagLLGa gggLccggca 20  
 15 <210> 612  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 612  
 cacatatgaa ctggacctgc 20  
 20 <210> 613  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 613  
 tetgaaactca gaacgatggc 20  
 25 <210> 614  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 614  
 ggtgcgaagc agactgagcc 20  
 30 <210> 615  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 615  
 tccaccggg acctgcgggg 20  
 35 <210> 616  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 616  
 caccgggacc tcggggctg 20  
 40 <210> 617  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 617  
 ctgagtgcc ttctcggttg 20  
 45 <210> 618  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 <400> 618  
 50  
 55

EP 2 336 318 A1

	c-cggttgct gccgctgagg	20
5	<210> 619 <211> 20 <212> DNA <213> H. sapiens	
	<400> 619 tcggttgctg ccgc-gagga	20
10	<210> 620 <211> 20 <212> DNA <213> H. sapiens	
	<400> 620 cggttgctgc cgctgaggag	20
15	<210> 621 <211> 20 <212> DNA <213> H. sapiens	
	<400> 621 gttgctgccg ctgaggagcc	20
20	<210> 622 <211> 20 <212> DNA <213> H. sapiens	
	<400> 622 c-gccgcTga ggagcccgc	20
25	<210> 623 <211> 20 <212> DNA <213> H. sapiens	
	<400> 623 accgcagctg gcgatggacc	20
30	<210> 624 <211> 20 <212> DNA <213> H. sapiens	
	<400> 624 cagctggcga tggaccgcc	20
35	<210> 625 <211> 20 <212> DNA <213> H. sapiens	
	<400> 625 gaacTLacla TcaTccTcLa	20
40	<210> 626 <211> 20 <212> DNA <213> H. sapiens	
	<400> 626 tccaattgaa ctttcacata	20
45	<210> 627 <211> 20 <212> DNA <213> H. sapiens	
	<400> 627 aaaallcaaa ctgcclala	20
50		
55		

EP 2 336 318 A1

<210> 628  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 5  
 <400> 628  
 cataaaacca tacagtgaac 20  
 <210> 629  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 10  
 <400> 629  
 aaaaaacccat acagtgagcc 20  
 <210> 630  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 15  
 <400> 630  
 aaccatacag tgagccagcc 20  
 <210> 631  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 20  
 <400> 631  
 accatacagt gagccagccc 20  
 <210> 632  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 25  
 <400> 632  
 ccatacagtg agccagcctc 20  
 <210> 633  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 30  
 <400> 633  
 gtagagccagc cttgcagtag 20  
 <210> 634  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 35  
 <400> 634  
 ccagccttgc agtaggcagc 20  
 <210> 635  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 40  
 <400> 635  
 taggcagtag actataagca 20  
 <210> 636  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 45  
 <400> 636  
 gcagtagact ataagcagaa 20  
 <210> 637  
 <211> 20

55

EP 2 336 318 A1

	<212> DNA	
	<213> H. sapiens	
5	<400> 637 tgaactggac ctgcacaaa	20
	<210> 638	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 638 caggacctgc accaaagctg	20
	<210> 639	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 639 ggacctgcac caaagctggc	20
	<210> 640	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 640 ctgcacaaa gctggcaca	20
	<210> 641	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 641 accaaagctg gcaccagggc	20
	<210> 642	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 642 ctcggaaaggt ctctqaactc	20
	<210> 643	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 643 aactcagaag gatggcattc	20
	<210> 644	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 644 ctcagaaggc tggcattttc	20
	<210> 645	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 645 atcaggatct gacttatttc	20
	<210> 646	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
50		
55		

EP 2 336 318 A1

	<400> 646 aggatctgag ttatcttgc	20
5	<210> 647 <211> 20 <212> DNA <213> H. sapiens	
	<400> 647 ctgagttatt ttgctaaact	20
10	<210> 648 <211> 20 <212> DNA <213> H. sapiens	
	<400> 648 atcttgctac acttggggga	20
15	<210> 649 <211> 20 <212> DNA <213> H. sapiens	
	<400> 649 laaacctggg ggagggagaa	20
20	<210> 650 <211> 20 <212> DNA <213> H. sapiens	
	<400> 650 ggaacaaata aatggagtc	20
25	<210> 651 <211> 20 <212> DNA <213> H. sapiens	
	<400> 651 gtttgtaact caagcagaag	20
30	<210> 652 <211> 20 <212> DNA <213> H. sapiens	
	<400> 652 ctgtaactca agcagaagc	20
35	<210> 653 <211> 20 <212> DNA <213> H. sapiens	
	<400> 653 gtaactcaag cagaaggtgc	20
40	<210> 654 <211> 20 <212> DNA <213> H. sapiens	
	<400> 654 aactcaagca gaaggtgcca	20
45	<210> 655 <211> 20 <212> DNA <213> H. sapiens	
	<400> 655 ctcaagcaga aggtgcgaag	20
50		
55		

EP 2 336 318 A1

5	<210> 656 <211> 20 <212> DNA <213> H. sapiens  <400> 656 caagcagaag gtgcgaagca	20
10	<210> 657 <211> 20 <212> DNA <213> H. sapiens  <400> 657 agcagaaggt gogaagcaga	20
15	<210> 658 <211> 20 <212> DNA <213> H. sapiens  <400> 658 cagaaggtgc gaagcagact	20
20	<210> 659 <211> 20 <212> DNA <213> H. sapiens  <400> 659 caaggtgcga agcagactga	20
25	<210> 660 <211> 20 <212> DNA <213> H. sapiens  <400> 660 aggtgogaag cagactgagg	20
30	<210> 661 <211> 20 <212> DNA <213> H. sapiens  <400> 661 gtgcgaagca gactgaggct	20
35	<210> 662 <211> 20 <212> DNA <213> H. sapiens  <400> 662 gcgaagcaga ctgaggctac	20
40	<210> 663 <211> 20 <212> DNA <213> H. sapiens  <400> 663 gaagcagact gagcctacca	20
45	<210> 664 <211> 20 <212> DNA <213> H. sapiens  <400> 664 agcagactga ggctaccatg	20
50	<210> 665	
55		



EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> H. sapiens  
 5 <400> 665  
 cagactgagg ctaccatgac 20  
 <210> 666  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 10 <400> 666  
 gactgaggct accatgacat 20  
 <210> 667  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 15 <400> 667  
 ctgaqgctac catcacattc 20  
 <210> 668  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 20 <400> 668  
 gaggctacca tgacattcaa 20  
 <210> 669  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 25 <400> 669  
 cgctaaccatg acattcaaat 20  
 <210> 670  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 30 <400> 670  
 ctaccatgac attcaaatat 20  
 <210> 671  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 35 <400> 671  
 cctgaagctg catctggctg 20  
 <210> 672  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 40 <400> 672  
 tgaagctgca tctggctggc 20  
 <210> 673  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 45 <400> 673  
 aagctgcatg tggctggtaa 20  
 <210> 674  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 50  
 55

EP 2 336 318 A1

	<400> 674 gclgcaglg gclgglacc	20
5	<210> 675 <211> 20 <212> DNA <213> H. sapiens	
	<400> 675 tgcatgtggc tggtaaccta	20
10	<210> 676 <211> 20 <212> DNA <213> H. sapiens	
	<400> 676 calglggclg glaaccclaaa	20
15	<210> 677 <211> 20 <212> DNA <213> H. sapiens	
	<400> 677 tgtggctggc aacctaaaag	20
20	<210> 678 <211> 20 <212> DNA <213> H. sapiens	
	<400> 678 tggctggtac cctaaaagga	20
25	<210> 679 <211> 20 <212> DNA <213> H. sapiens	
	<400> 679 gctggtaacc taaaaggagc	20
30	<210> 680 <211> 20 <212> DNA <213> H. sapiens	
	<400> 680 tggtaaccta aaaggagcc	20
35	<210> 681 <211> 20 <212> DNA <213> H. sapiens	
	<400> 681 gtaacctaaa aggagcctac	20
40	<210> 682 <211> 20 <212> DNA <213> H. sapiens	
	<400> 682 aacctaaaag gagcctacca	20
45	<210> 683 <211> 20 <212> DNA <213> H. sapiens	
	<400> 683	
50		
55		

EP 2 336 318 A1

	cctaaaagga gctacccaaa	20
5	<210> 684 <211> 20 <212> DNA <213> H. sapiens	
	<400> 684 ggcgcgaagc agacggaggc	20
10	<210> 685 <211> 20 <212> DNA <213> H. sapiens	
	<400> 685 cactatgttc atgagggagg	20
15	<210> 686 <211> 20 <212> DNA <213> H. sapiens	
	<400> 686 ccatcatagg ttctgacgtc	20
20	<210> 687 <211> 20 <212> DNA <213> H. sapiens	
	<400> 687 gaagctgatt gactcactca	20
25	<210> 688 <211> 20 <212> DNA <213> H. sapiens	
	<400> 688 ttgtaactca agcagaagcc	20
30	<210> 689 <211> 20 <212> DNA <213> H. sapiens	
	<400> 689 gtaactcaag cagaaggccc	20
35	<210> 690 <211> 20 <212> DNA <213> H. sapiens	
	<400> 690 aacccaagca gaaagcgcga	20
40	<210> 691 <211> 20 <212> DNA <213> H. sapiens	
	<400> 691 ctcaagcagc aggcgcgaag	20
45	<210> 692 <211> 20 <212> DNA <213> H. sapiens	
	<400> 692 cagaaggcgc gaaagcagac	20
50		
55		

EP 2 336 318 A1

	<210> 693	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
5	<400> 693	
	caagqccqca agcaqactca	20
	<210> 694	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 694	
	agqccqqaq caqactqacg	20
	<210> 695	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 695	
	gcgcgaagca gactgagga	20
	<210> 696	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 696	
	caagcaqact gacqctacca	20
	<210> 697	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 697	
	cagaaggcgc gaacqagact	20
	<210> 698	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 698	
	tctttctcct gtcttacaga	20
	<210> 699	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 699	
	cccacgtag aagatgcgac	20
	<210> 700	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 700	
	aatatggtag acatgagcca	20
	<210> 701	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 701	
	cattagctgc atttcaactg	20
	<210> 702	
	<211> 20	
	<212> DNA	
50		
55		

EP 2 336 318 A1

	<213> H. sapiens	
	<400> 702	
5	actttcactc ctagtctgca	20
	<210> 703	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 703	
10	ccatgtccag gtaagtcattg	20
	<210> 704	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 704	
15	gcaccaggca cggatgtgac	20
	<210> 705	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
	<400> 705	
20	gtgggggtccc agaggcactg	20
	<210> 706	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 706	
	gctgatcggc cactgcagcc	20
	<210> 707	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 707	
	tccacgtggc tggcgaqgc	20
	<210> 708	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 708	
	tgtaattgat ggtgatcaga	20
	<210> 709	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 709	
	gagagtacc agtgggaaat	20
	<210> 710	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 710	
	agtcagcatg ggtccagcc	20
	<210> 711	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
50		
55		

EP 2 336 318 A1

	<400> 711 caaaagaatg actgtccaac	20
5	<210> 712 <211> 20 <212> DNA <213> H. sapiens	
	<400> 712 gaatgactgt ccaacaagtg	20
10	<210> 713 <211> 20 <212> DNA <213> H. sapiens	
	<400> 713 tatctactgt aatttaaaat	20
15	<210> 714 <211> 20 <212> DNA <213> H. sapiens	
	<400> 714 cagataaggg lggagaacag	20
20	<210> 715 <211> 20 <212> DNA <213> H. sapiens	
	<400> 715 lclgggacag glalgaactc	20
25	<210> 716 <211> 20 <212> DNA <213> H. sapiens	
	<400> 716 tgatagcagt ggcccttgaa	20
30	<210> 717 <211> 20 <212> DNA <213> H. sapiens	
	<400> 717 ggallggcgl gaaaactcg	20
35	<210> 718 <211> 20 <212> DNA <213> H. sapiens	
	<400> 718 tgcccagggt tctcctgcc	20
40	<210> 719 <211> 20 <212> DNA <213> H. sapiens	
	<400> 719 gcactagcaa gaccacactc	20
45	<210> 720 <211> 20 <212> DNA <213> H. sapiens	
	<400> 720 caagaccaca ctctgcatag	20
50		
55		

EP 2 336 318 A1

5 <210> 721  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

<400> 721  
 tcctccatag gataccgtg 20  
 <210> 722  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

10 <400> 722  
 ggatgtaggg cagcaaaacc 20

<210> 723  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

15 <400> 723  
 totgcacaag gactccttg 20

<210> 724  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

20 <400> 724  
 cagcctgtct cagtgaaca 20

<210> 725  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

25 <400> 725  
 caggatgctt ccagtctaac 20

30 <210> 726  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

<400> 726  
 aaatgctcgt ctccaatctc 20

35 <210> 727  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

<400> 727  
 aacttggtga tccaaatcca 20

40 <210> 728  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

<400> 728  
 tgacatgggtg tgctcccttg 20  
 <210> 729  
 <211> 20  
 <212> DNA  
 <213> H. sapiens

45 <400> 729  
 acactgggtg tctggetacc 20

50 <210> 730  
 <211> 20

55

EP 2 336 318 A1

	<212> DNA	
	<213> H. sapiens	
5	<400> 730 g-gttcttggc taccctctagc	20
	<210> 731	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 731 tcctggcata ggtcacagta	20
	<210> 732	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 732 atgtcaacag taqcacctcc	20
	<210> 733	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 733 tagactcaga caagctctgga	20
	<210> 734	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 734 cctaagttgc tcctctctcg	20
	<210> 735	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 735 tgcgctgagt tccatgaaac	20
	<210> 736	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 736 ctattgcagg tgctctccag	20
	<210> 737	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 737 caqaqgaaca tottgcacct	20
	<210> 738	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 738 cctgtctcct tactcttctg	20
	<210> 739	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
50		
55		



EP 2 336 318 A1

	<400> 739 g_aaLLLLcLc accaLccalc	20
5	<210> 740 <211> 20 <212> DNA <213> H. sapiens	
	<400> 740 t-tctgagtct caat-gtcta	20
10	<210> 741 <211> 20 <212> DNA <213> H. sapiens	
	<400> 741 c_aatgLccLL gLgLGcaca_	20
15	<210> 742 <211> 20 <212> DNA <213> H. sapiens	
	<400> 742 gcctattgcc atttgtatg-	20
20	<210> 743 <211> 20 <212> DNA <213> H. sapiens	
	<400> 743 cattcatgt cott-gccta	20
25	<210> 744 <211> 20 <212> DNA <213> H. sapiens	
	<400> 744 gattctgagg gtaa-ctcag	20
30	<210> 745 <211> 20 <212> DNA <213> H. sapiens	
	<400> 745 tcaatgcagg tcat-ggaaa	20
35	<210> 746 <211> 20 <212> DNA <213> H. sapiens	
	<400> 746 c-accagatt gaccatccc-	20
40	<210> 747 <211> 20 <212> DNA <213> H. sapiens	
	<400> 747 tacttgatag tgctctagga	20
45	<210> 748 <211> 20 <212> DNA <213> H. sapiens	
	<400> 748	
50		
55		

EP 2 336 318 A1

	ttgactgcag gaccaggagg	20
5	<210> 749 <211> 20 <212> DNA <213> H. sapiens	
	<400> 749 aacaacact tgtcaaatg	20
10	<210> 750 <211> 20 <212> DNA <213> H. sapiens	
	<400> 750 aatgagacca aactccact	20
15	<210> 751 <211> 20 <212> DNA <213> H. sapiens	
	<400> 751 tccactttg aagctagcaa	20
20	<210> 752 <211> 20 <212> DNA <213> H. sapiens	
	<400> 752 gatctggagc ttattcttga	20
25	<210> 753 <211> 20 <212> DNA <213> H. sapiens	
	<400> 753 acctcatgtg acttgatgc	20
30	<210> 754 <211> 20 <212> DNA <213> H. sapiens	
	<400> 754 tcttaagas acaccttga	20
35	<210> 755 <211> 20 <212> DNA <213> H. sapiens	
	<400> 755 taggccalc ctggctgca	20
40	<210> 756 <211> 20 <212> DNA <213> H. sapiens	
	<400> 756 aaactctcag gatacgtaa	20
45	<210> 757 <211> 20 <212> DNA <213> H. sapiens	
	<400> 757 atacctcct ctaccttgc	20
50	<210> 758	
55		

EP 2 336 318 A1

	<211> 20	
	<212> DNA	
	<213> H. sapiens	
5	<400> 758 tacctttgct gaaggtcctc	20
	<210> 759	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 759 a-ctatctag tqaatttc	20
	<210> 760	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 760 tcagctcatc aaaatgct	20
	<210> 761	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 761 a-atgctagt ccttccttc	20
	<210> 762	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 762 caaaggtctg agttatccag	20
	<210> 763	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 763 tgacttatag atgcaggctg	20
	<210> 764	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 764 tcagtgagg gtaattcttc	20
	<210> 765	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 765 tgcctagcca gttgaaaga	20
	<210> 766	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 766 cctgcagaat ttgcccagcc	20
	<210> 767	
	<211> 20	
	<212> DNA	
50		
55		

EP 2 336 318 A1

	<213> H. sapiens	
5	<400> 767 gtagctaggt aggtaaagca	20
	<210> 768 <211> 20 <212> DNA <213> H. sapiens	
10	<400> 768 ttgagtgaga cacacaaggc	20
	<210> 769 <211> 20 <212> DNA <213> H. sapiens	
15	<400> 769 gtgctagtca gggaatgca	20
	<210> 770 <211> 20 <212> DNA <213> H. sapiens	
20	<400> 770 ggggagagag catgcccagc	20
	<210> 771 <211> 20 <212> DNA <213> H. sapiens	
25	<400> 771 gcatgcccag ctgcgaaagc	20
	<210> 772 <211> 20 <212> DNA <213> H. sapiens	
30	<400> 772 agccaggat agaaaggagc	20
	<210> 773 <211> 20 <212> DNA <213> H. sapiens	
35	<400> 773 aactttctaa gaggcagaa	20
	<210> 774 <211> 20 <212> DNA <213> H. sapiens	
40	<400> 774 tcttagtctg gtcaagagtg	20
	<210> 775 <211> 20 <212> DNA <213> H. sapiens	
45	<400> 775 agtaggagat ttcaatgaa	20
	<210> 776 <211> 20 <212> DNA <213> H. sapiens	
50		
55		

EP 2 336 318 A1

	<400> 776 tcttcaccag caacacatta	20
5	<210> 777 <211> 20 <212> DNA <213> H. sapiens	
10	<400> 777 atggccacct agcatggcac	20
	<210> 778 <211> 20 <212> DNA <213> H. sapiens	
15	<400> 778 catgtttctg agccaccaga	20
	<210> 779 <211> 20 <212> DNA <213> H. sapiens	
20	<400> 779 taggtggctc cctgacctca	20
	<210> 780 <211> 20 <212> DNA <213> H. sapiens	
25	<400> 780 lccaaaglcl lggaalcc	20
	<210> 781 <211> 20 <212> DNA <213> H. sapiens	
30	<400> 781 acaaagaaag gggcagttcg	20
	<210> 782 <211> 20 <212> DNA <213> H. sapiens	
35	<400> 782 lcgtglcllc clggcccaga	20
	<210> 783 <211> 20 <212> DNA <213> H. sapiens	
40	<400> 783 gcagtgccca gcacacaata	20
	<210> 784 <211> 20 <212> DNA <213> H. sapiens	
45	<400> 784 actcgtccag gtgogaagca	20
	<210> 785 <211> 20 <212> DNA <213> H. sapiens	
50	<400> 785 gccacclaag glaaagaag	20
55		

EP 2 336 318 A1

<210> 786  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 5  
 <400> 786  
 atcaqaqtgg cagaqagaac 20  
 <210> 787  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 10  
 <400> 787  
 ttaccatag ttgtgacaca 20  
 <210> 788  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 15  
 <400> 788  
 cattttgtag gcaatgagcc 20  
 <210> 789  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 20  
 <400> 789  
 gcattagtaa acatgagaac 20  
 <210> 790  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 25  
 <400> 790  
 tcatttcag cgatggcccg 20  
 <210> 791  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 30  
 <400> 791  
 gaaaatctag tgcattcaa 20  
 <210> 792  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 35  
 <400> 792  
 tcctatacag ttttgggaac 20  
 <210> 793  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 40  
 <400> 793  
 aaggacttca gtatggagcc 20  
 <210> 794  
 <211> 20  
 <212> DNA  
 <213> H. sapiens  
 45  
 <400> 794  
 atggagcttt tattgaattg 20  
 <210> 795  
 <211> 20

55

EP 2 336 318 A1

	<212> DNA	
	<213> H. sapiens	
5	<400> 795 ccatcagcac tattatttat	20
	<210> 796	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
10	<400> 796 ataggcaagc tcaçccatag	20
	<210> 797	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
15	<400> 797 tçctaqatqa qatacatcaa	20
	<210> 798	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
20	<400> 798 gaagaccaaa catggttcta	20
	<210> 799	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
25	<400> 799 ctctgttttag tçctctccag	20
	<210> 800	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
30	<400> 800 cattqataaa atgtçctgçc	20
	<210> 801	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
35	<400> 801 tctggcacag caaacctctç	20
	<210> 802	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
40	<400> 802 tggcacagca aaacçtctag	20
	<210> 803	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
45	<400> 803 tagaacacat agtççgattç	20
	<210> 804	
	<211> 20	
	<212> DNA	
	<213> H. sapiens	
50		
55		

EP 2 336 318 A1

5 <400> 804  
aacacalagl glgaLLaag 20

<210> 805  
<211> 20  
<212> DNA

<213> Artificial Sequence

10 <220>  
<223> Antisense Oligonucleotide

<400> 805  
c-tttccgttg gaccocctggg 20

<210> 806  
<211> 20  
<212> DNA

15 <213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

20 <400> 806  
Lcccgcclgl gacaLgcalL 20

<210> 807  
<211> 20  
<212> DNA

25 <213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 807  
t-ttaactcg cgcatttac 20

30 <210> 808  
<211> 432  
<212> DNA  
<213> O. cuniculus

<400> 808  
gatcttacct totccaagca aaatgcattg ctacgtgctg agtatcaggc tgattacaag 60

35 tcaactgaggt tottcaacct gctttctggg t-gttgaata cccatggtct tgaattaat 120  
gctgacatct tgggcactga caaaatgaat actgctgctc acaaggcaac tctaagaatt 180  
ggccaaaatg gagtatctac cagtgcaca accagcttga ggtacagtc cctgatgctg 240  
gagaatgagc tgaacgcaga gcttgcctt totggggcat ctatgaaatt agcaacaaat 300

40 ggccgcllc aggaacacaa lgcaaaellc agcclagalg gaaagclac cclcacagag 360  
t-atcoctgg gaagcgctta ccaggccatg atctgggtg ctgacagcaa gaacatttc 420  
aacttcaaga tc 432

45 <210> 809  
<211> 660  
<212> DNA  
<213> O. cuniculus

<400> 809  
c-gggaaaac toccacagca agttaatgat tatctgagta cattcaattg ggagagacaa 60

50 g-ttccagtg ccaaggagaa actaactact t-cacaaaaa attataaat tacagagaat 120  
gatatacaaa ctgcattgga taatgcacaa atcaacttae atgaaaaac gtctcaactt 180

55



EP 2 336 318 A1

5 cagacatag tgatataatt tgatcagtat actaaagata attttgatct acatgatatt 240  
 aaaatagcta tagcagta= tatagatcaa acatggaaa aattaaaa= tottgatgaa 300  
 10 ogttatcata tccgngcaca ttttaattaaa tcaatccata atttatatatt gtttattgaa 360  
 gctattgatt ttaacaaaa= tggaagtagt actgcatott ggattcaaaa tgggatacc 420  
 aagtatcaag tcagaactcg gatacaagaa atattgcaac agtttaagac acagattcag 480  
 aatacaaaaca toccataccc ggctgaaaaa cggaaacaac agatggaggc tattgatgtc 540  
 agagtqcttt tagacaaat= gagaaactaca atccatttc gtataataaa gacattatt 600  
 gaacatttca aatactttgt tataaatatt atgaaaatt ttgaagtaat tcacaaaaac 660

15 <210> 810  
 <211> 543  
 <212> DNA  
 <213> O. cuniculus

20 <220>  
 <221> misc\_feature  
 <222> (45)  
 <223> n = a, c, g, or t

<220>  
 <221> misc\_feature  
 <222> (118)  
 <223> n = a, c, g, or t

25 <220>  
 <221> misc\_feature  
 <222> (148)  
 <223> n = a, c, g, or t

30 <220>  
 <221> misc feature  
 <222> (173)  
 <223> n = a, c, g, or t

<220>  
 <221> misc\_feature  
 <222> (180)  
 <223> n = a, c, g, or t

35 <400> 810  
 cagaacatcg gagacaaccg attggatttt ccactaaat cttanaatga agcaaaaatt 60  
 aagtttgata agtacaagg= tgaaaaatcg ccaaacaggc tcccaggac ctctcagct 120  
 40 cctggataca ttattccaat ttccaatntt gaagtatctc cactcacaat agnagacgtn 180  
 agcallcagl calgagalec caaaalcaal aagcacccc aalgtaacca tcclga\_lc 240  
 aagcttctat gtgccttca= atacattggc tetgccaac ctagagctgc cagtcttcca 300  
 tqtccccagg aatccactca aqgtctctct tccaqatttc aagcaattca aaaccattaa 360  
 45 caatattttt attocagca tgggcaccat tacctatgac ttttccttca aacaaagat 420  
 cattacactg aataccaatg ctggacttta taaccaatca gacattgttg cccatctct 480  
 tcttctctct tcatctgtca ttgatgcact acagtacaaa tttagaggga cgtcaagtt 540  
 tga 543

50 <210> 811  
 <211> 19  
 <212> DNA

55

EP 2 336 318 A1

<213> Artificial Sequence  
 <220>  
 <223> Primer  
 5  
 <400> 811  
 aaqcaacccc aatqtcacc 19  
 <210> 812  
 <211> 20  
 <212> DNA  
 10  
 <213> Artificial Sequence  
 <220>  
 <223> Primer  
 <400> 812  
 gggatggcag agccaatgta 20  
 15  
 <210> 813  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Probe  
 20  
 <400> 813  
 tctctggattc aagcttctat gtgccttca 29  
 <210> 814  
 <211> 20  
 <212> DNA  
 25  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 814  
 tgcttggaga aggtaagatc 20  
 <210> 815  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 815  
 gcgttgtctc cgatgttctg 20  
 <210> 816  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 45  
 <400> 816  
 taatcattaa ctctctgtgg 20  
 <210> 817  
 <211> 20  
 <212> DNA  
 50  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 55

EP 2 336 318 A1

5 <400> 817  
tcagcagcta gcaatgcatc 20

<210> 818  
<211> 20  
<212> DNA  
<213> Artificial Sequence

10 <220>  
<223> Antisense Oligonucleotide

<400> 818  
gcctgatact cagcagctag 20

15 <210> 819  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

20 <400> 819  
caattgaatg tactcagata 20

<210> 820  
<211> 20  
<212> DNA  
<213> Artificial Sequence

25 <220>  
<223> Antisense Oligonucleotide

<400> 820  
acctcagtga cttgtaatca 20

30 <210> 821  
<211> 20  
<212> DNA  
<213> Artificial Sequence

35 <220>  
<223> Antisense Oligonucleotide

<400> 821  
cactggaaac ttgtctctcc 20

40 <210> 822  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

45 <400> 822  
agtagttagt ttctccttgg 20

<210> 823  
<211> 20  
<212> DNA  
<213> Artificial Sequence

50 <220>  
<223> Antisense Oligonucleotide

55

EP 2 336 318 A1

	<400> 823 tcagtgcccc agatgtcagc	20
5	<210> 824 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 824 atgqaataa tgtaaccaag	20
15	<210> 825 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
20	<400> 825 tggcattat ccaatgcaag	20
25	<210> 826 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
30	<400> 826 gtgccttgt gacagcagc	20
35	<210> 827 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 827 atgtgagtg gagaacttc	20
45	<210> 828 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
50	<400> 828 catatgtctg aagtggagac	20
55	<210> 829 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
	<400> 829	

EP 2 336 318 A1

	g-agatactc cattttggcc	20
5	<210> 830 <211> 20 <212> DNA <213> Artificial Sequence  <220> <223> Antisense Oligonucleotide	
10	<400> 830 ggatcacatg actgaatgc	20
15	<210> 831 <211> 20 <212> DNA <213> Artificial Sequence  <220> <223> Antisense Oligonucleotide	
20	<400> 831 tcaagctggt tgttgactg	20
25	<210> 832 <211> 20 <212> DNA <213> Artificial Sequence  <220> <223> Antisense Oligonucleotide	
30	<400> 832 cgactgtacc tcaagctgc	20
35	<210> 833 <211> 20 <212> DNA <213> Artificial Sequence  <220> <223> Antisense Oligonucleotide	
40	<400> 833 gctcattcgc cagcaccgg	20
45	<210> 834 <211> 20 <212> DNA <213> Artificial Sequence  <220> <223> Antisense Oligonucleotide	
50	<400> 834 t-gatctata ataccagcta	20
55	<210> 835 <211> 20 <212> DNA <213> Artificial Sequence  <220> <223> Antisense Oligonucleotide	
	<400> 835 a-ggaagact ggcagctcta	20

EP 2 336 318 A1

5  
 <210> 836  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

10  
 <400> 836  
 ttgtgttcct tgaagcggcc 20  
 <210> 837  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

15  
 <400> 837  
 tgtgcacgga tatgataaag 20  
 <210> 838  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

20  
 <400> 838  
 gaccttgagt agatccctgg 20  
 <210> 839  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

25  
 <400> 839  
 gaaatctggc agagagacct 20  
 <210> 840  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

30  
 <400> 840  
 gtaagctttcc catctaagcc 20  
 <210> 841  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

35  
 <400> 841  
 gataacclcg tgaggglacc 20  
 <210> 841  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

40  
 <400> 841  
 gataacclcg tgaggglacc 20  
 <210> 841  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

45  
 <400> 841  
 gataacclcg tgaggglacc 20  
 <210> 841  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

50  
 <400> 841  
 gataacclcg tgaggglacc 20  
 <210> 841  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide

55

EP 2 336 318 A1

<210> 842  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 842  
 a-gttgcca tggc-ggaa- 20  
 10  
 <210> 843  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Antisense Oligonucleotide  
 15  
 <400> 843  
 aagatgcagt actactcca 20  
 <210> 844  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 844  
 gcacccagaa tcatggcctg 20  
 <210> 845  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 845  
 ctgatactt ggta-ccaca 20  
 <210> 846  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 846  
 cagtgtaatg atcgtgat- 20  
 <210> 847  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 847  
 taaagtccag cattggtat- 20  
 <210> 848  
 50  
 55

EP 2 336 318 A1

<211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 848  
 caacaatgtc tgatgggta 20  
 10  
 <210> 849  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 15  
 <400> 849  
 gaagaggaag aaaggatatg 20  
  
 <210> 850  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 850  
 lgacagatga agaggaagaa 20  
 25  
 <210> 851  
  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 851  
 t-gtactgta gtgcatcaat 20  
 35  
 <210> 852  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Antisense Oligonucleotide  
 40  
 <400> 852  
 gccacaalcl gllgllcag 20  
  
 <210> 853  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Antisense Oligonucleotide  
  
 <400> 853  
 acttgaqcgt gccctctaat 20  
 50  
 <210> 854  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 55





EP 2 336 318 A1

<400> 858  
ccttccctga aggttccctc 20

<210> 859  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 859  
ccttactgt gctgtggaca 20

<210> 860  
<211> 13938  
<212> DNA  
<213> H. sapiens

<400> 860  
ctggattgq qacacacttt ctggacactg ctggccagtc ccaaaatgga acataacgaa 80  
gtgggtcttt tacttctttt atttctgaaa tcagcagcac ctgagcaaag caatgttggtc 120  
caggatttgt accatgggtg tggacagagt tatcgaggca cgtactccac cactgtcaca 180  
ggaaggacct gccaaagcttg gtcatctatg acaccacata aacataatag gaccacagaa 240  
aactaoccaa atgctggctt gatcatgaac taactgcagga atccagatgc tctggcagct 300  
ccllalgl ll alacgaggga lcccgglglc agglgggagl aclgcaacct gacgcaa\_gc 360  
tcagaocgca aagggaactgc cgtcgcgcct cggactgtta ccccggttcc aagcctcag 420  
gctccttccg aacaagcaac gactgagcaa aggcctgggg tgcaggagtg ctaccatggt 480  
aatggacaga gttatcgagg cacatactcc accactgtca caggaagaac ctgccaaagct 540  
ttgtcatcta tgacaccaca ctgcataagt cggaccocaa aatactaccc aaatgctggc 600  
ttgatcatga actactgcag gaatccagat gctgtggcag ctcttatttg ttatccagg 660  
gatcccggtg tcaagtgagg gtactgcaac ctgacgcaat gctcagaagc agaagggact 720  
cccgltcggc clccgactg lccccggll ccaagcclag agglccllc cgaacaagca 780  
ccgactgagc aaaggcctgg ggtgcaggag tgcatacatg gtaatggaca gsgttatcga 840  
ggcacatact ccaaccctgt cacaggaaga acctgccaaag ctgtgtcctc tatgacacca 900  
cactcgcata gtcggaccac agaatactac ccaaatgctg gcttgcatac gaactacgac 960  
aggaatccag atgctgtgac agctccttat tgttatacga gggatcccgg tccagctgg 1020  
gagtaactgc acctgacgca atgctcagac gcagaaggga ctgcctcgc gccctccgact 1080  
gttaccocgg tccaagcct agaggctcct tcogaacaag caccgactga gcaaaaggcct 1140  
ggggtgcagg agtgcataca tggtaatgga cagagttatc gaggcacata ctccaccact 1200  
gtcacaggaa gaacctgcca agcttggtca totatgacac cacactcgca tagtggacc 1260  
ccagaatact acccaaatgc tggcttgatc atgaactact gcaggaatcc agatgctgtg 1320  
cgactcctt attgtatac gaggatccc ggtgtcaggt gggagtastg caacctcagc 1380  
caatgctcag acgcagaaag gactgcctgc ccgctccga ctgttacccc gctccaagc 1440  
ctagaggctc ctccgaaaca agcaccgact gagcaaaagg ctggggtgca gtagtgcac 1500  
catggttaat gacagagtta togaggcaca taactcacca ctgtcacagg aagaacctgc 1560  
caagcttggc catctatgac accacactcg catagtggga ccccagaata ctacccaat 1620  
gctggcttga toatgaacta ctgcaggaat ccagatgctg tggcagctcc ttattgttat 1680  
acgagggalc ccgg\_glcag glgggaglac lgcacclga cgcgaalglc agcgcagaa 1740  
gggactgccc tggcctccac gactgttacc ccggttccaa gccctagaggc tccctccgaa 1800  
caagcaccga ctgagcaaaag gctgggggtg caggagtgtc acctggttaa tggacagagt 1860  
tatcgaggca cactactccac cactgtcaca ggaagaacct gccaaagcttg gtcactatg 1920  
acaccacact cgcatagttg gacccagaa taactcccaa atgctggctt gatcatgaa 1980  
tactgcagga atccagatgc tgtggcagct ccttattggt atcagaggga tcccggtgtc 2040  
aggtgggagt actgcaacct gacgcaatgc tcagacgag aagggactgc cgtcgcgcct 2100  
ccgactglle ccccgllcc aagcclagag cclcllccg aacaagcacc gactlgagaa 2160  
agcctggggg tgcaggagtg ctaccatggt aatggacaga gttatcaggg cacataccc 2220  
accactgtca caggaagaac ctgcocagct tggcatcta tgacaccaca ctgcataagt 2280  
cggaccocag aatactaccc aeatgctggc ttgataatga actactgcag gaatccagat 2340  
cctgtgqcaq ctcttatttg ttatacagag gatcccgtg tcaggtggga gtaactgaa 2400  
ctgacgcaat gctcagagcc agaagggact gccgtcggc ctccgactgt taccocggt 2460  
ccaagcctag aggcctcttc cgaacaagca cggactgagc aaaggcctgg gg-gcaggag 2520  
tgctaccatg gtaatggaca gacttatcga ggcacatact ccaccactgt cacaggaaga 2580  
acctgccaaag ctgtgtcctc tatgacacca cactcgcata gtcggacccc agaatactac 2640  
ccaaatgctg gcttgatcat gaactactgc aggaatccag atgctgtggc agctccttat 2700  
tgttatacga gggaccccgg tgtcaggtgg gactactgca acctgacgca atgctcagac 2760  
ccagaaggga ctgcctcctc gctccagact ctaccccgg ttccaagcct acaggtcct 2820  
tcogaaccaag caccgactga gcaaaaggcct ggggtgcagg agtgcatacca tggtaatgga 2880  
cagagttatc gaggcacata ctccaccact gtcacaggaa gaacctgcca agcttggca 2940  
tctatgacac cacactcgcata tagtcggacc ccagaatact acccaaatgc tggcttcatc 3000  
atgaactact gcaggaatcc agatgctgtg ccagctcctt attgttatac gagggatccc 3060  
gglglcaggl gggaglacg caacclgag acgcagaaag gactgcccgc 3120  
gcgcctccga ctgttaccoc ggttccaagc ctgagggctc ctccgaaaca agcaccgact 3180

EP 2 336 318 A1

	gagcaaaaggc	ctgggggtgca	ggagtgctac	catggtaatg	gacagagtta	togaggcaca	3240
	tactocacca	ctgtocacagg	asgaacctgc	caagcttggg	catctatgac	accacaotcg	3300
	calaglcggg	ccccagaala	claccccaal	gclggcllga	lcalgaacfa	clgcaggaa	3360
	ccagatgctg	tggcagctcc	ttattgttat	acgagggato	ccgggtgcag	gtggggagtac	3420
5	tgcaacctga	cgcaatgctc	agaocgagaa	gggactgcog	tccgcoctcc	gactgttacc	3480
	ccgggtccaa	gocctagaggc	tccctccgaa	caagcaccga	ctgagcazag	gocctggggtg	3540
	caqgagtgct	accaatggtaa	tggacagagt	tatocgaggca	catactccac	cactgtocaca	3600
	ggaagaacct	gccaagcttg	gtcatctatg	acaccacact	cgcatagtccg	gaccccagaa	3660
	tactacccaa	atgcctggctc	gatcatgaac	tactgcagga	atccagatgc	tctggcagct	3720
	ccttatgtgt	atacagaggga	tcccgggtgc	agggtgggagt	actgcaacct	gacgcaatgc	3780
	tcagaocgag	aagggaactgc	cgtocogcct	ccgactgtta	ccccgggtcc	aagcctagag	3840
10	gctccttccg	aacaagcaacc	gactgagcaa	aggcctgggg	tgcaaggagt	ctaccatggt	3900
	aatggacaga	gttatcgagg	cacatactcc	accactgtca	caggaaagac	ctgccaagct	3960
	tqqtcatcta	tqacaccaca	ctcgcatagt	ccgaccccag	aatactaacc	aatatgctggc	4020
	t-gatcatga	actactgcag	gaatccagat	gctgtggcag	ctccctattg	ttataccagg	4080
	gatccocggg	toagggtggga	gtactgcaac	ctgacgcaat	gctcagacgc	agaagggaact	4140
	gocgtgcggc	ctccgactgc	tccccgggt	ccaagcctag	aggctccttc	cgaaacaagca	4200
	ccgactgagc	aaaggcoctgg	gggtgcaggag	tgctaccatg	gtaatggaca	gagttatcga	4260
15	ggcacaiacl	ccaccactgc	ccagggaaga	acclgccaag	cllglgclalc	la_lgacacca	4320
	caactcgata	gtccgacccc	agaatactac	ccaaatgctg	gcttgatcat	gaactactgc	4380
	agqaatccag	atgcctgtgoc	aqctccttat	tqttatacga	ggqatccccq	tqtcaqgtqq	4440
	gagtaactgca	acctgacgca	atgctcagac	gcagaaggga	ctgocgtcgc	gocctccgact	4500
	gttacocccg	ttccaagcct	agaggctcct	tcogaaacaag	caocgactga	gcaaaagcct	4560
	gggggtcagg	agtgtaccca	tggtaatgga	ccagagttatc	gaggcacata	ctccaccact	4620
	gtcacaggaa	gaacctgcca	agcttggtca	tctatgacac	cacactcga	tagtccgacc	4680
20	ccagaalaccl	accocaaalg	lggclllgalc	a_lgaacclac	gcaggaaacc	agalglclglg	4740
	gcagctcctt	attgtatatac	ggggatcccc	gggtgcagg	gggagtactg	caacctgacg	4800
	caatgctcag	acgcagaagg	gactgcogtc	ggcctccga	ctgttaccoc	ggttccaagc	4860
	ctagaggctc	cttcogaaana	agcaccgact	gagcaaaagg	ctgggggtgca	ggagtgttac	4920
	catggtaatg	gacagagttta	togagccaca	tactccacca	ctgtcacaag	aaqaacctgc	4980
	caagcttggg	catctatgac	accacactcg	catagtggga	ccccagaeta	ctaccccaat	5040
	gctggcttga	toatgaacta	ctgcaggaat	ccagatgctg	tggcagctcc	ttattgttat	5100
25	acgagggalc	ccgc_lglcag	glgggaglac	lgcaacclga	cgcaalgclc	agcgcagaa	5160
	gggactgccc	tccgcoctcc	gactgttacc	ccggttccaa	gocctagagg	tccttccgaa	5220
	caagcaccga	ctgagcaaaag	gocctggggg	caggagtgct	accatggtaa	tggacagagt	5280
	tatcgaggca	catactccac	cactgtcaca	ggaaagacct	gccaagcttg	gtcatctatg	5340
	acaccacact	cgcatagtcc	gaccccagaa	taactaccaca	atgctggctt	gaacataaac	5400
	tactgcaggg	atccagatgc	tgtggcagct	ccttatgtgt	atacagggga	tcccgggtgc	5460
	agggtgggagt	actgcaacct	gaocgaatgc	tcagaocgag	aagggaactgc	cgtocogcct	5520
	ccgactgtta	ccccgggtcc	aaocctagag	gctccttccg	aaacaagcacc	gactgagcaa	5580
30	aggcctgggg	tgcaggagtg	ctaccatggt	aatggacaga	ggtatogagg	cacatactcc	5640
	accactgtca	caggaagaac	ctgccaagct	tggtcatcta	tgacaccaca	ctcgcatagt	5700
	oggaccccag	aatactacc	aaatgctggc	t-gateatga	actactgcag	gaatccagat	5760
	cctgtgqcaq	ctccctattg	ttatacgaag	catcccgggt	tcaggtggga	gtactgcaac	5820
	ctgacgcaat	gctcagaccg	agaagggaact	gcccgtccgc	ctccgactgt	tcccccggt	5880
	ccaggcoctag	aggctccttc	cgaacaagca	ccgactgagc	aaaggcoctg	ggcgcaaggag	5940
	tgctaaccatg	gtaaaggaca	gagttatcga	ggcacatact	ccaccactgt	cacagggaaga	6000
35	acctgccaag	cttggtcatc	tatgaaccca	cactcgcaat	gtcgggacccc	agaatactac	6060
	ccaaalgclg	gcllgalca_l	ggaclacclgc	aggaalccag	algc_glgcc	agclcccllal	6120
	tggtatacga	gggaaccocg	tgtaaggtgg	gagtaactgca	acctgacgca	atgctcagac	6180
	gcagaaggga	ctgcogctcc	gocctccgact	cttaccocgg	ttccaagcct	agaggctcct	6240
	tcogaacaaag	caccgactga	gcaaaaggcct	gggggtgcagg	agtgctacca	tggtaatgga	6300
	caaggttatc	gagggacata	ctccaccaact	gtcacaggaa	gaaocctgcca	agcttggcca	6360
	tctatgacac	cacactcga	tagtccgacc	ccagaatact	accocaaatgc	tggcttgatc	6420
40	atgaactact	gcaggaatcc	agatgctgtg	ccagctcctt	attgttatatac	gagggatccc	6480
	gglgclcaggl	gggaglacly	caacclgacg	caalgclcag	acgcagaagg	gaclgccglc	6540
	ggcctccga	ctgttaccoc	ggttccagc	ctagaggctc	cttccgacca	agcaccgact	6600
	gagcaaaagg	ctgggggtgca	ggagtgctac	catggtaatg	gacagagtta	togaggcaca	6660
	tactocacca	ctgtcaccag	asgaacctgc	caagcttggg	catctatgac	accacaotcg	6720
	catagtccga	ccccagaata	ctaccocaaat	cctggcttga	toatgaacta	ctcagaggaat	6780
	ccagatgctg	tggcagctcc	ttattgttat	acgagggato	ccgggtgcag	gtggggagtac	6840
45	tgcaacctga	cgcaatgctc	agaocgagaa	gggactgcog	tccgcoctcc	gactgttacc	6900
	ccgggtccaa	gocctagaggc	tccctccgaa	caagcaccga	ctgagcazag	gocctggggtg	6960
	caqgagtgct	accaatggtaa	tggacagagt	tatocgaggca	catactccac	cactgtocaca	7020
	ggaagaacct	gccaagcttg	gtcatctatg	acaccacact	cgcatagtccg	gaccccagaa	7080
	tactacccaa	atgcctggctc	gatcatgaac	tactgcagga	atccagatgc	tctggcagct	7140
	ccttatgtgt	atacagaggga	tcccgggtgc	agggtgggagt	actgcaacct	gacgcaatgc	7200
	tcagaocgag	aagggaactgc	cgtocogcct	ccgactgtta	ccccgggtcc	aagcctagag	7260
	gctccttccg	aacaagcaacc	gactgagcaa	aggcctgggg	tgcaaggagt	ctaccatggt	7320
50	aatggacaga	gttatcgagg	cacatactcc	accactgtca	caggaaagac	ctgccaagct	7380
	tggtcatcta	tqacaccaca	ctcgcatagt	ccgaccccag	aatactaacc	aatatgctggc	7440
	l_galcalga	aclaclgcag	gaalccagal	gclglgclcag	ctcccllallg	lla_laccagg	7500
	gatccocggg	toagggtggga	gtactgcaac	ctgacgcaat	gctcagacgc	agaagggaact	7560

55

EP 2 336 318 A1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

gcogtgcgc ctccgactgt tccccgggt ccaagcctag aggtccttc cgaacaagca 7620  
 ccgactgagc aaaggcctgg ggtgcaaggag tgcatacatg gtaatggaca gacttatoga 7680  
 cgcacalacL ccaccaclyL cacaggaaga acclgcacag clLggLcclL laLgacacca 7740  
 cactgcata gtccgacccc agaatactac ccaaatgctg gcttgatcat gaactactgc 7800  
 aggaatccag atgcctgtgc agctccttat tgttatacga gggatccogg tgcagcgg 7860  
 gagtactgca acctgacgca atgctcagac gcagaaggga ctgocgtgoc goctccgact 7920  
 ctacccocgg ttccaaagcc agaggctcct tccgaacaaq caocgactga gcaqagccct 7980  
 ggggtgcagg agtgcacca cggtaatgga cagagttatc gaggcacata ctccaccact 8040  
 ctcaactggaa gaacctgcca agcttggtca tctatgacac cacactogca tagtccgacc 8100  
 ccagaatact acccaaatgc tggcttgatc atgaactact gcaggaaatcc agatgctgtg 8160  
 gcagctcctt attgtatata gaggatccc ggtgtcaggt gggagtactg caacctgacg 8220  
 caatgctcag acgcagaagg gactgcogtc gcgctccga ctgttacccc ggtccaagc 8280  
 ctagaggctc ctccgaaaca agcaccgact gagcaaaagg ctgggtgca gtagtgcac 8340  
 catggtaatg gacagatgta tcaaggcaca tactccacca ctgtcacagg aagaacctgc 8400  
 caagcttggt catctatgac accacactcg catagtggga ccccagaata ctaccccaat 8460  
 gctggcttga toatgaacta ctgcaggaaat ccagatgctg tggcagctcc ttattgttat 8520  
 acgagggatc ccgctgtcag gtgggagtac tgcacctga cgcactgctc agacgcagaa 8580  
 gggactgccc togcctccc gactgttacc ccggttccaa gcctagaggc tocttccgaa 8640  
 caagcaccga clgagcaaaq gccLggggLg caggagLgcl accalggLaa LggacagL 8700  
 taticgagga catactccac cactgtcaca ggaagaacct gccaaagctg gtcactatg 8760  
 aacccactac ccgatagtoc qa.ccccagaa tactacccaa atgtctgctt gacataaac 8820  
 tactgcagga atccagatgc tgtggcagct ccttat.tgtt. ata.cgaggga tcccgggtgct 8880  
 aggtgggagt actgcacctc gacgcaatgc tcaagocgag aagggactgc cgtccgocct 8940  
 ccgactgtta ccccggttcc aagcctagag gctcctccgg aacaagcacc gactgagcag 9000  
 aggcctgggg tgcaggagtg ctaccaaggt aatggacaga gttatogagg cacataccc 9060  
 accalglca clggagaagc clgcccagcl LggLcclLla Lgacaccaca clcgcalagL 9120  
 cggaccocag aatactacc aaatgctggc tctgactgca actactgca gaa.tccagat 9180  
 gctgtggcag ctccctattg ttatacaggg gatcccggtg tcaggtggga gtactgcaac 9240  
 ctgacgcaat gctcagagcc agaagggact gcogtgcgc ctccgactgt taccccgggt 9300  
 ccaagcctag aggcctcttc cgaacaagca ccgactgagc agaggcctgg gctgcaagag 9360  
 tgcataccag gtaatggaca gcttatcga ggcacatact ccaccactgt cactggaga 9420  
 acctgccaag ctggctcatc tatgaca.cca cactgcata gtcggacccc agaatactac 9480  
 ccaaalglcl gclLgalcaL gaclacLlgc aggaalccag alglclLggc agclccLlal 9540  
 tgttatacga gggatcccg tgtcaggtgg agtactgca acctgacgca atgtctcagc 9600  
 gcagaaggga ctgcogtccc gctccgact gttaccocgg ttccaagcct agaggctcct 9660  
 tccgaacaag caccgactga gcagagccct ggggtgcagg agtgcctaca cggtaatgga 9720  
 caagcttatc gacgcacata ctccaccact ctcaactggaa gaacctgcca acctgtca 9780  
 tctatgacac caccctgca tegtccgacc ccagaatact acccaaatgc tggcttcatc 9840  
 atgaactact gcaggaaatcc agatgctgtg ccagctcctt attgttatata gagggatccc 9900  
 ggtgtcaggt gggagtctly caacctgacg caatgctcag accgagaagg gactgcccgc 9960  
 gcgctccga ctgttacccc ggttccagc ctagaggctc ctccgaaca agcaccgact 10020  
 gagcagaggc ctggggtgca gtagtctac cacggtaatg gacagagtta tccaggcaca 10080  
 tactccacca ctgtcactgg aegaacctgc caagcttggc catctatgac accacactcg 10140  
 catagtcggc ccccaagata ctaccaaat cctggcttga tcatgaaata ctgcagcaat 10200  
 ccagatcctg tccagcccc ttattgttat acgagggata ccagtgtcag gtagggatc 10260  
 tgcacactga caaatgctc agaocgagaa gggactgocg tgcgocctcc aactattacc 10320  
 ccgattccaa gcctagaggc tcttctgaa caagcaccac ctgagcaaaq gctgggggtg 10380  
 caggagtgcL accacggaaa tggacagagt tatcaaggca catacttoat tactgtaca 10440  
 ggaagaacct gccagclly gatcaagaac taactgcgaa atccagatcc tctggcagcc 10500  
 tactacccaa atgcggctL gatcaagaac taactgcgaa atccagatcc tctggcagcc 10560  
 ccttgggtgt atacaacaga tcccaggtgc aggtgggagt actgcaacct gacacgatc 10620  
 t.cagatgcag aatggactgc ct.t.cct.cct. ccgaatgta t.t.ctggct.c. aagcctagag 10680  
 cctttttttg aacaagcaat gactgaggaa acccccggg tacaggactg ctaactocat 10740  
 tatggacaga gttaccgagg cacatactcc accactgtca caggaaagaa ttgccagct 10800  
 tggctatcta tgacaccaca ccagcatagt ccgaccccag aaaactaacc aaatgctggc 10860  
 c.gaccagga acLacLgcaq gaalccagal gclgagallc gccLlggly llacacualg 10920  
 gatcccagtg tcaagtgga gtactgcaac ctgacaacat gcctggtgac agaatcaagt 10980  
 ctccctgcaa ctctcagct ggtcccagat ccaagcacag aggtctcttc tgaagaagca 11040  
 ccaacggagc aaagccccgg ggtcccagat tgcatacatg gtgatggaca gacttatoga 11100  
 gqctcattct ctaccactcc cacagggaag acatgtcagt ctgtgtctc taTgacaoca 11160  
 cactggcatc agaggacaac agaataatat ccaaatgggt gcctgaccag gaactatgc 11220  
 aggaatccag atgcTgagat tsgtccctgg tgttatacca tggatcccaa tgcagatgg 11280  
 gagtactgca acctgacaca atgtccagtg acagaatcaa gtgtccttgc gacgtccagc 11340  
 gctgtttctg aacaagcaac acggagcaa agccccacag tccaggactg ctacatgggt 11400  
 gatggacaga gttatcgagg ct.cattctcc accactgtta caggaaaggac atgtcagct 11460  
 tggctctcta tgacaccaca ctggcatcag agaaccacag aatactacc aaatgggtggc 11520  
 ctgaccagga actactgcaq gaatccagat cctgagatcc gcccttgggt ttataccatg 11580  
 gatcccagtg tcaagtgga gtactgcaac ctgacgcaat gtcagtgat ggaatcaact 11640  
 ctccctacaa ctcccagct ggtcccagtt ccaagcacag agcttccctc tgaagaagca 11700  
 ccaactgaaa acagcactgg ggtcccagac tgcataccag gtgatggaca gacttatoga 11760  
 ggcacactct ccaaccata cacaggcaga acatgtcagt cttggtctgc tatgacaoca 11820  
 calLggcalc ggaagclcc allalacLal ccaaalglcl gccLgaccag gaactacLgc 11880  
 aggaatccag atgcTgagat togccctgg tgttacacca tggatcccag tgtcaggtgg 11940

EP 2 336 318 A1

gagtactgca acctgacacg atgtccagtg acagaatcga gtgtccctcac aactcccaca 12000  
 g-gggccccgg ttccaagcac agaggctcct totgaaacaag caccacctga gaaaagccct 12060  
 g-ggglccagg allgclacca lgglgalgga cggagllalc gaggcatalc clccaccac 12120  
 g-cacacaggaa ggacctgtca atcttgggtca totatgatac cacactggca tcagaggacc 12180  
 ccagaaaaact acccaaatgc tggcctgacc gagaactact gcaggaatcc agattctggg 12240  
 aaacaacctt ggtgttacac aacogatccg tgtgtgaggt gggagtactg caatctgaca 12300  
 caatqctcag aaacagaatc aggtgtccta gagactccca ctattgttcc actccaagc 12360  
 atggaggctc attctgaagc agcaccactc gagcaaaccc ctgtgttcog gcagtgcac 12420  
 catggtaatg gccagagtta togaggcaaca tctccacca ctgtcacagg aaggacatgt 12480  
 caatcttggg catccatgac accacaccgg catcagagga cccagaaaa ctaccccaat 12540  
 gatggcctga caatgaaact ctgcaggaat ccagatgcog atacaggccc ttgggtgttt 12600  
 accatggacc ccagcatcag gtgggagtac tgcaacctga ccgatgtctc agacacagaa 12660  
 gggactgtgg tcgctcctcc gactgtcctc caggttccaa gccataggcc tctttctgaa 12720  
 caagactgta tqtllqqgaa tqggaaagga taccggggca aqaaggcaac cactqttact 12780  
 gggacgccc at gccaggaatg gctgcaccag gagccccata gcacacgcac gttcattcca 12840  
 gggacaaaaa aatgggcacg totggaaaaa aattaactgc gtaacctga tegtgaatc 12900  
 aatggctccct ggtgctacac aatgaatcca agaaaacttt ttgactactg tgatatccct 12960  
 ctctgtgcat cctctcatt tgattgtggg aagcctcaag tggagccgaa gaaatgtcct 13020  
 ggaagcally lagggggly lgggccac ccacalcccl ggccllyga aglcagllc 13080  
 agaacaaggt ttgaaagca cttctgtgga ggcaccttae tatcccaga gtgggtgctg 13140  
 actqctqctc actcctttaa qaagtctca aqqcctcat cctacaaqgt catcctcgt 13200  
 gcacaccaag aagtgaacct cgaatctcat gtcaggaaa tagaagtgtc taggctctc 13260  
 t-gggagccca cacaagcaga tattgccttg ctaaaagctaa gcaggcctgc cgtcctcact 13320  
 gcacaaagtaa tgcagcttg totgcatcc ccagactaca tggtcacgc caggactgaa 13380  
 tgttacatca ctggctgggg agaaacccaa ggtaccttg ggactggcct tccaaggaa 13440  
 ccccagclcc llglallga gaalgaagly lgcaalcacl alaaglala llglgclgag 13500  
 catttggcca gaggcactga cagttgpcag ggtgacagt gaggcctct gcttgcctc 13560  
 gagaaggaca aatacattt acaaggagtc acttcttggg gtcttggctg tgcacgccc 13620  
 aataagcctg gtgtctatgc tctgtttca aggtttgtta ctggattga gggaaatgat 13680  
 aqaaataatt aattggacg qagacagagt caagcatcaa cctacttaca aactgaaacg 13740  
 tgggtaagga tttgcatgc tggaaataat agacagcaat caaacgaaga cactgttccc 13800  
 agctaccagc tatcccaaac ctggcattt ttggatttt tgtgtataag ct-ttaaggt 13860  
 c-gaactgaca aallclgla laagglglca lagclalgac allglleaa aalaaaccl 13920  
 gcacttattt tgatttga 13938

<210> 861  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

<400> 861  
 cagctccttc ttgtatacg aggga 25

<210> 862  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

<400> 862  
 lgcglclgag callgcgl 18

<210> 863  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Probe

<400> 863  
 cccagatgca ggtcggata ctgc 24

<210> 864  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

55

EP 2 336 318 A1

	<220> <223> Antisense Oligonucleotide	
5	<400> 864 gcctcagtct tcttcgcacc	20
	<210> 865 <211> 20 <212> DNA <213> Artificial Sequence	
10	<220> <223> Antisense Oligonucleotide	
	<400> 865 gcctcagtct tattcgcacc	20
15	<210> 866 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
20	<400> 866 gcctcagtat tattcgcacc	20
	<210> 867 <211> 20 <212> DNA <213> Artificial Sequence	
25	<220> <223> Antisense Oligonucleotide	
	<400> 867 gcctcattat tattcgcacc	20
30	<210> 868 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
35	<400> 868 gcctcattat tattcgcacc	20
	<210> 869 <211> 20 <212> DNA <213> Artificial Sequence	
40	<220> <223> Antisense Oligonucleotide	
	<400> 869 gcctcattat tattatcacc	20
45	<210> 870 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
50	<400> 870 gcctaattat tattatcacc	20
	<210> 871	
55		

EP 2 336 318 A1

<211> 19  
 <212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 871  
 gcctcagtct gcttgcac 19  
 <210> 872  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 872  
 gcctcagtct gcttgcac 18  
 <210> 873  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 873  
 gcctcagtct gcttc 15  
 <210> 874  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 874  
 cctcagtctg cttgcac 18  
 <210> 875  
 <211> 16  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 875  
 ctcagtctgc ttgcac 16  
 <210> 876  
 <211> 14  
 <212> DNA  
 <213> Artificial Sequence  
 40  
 <220>  
 <223> Antisense Oligonucleotide  
 <400> 876  
 tcagtctgct tcgc 14  
 <210> 877  
 <211> 10  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Antisense Oligonucleotide  
 55

EP 2 336 318 A1

	<400> 877 gcctcagtct	10
5	<210> 878 <211> 10 <212> DNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
10	<400> 878 gcttcgcacc	10
	<210> 879 <211> 20 <212> RNA <213> Artificial Sequence	
15	<220> <223> Antisense Oligonucleotide	
	<400> 879 uggaagccau acaccucuu	20
20	<210> 880 <211> 20 <212> RNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
25	<400> 880 ugaccaggac ugccuguucu	20
	<210> 881 <211> 20 <212> RNA <213> Artificial Sequence	
30	<220> <223> Antisense Oligonucleotide	
	<400> 881 gaaauagggcu guaꝑꝑꝑꝑuaa	20
35	<210> 882 <211> 20 <212> RNA <213> Artificial Sequence	
	<220> <223> Antisense Oligonucleotide	
40	<400> 882 uuuacugauc aaauuguauc	20
	<210> 883 <211> 20 <212> RNA <213> Artificial Sequence	
45	<220> <223> Antisense Oligonucleotide	
	<400> 883 uggaauucug guaꝑꝑꝑꝑaag	20
50	<210> 884 <211> 20 <212> RNA <213> Artificial Sequence	
55		



EP 2 336 318 A1

<220>  
 <223> Antisense Oligonucleotide  
 5 <400> 884  
 aaaucaaaug auugcuuuugu 20  
 <210> 885  
 <211> 20  
 <212> RNA  
 <213> Artificial Sequence  
 10 <220>  
 <223> Antisense Oligonucleotide  
 <400> 885  
 gugaugacac uugauuuuaaa 20  
 <210> 886  
 <211> 20  
 <212> RNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> Antisense Oligonucleotide  
 <400> 886  
 gaagcugccu cuucuuccca 20  
 <210> 887  
 <211> 20  
 <212> RNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Antisense Oligonucleotide  
 <400> 887  
 gagaguuggu cugaaaaaac 20  
 <210> 888  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 25 <220>  
 <223> Antisense Oligonucleotide  
 <400> 888  
 g-gcgcgcga gcccgaaatc 20  
 <210> 889  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> Antisense Oligonucleotide  
 <400> 889  
 cuucuggcau ccgguuuagt t 21  
 <210> 890  
 <211> 466  
 <212> DNA  
 <213> M. fascicularis  
 35 <220>  
 <221> misc\_feature  
 <222> 9  
 <223> n = A,T,U or G  
 <400> 890  
 40  
 45  
 50  
 55

## EP 2 336 318 A1

5 ggatcggcng accctgagct gcatatggct ggtaatc-aa aaggagccta ccaaataat 60  
gaaataaaac acatctatac catctcttct gctgccttat cagcaagcta caaagcagac 120  
actggttgcta aggttcaggg tgtggagttt agccatcggc tcaacacaga catcgctggg 180  
ctggcttcag ccatgacat tagcacaac tataattcag actcattgca tttcagcaat 240  
gtcttcatt ctgtaatggc tccatttacc atgaccattg atacacatac aaatggcaac 300  
gggaaaacttg ttctctgggg agaacatact gggcagcgtg atagcaaatt cctggtgaaa 360  
gcagAACctc tggcattcac tttctctcat gattacaaag gctccacgag tcatcatctc 420  
atgtctagga aaagcatcag tgcagctctt gaacacaaag tcagta 456

10 <210> 891  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

15 <400> 891  
gctcagctct gctttacacc 20

20 <210> 892  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

25 <400> 892  
agattaccag ccatatgcag 20

### Claims

- 30
1. An antisense oligonucleotide compound 8 to 50 nucleobases in length comprising at least 8 contiguous nucleotides of SEQ ID NO:247.
  - 35 2. The antisense oligonucleotide compound of claim 1, wherein the antisense oligonucleotide compound has a sequence comprising SEQ ID NO:247.
  3. The antisense oligonucleotide compound of claim 1, wherein said compound is twelve to thirty nucleobases in length, or fourteen to twenty nucleobases in length.
  - 40 4. The antisense oligonucleotide compound of claim 2, wherein the antisense oligonucleotide compound has a sequence consisting of SEQ ID NO:247.
  5. The antisense oligonucleotide compound of any preceding claim, wherein the antisense oligonucleotide compound is an oligonucleotide mimetic compound.
  - 45 6. The antisense oligonucleotide compound of claim 5, wherein the oligonucleotide mimetic compound is a chimeric phosphorothioate oligonucleotide compound.
  7. The antisense oligonucleotide compound of claim 6, wherein the chimeric phosphorothioate oligonucleotide compound comprises 2'-methoxyethoxyl nucleotide wings and a 2'-deoxynucleotide gap.
  - 50 8. The antisense oligonucleotide compound of claim 7, wherein the chimeric phosphorothioate oligonucleotide compound comprises ten 2'-deoxynucleotides.
  - 55 9. The antisense oligonucleotide compound according to claim 8, wherein said compound is 20 nucleobases in length having a sequence of nucleobases as set forth in SEQ ID NO:247 and comprising 5 -methylcytidine at nucleobases 2, 3, 5, 9, 12, 15, 17, 19, and 20, wherein every internucleoside linkage is a phosphorothioate linkage, nucleobases 1-5 and 16-20 comprise a 2'-methoxyethoxyl modification, and nucleobases 6-15 are deoxynucleotides.

10. A pharmaceutically acceptable salt of a compound of any preceding claim.

11. A composition comprising the antisense oligonucleotide compound or pharmaceutically acceptable salt of any preceding claim and a pharmaceutically acceptable carrier or diluent.

5

12. The compound or pharmaceutically acceptable salt of any of claims 1-10, for use in therapy.

13. The compound or pharmaceutically acceptable salt of claim 12, for use in treating a human having a disease or condition associated with apolipoprotein B expression, wherein:

10

(i) the human has a condition associated with abnormal lipid metabolism

(ii) the human has a condition associated with abnormal cholesterol metabolism

(iii) the human has a cardiovascular disease, such as atherosclerosis

(iv) the human has an abnormal metabolic condition associated with apolipoprotein B expression, such as hyperlipidemia;

15

(v) the human has diabetes; or

(vi) the human is obese.

14. The compound or pharmaceutically acceptable salt of claim 12, for use in:

20

(i) preventing or delaying the onset of an increase in glucose levels in an animal;

(ii) modulating serum cholesterol levels in an animal; or

(iii) modulating lipoprotein levels in an animal.

15. The compound or pharmaceutically acceptable salt of claim 14, wherein:

25

(i) the glucose levels are serum glucose levels;

(ii) the glucose levels are plasma glucose levels;

(iii) the lipoprotein is VLDL;

(iv) the lipoprotein is HDL; or

(v) the lipoprotein is LDL.

30

35

40

45

50

55



Europäisches  
Patentamt  
European  
Patent Office  
Office européen  
des brevets

## EUROPEAN SEARCH REPORT

Application Number  
EP 10 18 0483

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	OSTRANDER E A ET AL: "DOG (CLONE: CXX.371) PRIMER FOR STS 371, 5' END, SEQUENCE TAGGED SITE", EMBL, 18 September 1993 (1993-09-18), XP002392182, * the whole document * -----	1,3,5,6	INV. C12N15/113 A61K48/00 C07H21/00
X	OSTRANDER E A ET AL: "DOG PRIMER FOR STS 610, 3' END, SEQUENCE TAGGED SITE", EMBL, 6 January 1994 (1994-01-06), XP002392183, * the whole document * -----	1,3,5,6	
X	TANG QIDONG ET AL: "The Inhibition of Antisense Oligodeoxynucleotides on the Expression of Apolipoprotein B in Rat Liver Cells", CHINESE JOURNAL OF ARTERIOSCLEROSIS,, vol. 7, no. 4, 1 January 1999 (1999-01-01), XP002632627, * the whole document * -----	1-15	
X	EGGERMAN ET AL: "Use of Oligonucleotides to Target Nucleic Acid Sequences Encoding Apolipoprotein B to Decrease Serum Apolipoprotein B and Cholesterol Levels", FEDERAL REGISTER,, vol. 65, no. 110, 7 June 2000 (2000-06-07), XP002632628, * the whole document * ----- -/--	1-15	
The present search report has been drawn up for all claims			
5	Place of search Munich	Date of completion of the search 14 April 2011	Examiner Persson, Anna
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document	

EPO FORM 1503 03.02 (P04C01)



## EUROPEAN SEARCH REPORT

Application Number  
EP 10 18 0483

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	GRAHAM M J ET AL: "INHIBITION OF APOB-100 AS A THERAPEUTIC STRATEGY FOR THE TREATMENT OF HYPERLIPIDEMIAS", CIRCULATION, LIPPINCOTT WILLIAMS & WILKINS, US, vol. 106, no. SUPPL. 19, 5 November 2002 (2002-11-05), pages II-109, XP009070023, ISSN: 0009-7322 * the whole document *	1-15	
X,D	WO 01/12789 A2 (CHAN LAWRENCE [US]; TENG BA BIE [US]; TIEBEL MARGRET [DE] BAYLOR COLLE) 22 February 2001 (2001-02-22) * the whole document *	1-15	
X,P	WO 03/011887 A2 (ISIS PHARMACEUTICALS INC [US]; CROOKE ROSANNE M [US]; GRAHAM MARK J [U]) 13 February 2003 (2003-02-13) * pages 95-98; examples 15-16; table 2 *	1-15	
A	LAW S W ET AL: "Human Apolipoprotein B-100: Cloning, Analysis of Liver mRNA, and Assignment of the Gene to Chromosome 2", 19851201, vol. 82, no. 24, 1 December 1985 (1985-12-01), pages 8340-8344, XP002979995, * the whole document *	1-15	TECHNICAL FIELDS SEARCHED (IPC)
----- -/--			
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 14 April 2011	Examiner Persson, Anna
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document	

5  
EPO FORM 1503 03.82 (P04C01)



EUROPEAN SEARCH REPORT

Application Number  
EP 10 18 0483

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	SMITH LISE ET AL: "Rational selection of antisense oligonucleotide sequences", EUROPEAN JOURNAL OF PHARMACEUTICAL SCIENCES, ELSEVIER, AMSTERDAM, NL, vol. 11, no. 3, 1 September 2000 (2000-09-01), pages 191-198, XP002372482, ISSN: 0928-0987, DOI: DOI:10.1016/S0928-0987(00)00100-7 * the whole document *	1-15	TECHNICAL FIELDS SEARCHED (IPC)
A	ROJANASAKUL Y: "ANTISENSE OLIGONUCLEOTIDE THERAPEUTICS: DRUG DELIVERY AND TARGETING", ADVANCED DRUG DELIVERY REVIEWS, ELSEVIER BV, AMSTERDAM, NL, vol. 18, 1 January 1996 (1996-01-01), pages 115-131, XP002910299, ISSN: 0169-409X, DOI: DOI:10.1016/0169-409X(95)00020-8 * the whole document *	1-15	
T	BRADLEY J ET AL: "Small, dense, LDL particle concentration is significantly reduced after treatment with an antisense inhibitor of ApoB in healthy volunteers", CIRCULATION, LIPPINCOTT WILLIAMS & WILKINS, US, vol. 112, no. 17, SUPPL. S, 1 October 2005 (2005-10-01), pages II-133/II, XP009069945, ISSN: 0009-7322 * the whole document *		
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 14 April 2011	Examiner Persson, Anna
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

5  
EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 10 18 0483

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-04-2011

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0112789	A2	22-02-2001	AU 6910100 A	13-03-2001
-----				
WO 03011887	A2	13-02-2003	AU 2002326481 B2	13-11-2008
			CA 2455228 A1	13-02-2003
			EP 1419168 A2	19-05-2004
			EP 2174945 A1	14-04-2010
			JP 4471650 B2	02-06-2010
			JP 2005503142 T	03-02-2005
			JP 2009055925 A	19-03-2009
			US 2003215943 A1	20-11-2003
			US 2008242629 A1	02-10-2008
-----				

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 0315493 W [0001]
- US 42623402 P [0001]
- US 14719602 A [0001]
- US 13598502 A [0001]
- US 92003301 A [0001]
- US 5786206 A [0011]
- US 6156315 A [0017]
- WO 9918986 A [0017]
- US 6096516 A [0018]
- EP 911344 A [0019]
- US 6309844 B [0019]
- WO 0130354 A [0020]
- US 5220006 A [0021]
- WO 0112789 A [0022]
- US 3687808 A [0095] [0104] [0105]
- US 4469863 A [0095] [0242]
- US 4476301 A [0095]
- US 5023243 A [0095] [0247]
- US 5177196 A [0095]
- US 5188897 A [0095]
- US 5264423 A [0095]
- US 5276019 A [0095]
- US 5278302 A [0095]
- US 5286717 A [0095]
- US 5321131 A [0095]
- US 5399676 A [0095]
- US 5405939 A [0095]
- US 5453496 A [0095]
- US 5455233 A [0095]
- US 5466677 A [0095] [0097]
- US 5476925 A [0095] [0246]
- US 5519126 A [0095]
- US 5536821 A [0095]
- US 5541306 A [0095]
- US 5550111 A [0095]
- US 5563253 A [0095]
- US 5571799 A [0095]
- US 5587361 A [0095]
- US 5194599 A [0095]
- US 5565555 A [0095]
- US 5527899 A [0095]
- US 5721218 A [0095]
- US 5672697 A [0095]
- US 5625050 A [0095] [0243]
- US 5034506 A [0097] [0100]
- US 5166315 A [0097]
- US 5185444 A [0097]
- US 5214134 A [0097]
- US 5216141 A [0097]
- US 5235033 A [0097]
- US 5264562 A [0097] [0250]
- US 5264564 A [0097] [0250]
- US 5405938 A [0097]
- US 5434257 A [0097]
- US 5470967 A [0097]
- US 5489677 A [0097] [0100] [0249]
- US 5541307 A [0097]
- US 5561225 A [0097]
- US 5596086 A [0097]
- US 5602240 A [0097] [0100] [0249]
- US 5610289 A [0097] [0243] [0249]
- US 5608046 A [0097] [0107]
- US 5618704 A [0097]
- US 5623070 A [0097]
- US 5663312 A [0097]
- US 5633360 A [0097]
- US 5677437 A [0097]
- US 5792608 A [0097]
- US 5646269 A [0097]
- US 5677439 A [0097]
- US 5539082 A [0099] [0252]
- US 5714331 A [0099]
- US 5719262 A [0099] [0252]
- WO 9839352 A [0102]
- WO 9914226 A [0102]
- US 4981957 A [0103]
- US 5118800 A [0103]
- US 5319080 A [0103]
- US 5359044 A [0103]
- US 5393878 A [0103]
- US 5446137 A [0103]
- US 5466786 A [0103]
- US 5514785 A [0103] [0107]
- US 5519134 A [0103]
- US 5567811 A [0103]
- US 5576427 A [0103]
- US 5591722 A [0103]
- US 5597909 A [0103]
- US 5610300 A [0103]
- US 5627053 A [0103]
- US 5639873 A [0103]
- US 5646265 A [0103]
- US 5658873 A [0103]
- US 5670633 A [0103] [0211]
- US 5792747 A [0103]
- US 5700920 A [0103]
- US 4845205 A [0105]
- US 5130302 A [0105] [0248]



EP 2 336 318 A1

- US 5134066 A [0105]
- US 5175273 A [0105]
- US 5367066 A [0105]
- US 5432272 A [0105]
- US 5457187 A [0105]
- US 5459255 A [0105]
- US 5484908 A [0105]
- US 5502177 A [0105]
- US 5525711 A [0105]
- US 5552540 A [0105]
- US 5587469 A [0105]
- US 5594121 A [0105]
- US 5596091 A [0105]
- US 5614617 A [0105]
- US 5645985 A [0105]
- US 5830653 A [0105]
- US 5763588 A [0105]
- US 6005096 A [0105]
- US 5681941 A [0105]
- US 5750692 A [0105]
- US 9209196 W [0106]
- US 09334130 B [0106]
- US 4828979 A [0107]
- US 4948882 A [0107]
- US 5218105 A [0107]
- US 5525465 A [0107]
- US 5541313 A [0107]
- US 5545730 A [0107]
- US 5552538 A [0107]
- US 5578717 A [0107]
- US 5580731 A [0107]
- US 5591584 A [0107]
- US 5109124 A [0107]
- US 5118802 A [0107]
- US 5138045 A [0107]
- US 5414077 A [0107]
- US 5486603 A [0107]
- US 5512439 A [0107]
- US 5578718 A [0107]
- US 4587044 A [0107]
- US 4605735 A [0107]
- US 4667025 A [0107]
- US 4762779 A [0107]
- US 4789737 A [0107]
- US 4824941 A [0107]
- US 4835263 A [0107]
- US 4876335 A [0107]
- US 4904582 A [0107]
- US 4958013 A [0107]
- US 5082830 A [0107]
- US 5112963 A [0107]
- US 5214136 A [0107]
- US 5245022 A [0107]
- US 5254469 A [0107]
- US 5258506 A [0107]
- US 5262536 A [0107]
- US 5272250 A [0107]
- US 5292873 A [0107]
- US 5317098 A [0107]
- US 5371241 A [0107]
- US 5391723 A [0107]
- US 5416203 A [0107]
- US 5451463 A [0107]
- US 5510475 A [0107]
- US 5512667 A [0107]
- US 5565552 A [0107]
- US 5567810 A [0107]
- US 5574142 A [0107]
- US 5585481 A [0107]
- US 5587371 A [0107]
- US 5595726 A [0107]
- US 5597696 A [0107]
- US 5599923 A [0107]
- US 5599928 A [0107]
- US 5688941 A [0107]
- US 5013830 A [0109]
- US 5149797 A [0109]
- US 5220007 A [0109]
- US 5256775 A [0109] [0244]
- US 5366878 A [0109] [0244]
- US 5403711 A [0109]
- US 5491133 A [0109]
- US 5565350 A [0109]
- US 5623065 A [0109] [0257]
- US 5652355 A [0109]
- US 5652356 A [0109]
- US 5700922 A [0109] [0252]
- US 5108921 A [0112]
- US 5354844 A [0112]
- US 5416016 A [0112]
- US 5459127 A [0112]
- US 5521291 A [0112]
- US 5543158 A [0112]
- US 5547932 A [0112]
- US 5583020 A [0112]
- US 5591721 A [0112]
- US 4426330 A [0112] [0155]
- US 4534899 A [0112] [0155]
- US 5013556 A [0112] [0155]
- US 5213804 A [0112] [0155]
- US 5227170 A [0112]
- US 5264221 A [0112] [0156]
- US 5356633 A [0112] [0155]
- US 5395619 A [0112]
- US 5417978 A [0112]
- US 5462854 A [0112]
- US 5469854 A [0112]
- US 5512295 A [0112]
- US 5527528 A [0112]
- US 5534259 A [0112]
- US 5543152 A [0112] [0154]
- US 5556948 A [0112] [0155]
- US 5580575 A [0112]
- US 5595756 A [0112]
- WO 9324510 A, Gosselin [0114]
- WO 9426764 A [0114]

## EP 2 336 318 A1

- US 5770713 A, Imbach [0114]
- US 09315298 B [0121] [0122]
- US 08886829 B [0122]
- US 09108673 B [0122]
- US 09256515 B [0122]
- US 09082624 B [0122]
- US 4837028 A [0154]
- WO 8804924 A [0154]
- WO 9713499 A, Lim [0154]
- EP 0445131 B1 [0155]
- WO 9004384 A [0155]
- EP 0496813 B1 [0155]
- WO 9105545 A [0155]
- US 5225212 A, Martin [0155]
- WO 9420073 A, Zalipsky [0155]
- WO 9610391 A, Choi [0155]
- US 5540935 A, Miyazaki [0155]
- WO 9640062 A, Thierry [0156]
- US 5665710 A, Rahman [0156]
- WO 9704787 A, Love [0156]
- US 5705188 A, Junichi [0171] [0529]
- WO 9730731 A, Lollo [0171] [0529]
- US 4501729 A [0186]
- US 6159946 A [0201]
- US 6547787 B [0202]
- US 6495532 B [0203]
- US 5399331 A [0203]
- US 4885172 A [0203]
- US 5059421 A [0203]
- US 5171578 A [0203]
- US 6217886 B [0203]
- US 4818542 A [0204]
- US 6562864 B [0205]
- US 6433012 B [0206]
- US 6368356 B [0207]
- US 6344035 B [0207]
- US 6344028 B [0207]
- US 6287285 B [0207]
- US 6200304 B [0207]
- US 5824049 B [0207]
- US 5749915 B [0207]
- US 5674242 B [0207]
- US 5670161 B [0207]
- US 5609629 B [0207]
- US 5593974 B [0207]
- US 5470307 B [0207]
- US 5506351 A [0209]
- WO 9402501 A1, McGee, D. P. C., Cook, P. D., Guinasso, C. J. [0235]
- US 5508270 A [0241]
- US 9400902 W [0245]
- US 9306976 W [0245]
- WO 9417093 A [0245]
- WO 9402499 A [0245]
- US 5177198 A [0248]
- US 5378825 A [0249]
- US 5386023 A [0249]
- US 5223618 A [0251]

### Non-patent literature cited in the description

- **Davidson ; Shelness.** *Annu. Rev. Nutr.*, 2000, vol. 20, 169-193 [0005] [0006] [0007] [0008] [0009]
- **Law et al.** *Proc. Natl. Acad. Sci. U. S. A.*, 1985, vol. 82, 8340-8344 [0010]
- **Deeb et al.** *Proc. Natl. Acad. Sci. U. S. A.*, 1986, vol. 83, 419-422 [0010]
- **Kim ; Young.** *J. Lipid Res.*, 1998, vol. 39, 703-723 [0012] [0013] [0014]
- **Nishina et al.** *J. Lipid Res.*, 1990, vol. 31, 859-869 [0012]
- **McCormick et al.** *J. Biol. Chem.*, 1997, vol. 272, 23616-23622 [0013]
- **Farese et al.** *Proc. Natl. Acad. Sci. U. S. A.*, vol. 92, 1774-1778 [0014]
- **Kim ; Young.** *J. Lipid Res.*, vol. 39, 703-723 [0014]
- **Veniant et al.** *J. Clin. Invest.*, 1997, vol. 100, 180-188 [0014]
- **Seed et al.** *N. Engl. J. Med.*, 1990, vol. 322, 1494-1499 [0015]
- **Sandkamp et al.** *Clin. Chem.*, 1990, vol. 36, 20-23 [0015]
- **Nowak-Gottl et al.** *Pediatrics*, 1997, vol. 99, E11 [0015]
- **Katan ; Beynen.** *Am. J. Epidemiol.*, 1987, vol. 125, 387-399 [0016]
- **Vessby et al.** *Atherosclerosis*, 1982, vol. 44, 61-71 [0016]
- **Hajjar ; Nachman.** *Annu. Rev. Med.*, 1996, vol. 47, 423-442 [0016]
- **Brazma ; Vilo.** *FEBS Lett.*, 2000, vol. 480, 17-24 [0088]
- **Celis et al.** *FEBS Lett.*, 2000, vol. 480, 2-16 [0088]
- **Madden et al.** *Drug Discov. Today*, 2000, vol. 5, 415-425 [0088]
- **Prashar ; Weissman.** *Methods Enzymol.*, 1999, vol. 303, 258-72 [0088]
- **Sutcliffe et al.** *Proc. Natl. Acad. Sci. U. S. A.*, 2000, vol. 97, 1976-81 [0088]
- **Celis et al.** *FEBS Lett.*, 2000, vol. 480, 2-16 [0088]
- **Jungblut et al.** *Electrophoresis*, 1999, vol. 20, 2100-10 [0088]
- **Larsson et al.** *J. Biotechnol.*, 2000, vol. 80, 143-57 [0088]
- **Fuchs et al.** *Anal. Biochem.*, 2000, vol. 286, 91-98 [0088]
- **Larson et al.** *Cytometry*, 2000, vol. 41, 203-208 [0088]
- **Jurecic ; Belmont.** *Curr. Opin. Microbiol.*, 2000, vol. 3, 316-21 [0088]

- **Carulli et al.** *J. Cell Biochem.*, 1998, vol. 31, 286-96 [0088]
- **Going ; Gusterson.** *Eur. J. Cancer*, 1999, vol. 35, 1895-904 [0088]
- **To.** *Comb. Chem. High Throughput Screen*, 2000, vol. 3, 235-41 [0088]
- **Nielsen et al.** *Science*, 1991, vol. 254, 1497-1500 [0099]
- **Martin et al.** *Helv. Chim. Acta*, 1995, vol. 78, 486-504 [0101]
- The Concise Encyclopedia Of Polymer Science And Engineering. John Wiley & Sons, 1990, 858-859 [0104]
- **Englisch et al.** *Angewandte Chemie*. 1991, vol. 30, 613 [0104]
- **Sanghvi, Y.S.** Antisense Research and Applications. CRC Press, 1993, 289-302 [0104]
- Antisense Research and Applications. CRC Press, 276-278 [0104]
- **Letsinger et al.** *Proc. Natl. Acad. Sci. USA*, 1989, vol. 86, 6553-6556 [0106]
- **Manoharan et al.** *Bioorg. Med. Chem. Lett.*, 1994, vol. 4, 1053-1060 [0106]
- **Manoharan et al.** *Ann. N.Y. Acad. Sci.*, 1992, vol. 660, 306-309 [0106]
- **Manoharan et al.** *Bioorg. Med. Chem. Lett.*, 1993, vol. 3, 2765-2770 [0106]
- **Oberhauser et al.** *Nucl. Acids Res.*, 1992, vol. 20, 533-538 [0106]
- **Saison-Behmoaras et al.** *EMBO J.*, 1991, vol. 10, 1111-1118 [0106]
- **Kabanov et al.** *FEBS Lett.*, 1990, vol. 259, 327-330 [0106]
- **Svinarchuk et al.** *Biochimie*, 1993, vol. 75, 49-54 [0106]
- **Manoharan et al.** *Tetrahedron Lett.*, 1995, vol. 36, 3651-3654 [0106]
- **Shea et al.** *Nucl. Acids Res.*, 1990, vol. 18, 3777-3783 [0106]
- **Manoharan et al.** *Nucleosides & Nucleotides*, 1995, vol. 14, 969-973 [0106]
- **Manoharan et al.** *Tetrahedron Lett*, 1995, vol. 36, 3651-3654 [0106]
- **Mishra et al.** *Biochim. Biophys. Acta*, 1995, vol. 1264, 229-237 [0106]
- **Crooke et al.** *J. Pharmacol. Exp. Ther.*, 1996, vol. 277, 923-937 [0106]
- **Berge et al.** *Pharmaceutical Salts. J. of Pharma Sci.*, 1977, vol. 66, 1-19 [0116]
- **Idson.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, 1988, vol. 1, 199 [0128] [0129] [0130] [0132] [0135]
- **Rosoff.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, 1988, vol. 1, 245 [0128] [0135] [0144]
- **Block.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, 1988, vol. 2, 335 [0128]
- **Higuchi et al.** *Remington's Pharmaceutical Sciences.* Mack Publishing Co, 1985, 301 [0128]
- **Rieger.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, 1988, vol. 1, 285 [0130]
- **Block.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, 1988, vol. 1, 335 [0132] [0137]
- **Rosoff.** *Pharmaceutical Dosage Forms.* Lieberman. Marcel Dekker, Inc, 1988, vol. 1, 245 [0136]
- **Leung ; Shah.** *Controlled Release of Drugs: Polymers and Aggregate Systems.* VCH Publishers, 1989, 185-215 [0136]
- **Schott.** *Remington's Pharmaceutical Sciences.* Mack Publishing Co, 1985, 271 [0136]
- **Rosoff.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, vol. 1, 245 [0137]
- **Constantinides et al.** *Pharmaceutical Research*, 1994, vol. 11, 1385-1390 [0139]
- **Ritschel.** *Meth. Find. Exp. Clin. Pharmacol.*, 1993, vol. 13, 205 [0139]
- **Constantinides et al.** *Pharmaceutical Research*, 1994, vol. 11, 1385 [0139]
- **Ho et al.** *J. Pharm. Sci.*, 1996, vol. 85, 138-143 [0139]
- **Lee et al.** *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, 92 [0140] [0166] [0167] [0168] [0169] [0170] [0191] [0521] [0523] [0525] [0527] [0528]
- **Wang et al.** *Biochem. Biophys. Res. Commun.*, 1987, vol. 147, 980-985 [0148]
- **Zhou et al.** *Journal of Controlled Release*, 1992, vol. 19, 269-274 [0149]
- **Weiner et al.** *Journal of Drug Targeting*, 1992, vol. 2, 405-410 [0151]
- **Plessis et al.** *Antiviral Research*, 1992, vol. 18, 259-265 [0151]
- **Hu et al.** *S.T.P. Pharma. Sci.*, 1994, vol. 4 (6), 466 [0152]
- **Allen et al.** *FEBS Letters*, 1987, vol. 223, 42 [0153]
- **Wu et al.** *Cancer Research*, 1993, vol. 53, 3765 [0153]
- **Papahadjopoulos et al.** *Ann. N.Y. Acad. Sci.*, 1987, vol. 507, 64 [0154]
- **Gabizon et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1988, vol. 85, 6949 [0154]
- **Sunamoto et al.** *Bull. Chem. Soc. Jpn.*, 1980, vol. 53, 2778 [0155]
- **Illum et al.** *FEBS Lett.*, 1984, vol. 167, 79 [0155]
- **Klibanov et al.** *FEBS Lett.*, 1990, vol. 268, 235 [0155]
- **Blume et al.** *Biochimica et Biophysica Acta*, 1990, vol. 1029, 91 [0155]
- **Rieger.** *Pharmaceutical Dosage Forms.* Marcel Dekker, Inc, 1988, 285 [0158] [0163]
- **Lee et al.** *Critical Reviews in Therapeutic Drug Carrier Systems.* 1991, 92 [0165]
- **Takahashi et al.** *J. Pharm. Pharmacol.*, 1988, vol. 40, 252 [0166]
- **Muranishi.** *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, vol. 7, 1-33 [0167] [0168] [0169]

- **El Hariri et al.** *J. Pharm. Pharmacol.*, 1992, vol. 44, 651-654 [0167]
- **Brunton et al.** Goodman & Gilman's The Pharmacological Basis of Therapeutics. McGraw-Hill, 1996, 934-935 [0168]
- **Swinyard.** Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990, 782-783 [0168]
- **Yamamoto et al.** *J. Pharm. Exp. Ther.*, 1992, vol. 263, 25 [0168] [0525]
- **Yamashita et al.** *J. Pharm. Sci.*, 1990, vol. 79, 579-583 [0168]
- **Jarrett.** *J. Chromatogr.*, 1993, vol. 618, 315-339 [0169]
- **Buur et al.** *J. Control Rel.*, 1990, vol. 14, 43-51 [0169]
- **Muranishi.** *Critical Reviews in Therapeutic Drug Carrier Systems*, 1980, vol. 7, 1-33 [0170]
- **Yamashita et al.** *J. Pharm. Pharmacol.*, 1987, vol. 39, 621-626 [0170]
- **Miyao et al.** *Antisense Res. Dev.*, 1995, vol. 5, 115-121 [0173]
- **Takakura et al.** *Antisense & Nuc1 . Acid Drug Dev.*, 1996, vol. 6, 177-183 [0173]
- The Merck Manual of Diagnosis and Therapy. 1987, 1206-1228 [0194]
- The Merck Manual of Diagnosis and Therapy. 1987, 2499-250646-49 [0194]
- **Simons et al.** *Nature*, 1992, vol. 359, 67-70 [0201]
- **Sanghvi.** *Nucleic Acids Research*, 1993, vol. 21, 3197-3203 [0210]
- **Kawasaki.** *J. Med. Chem.*, 1993, vol. 36, 831-841 [0211]
- **Martin, P.** *Helvetica Chimica Acta*, 1995, vol. 78, 486-504 [0216]
- Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications. *Bioorganic & Medicinal Chemistry*, 1996, vol. 4, 5-23 [0252]
- **Chiang et al.** *J. Biol. Chem.*, 1991, vol. 266, 18162-18171 [0258]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1993, vol. 1, 4.1.1-4.2.94.5.1-4.5.3 [0283]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1996, vol. 1, 4.2.1-4.2.9 [0283]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1997, vol. 2, 11.12.1-11.12.9 [0284]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1997, vol. 2, 11.4.1-11.11.5 [0284]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1998, vol. 2, 10.16.1-10.16.11 [0285]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1997, vol. 2, 10.8.1-10.8.21 [0285]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1991, vol. 2, 11.2.1-11.2.22 [0285]
- **Miura et al.** *Clin. Chem.*, 1996, vol. 42, 1758-1764 [0286]
- **Ausubel, F.M. et al.** Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1993, vol. 1, 4.5.1-4.5.3 [0286]
- **Jones, L.J. et al.** *Analytical Biochemistry*, 1998, vol. 265, 368-374 [0293]
- **Scaringe, S. A. et al.** *J. Am. Chem. Soc.*, 1998, vol. 120, 11820-11821 [0366]
- **Matteucci, M. D. ; Caruthers, M. H.** *J. Am. Chem. Soc.*, 1981, vol. 103, 3185-3191 [0366]
- **Beaucage, S. L. ; Caruthers, M. H.** *Tetrahedron Lett.*, 1981, vol. 22, 1859-1862 [0366]
- **Dahl, B. J. et al.** *Acta Chem. Scand*, 1990, vol. 44, 639-641 [0366]
- **Reddy, M. P. et al.** *Tetrahedron Lett.*, 1994, vol. 25, 4311-4314 [0366]
- **Wincott, F. et al.** *Nucleic Acids Res.*, 1995, vol. 23, 2677-2684 [0366]
- **Griffin, B. E. et al.** *Tetrahedron*, 1967, vol. 23, 2301-2313 [0366]
- **Griffin, B. E. et al.** *Tetrahedron*, 1967, vol. 23, 2315-2331 [0366]
- **Tanaka.** *Journ. Biol. Chem.*, 1993, vol. 268, 12713-12718 [0384]
- **Altschul et al.** *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0426]
- **Zhang ; Madden.** *Genome Res.*, 1997, vol. 7, 649-656 [0426]
- **Rainwater ; Kammerer.** *J. Exp. Zool.*, 1998, vol. 282, 54-61 [0439]
- **van Berge-Henegouwen et al.** *Gastroenterol.*, 1977, vol. 73, 300 [0508]
- **DiSanto.** Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990, 1451-1458 [0508]
- **Muranishi.** *Crit. Rev. Ther. Drug Carrier Systems*, 1990, vol. 7, 1 [0512]
- **Lee et al.** *Crit. Rev. Ther. Drug Carrier Systems*, 1991, vol. 8, 91 [0512]
- **Harvey.** Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990, 711 [0518] [0520]
- **Hirahata et al.** *Gan To Kagaku Ryoho*, 1992, vol. 19 (10), 1591 [0519]
- **Somogyi et al.** *Pharm. Res.*, 1995, vol. 12, 149 [0519]
- **Akamo et al.** *Japanese J. Cancer Res.*, 1994, vol. 85, 652 [0519]
- **Inoue et al.** *Artif. Organs*, 1997, vol. 21, 28 [0519]
- **Pennington et al.** *Ailment Pharmacol. Ther.*, 1995, vol. 9, 471 [0519]
- **Benet et al.** Goodman & Gilman's The Pharmacological Basis of Therapeutics. McGraw-Hill, 1996 [0520]
- **Muranishi.** *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, vol. 7, 1 [0523] [0525] [0528]

## EP 2 336 318 A1

- **El-Hariri et al.** *J. Pharm. Pharmacol.*, 1992, vol. 44, 651 [0523]
- **Brunton et al.** Goodman & Gilman=s The Pharmacological Basis of Therapeutics. McGraw-Hill, 1996, 934-935 [0525]
- **Swinyard.** Remington=s Pharmaceutical Sciences. Mack Publishing Co, 1990, 782-783 [0525]
- **Yamashita et al.** *J. Pharm. Sci.*, 1990, vol. 79, 579 [0525]
- **Jarrett.** *J. Chromatogr.*, 1993, vol. 618, 315 [0527]
- **Muranishi.** *Critical Reviews in Therapeutic Drug Carrier Systems*, vol. 7, 1 [0527]
- **Buur et al.** *J. Control Rel.*, 1990, vol. 14, 43 [0527]
- **Yamashita et al.** *J. Pharm. Pharmacol.*, 1987, vol. 39, 621 [0528]
- **Miyao et al.** *Antisense Res. Dev.*, 1995, vol. 5, 115 [0530]
- **Takakura et al.** *Antisense & Nucl. Acid Drug Dev.*, 1996, vol. 6, 177 [0530]