

RESTRICTED USE PESTICIDE

DUE TO POTENTIAL SECONDARY TOXICITY TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Section 24(c) Special Local Need Label FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEVADA

rozol[®] VOLE BAIT

FOR CONTROL OF TOWNSEND'S GROUND SQUIRRELS (*Spermophilus townsendii*) IN FALLOW AGRICULTURAL LAND, RANGELAND, AND BORDER AREAS/BUFFER STRIPS ADJACENT TO CROPS

EPA SLN No. NV-180001

EPA Reg. No. 7173-242

This label expires and must not be distributed or used in accordance with this SLN registration after Dec. 31, 2023.

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on the EPA registered label.

DIRECTIONS FOR USE

Spot baiting: Scatter ¼ cup of bait (approximately 2 ounces) around burrows and active runways. Do not pile the bait. Make a second application 5 to 7 days after the first, following the same procedure as the first application. If squirrel activity persists, a third application may be made 5 to 7 days later. Do not exceed 10 pounds of bait per acre per application. Do not exceed 30 pounds of bait per acre per year.

Bait station baiting: Apply bait in secured tamper-resistant bait stations. Fill bait stations with 1 to 4 pounds of bait, and secure the stations near active burrows, runways, and other locations where signs of ground squirrel activity are observed. Space stations 20 to 100 feet apart, depending on the density of the squirrel population. Check stations at least weekly to ensure an uninterrupted supply of fresh bait for at least 30 days or until all signs of squirrel feeding have stopped. Do not exceed 4 pounds of bait per station.

Surveillance and Follow-Up: Collect and dispose of dead ground squirrels and search for non-target animals 4 days after the first application and at subsequent intervals of 1 to 2 days for at least 2 weeks after the last bait application, or longer if carcasses are still being found. Wearing gloves, collect and properly dispose of visible carcasses of ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Wear gloves and use leakproof plastic bags or other suitable containers for transporting carcasses not buried on site.

RESTRICTIONS / PRECAUTIONS:

- Apply this product when ground squirrels are active, approximately mid-February to mid-July.
- Do not apply this product directly to water or within 10 feet of surface water (i.e. streams, rivers, lakes).
- Do not broadcast bait. Do not apply this product by any method not specified on this label.
- Keep away from humans, domestic animals and pets during application, handling and storage.
- This product is extremely toxic to mammals and birds. This SLN label should not be used where impact on listed threatened or endangered species is likely. Consult the Federal label for additional restrictions and precautions to protect fish and wildlife.

24 (c) registrant:
Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(800)-351-1476

LIPHATECH[®]

12/14/2017